# \*Academic Year 3\*



# **Complications of Exodontia**

# Dr. Hamid Hammad Enezei

# **Ph.D in Oral & Maxillofacial Surgery**



# **Objectives:**

# **1-Understand the technique of:**

- A- Extraction of single-rooted teeth with destroyed Crown.
- **B-** Extraction of multi-rooted teeth with destroyed Crown.
- C- Root tips extraction.
- 2 List the indications and contraindications for surgical tooth extraction.
- **3** Outline the steps of surgical extraction of tooth / root.



Further reading Fragiskos D. Fragiskos (2007): Oral Surgery. Springer Robinson (2007): Tooth Extraction-A Practical Guide. Wright H<sup>2</sup>upp(2014), Contemporary Oral and Maxillofacial Surgery. Alsevier

# **Methods of Tooth Extraction**

Erupted teeth could be extracted by two ways:

# 1- The closed technique (routine)

The most frequent technique.



# 2- The open technique (flap or surgical)

It is indicated when the routine method fails to extract the tooth.

# **Clinical Steps of the Closed Extraction Technique**









4- Buccal and lingual pressure, with tractional force buccally



## **Extraction of Single Rooted Teeth with Destroyed Crown**

**1-** It is accomplished with the help of the straight elevator.

2- The blade of the elevator is seated between the root and alveolar bone, with the concave surface of the blade is in contact with the mesiobuccal or distobuccal corner of the root.

**3-** Using the alveolar bone as a fulcrum, rotational forces are applied around the axis of the elevator, resulting in displacement of the root out of the socket.

### **Indication:**

When the root could not be removed by root forceps & the shape is favorable, and its level is above or with the level of the bone. 6



Luxation of root with straight elevator



## **Extraction of Single Rooted Teeth with Destroyed Crown**

4- Sometimes a small portion of the alveolar bone may be grasped with the root forceps and the root then is removed along with that small portion of the alveolar bone ( alveolar application of the forceps ).

### Note:

If you are planning to place an implant, this procedure is not recommended since the bone is very precious in implant surgery.

### **Indication:**

When the root could not be grasped by root forceps & the shape is favorable, and its level is with or little below the alveolar bone.



## **Extraction of Multi Rooted Teeth with Destroyed Crown**

**1-** It is achieved by root separation.





Mandibular first molar with destroyed crown

2- The roots are sectioned and separated after creating a deep perpendicular buccolingual groove using a fissure bur, which reaches the intraradicular bone.

### **Indication:**

When you can not get grasp by forceps & the roots shape is favorable, and its level is above or with the level of the bone.





Separation of roots of the mandibular first molar with fissure bur

**3-** Sectioning may also be accomplished using the straight elevator, after placing its blade in the root bifurcation.





**4-** The roots then separated by employing rotational movements.



**5-** At the same time by this rotational movement, one root could be elevated easily from its socket, otherwise root forceps could be used to remove each root separately.

a







**6-** The other root may then be removed using the cryer's elevator.

7- The elevator is placed so that the blade is positioned in the empty socket with its end facing the root. The intraradicular bone is removed first (if it is higher than the root) and then the tip of the elevator comes into contact with the root, which is removed after applying rotational pressure upwards.

### Note:

In this example, Straight elevator could be used by placing it in the mesial corner of the root and pushing it toward distal socket.









8- For upper molars ( have three roots ), the roots are sectioned, first separate palatal from buccal roots, then section the buccal roots into mesiobuccal and distobuccal one. Sectioning either by bur or elevator. Then each root is removed separately by elevator or root forceps.

### Note:

During sectioning of the roots by bur, attention should be given not to extend the bur deep since maxillary sinus sometimes extends into the furcation area of the tooth.



**Roots sectioning** 





## **Extraction of Root Tips**

1-In order to extract root tips from the maxilla and mandible, fine apexo elevator (with narrow blade) is the most appropriate instrument, due to its sharp tip, which fits easily between the root tip and the alveolar bone, luxating the root tip from the socket.

2-Sometimes extraction of the root tip requires part of the intraradicular bone should be removed from inside the socket with a round bur or sharp instrument, creating room that will facilitate its luxation.

**3-** Special care required for extraction of root tips of upper posterior teeth because there is an increased risk of displacing the root tip into the maxillary sinus.



4- The root tip may also be removed with the aid of an endodontic file, which is positioned inside the socket and then screwed into the root canal, upon which the root tip is delivered either by hand or with a needle holder.

When a needle holder is used, a protective gauze may be placed between it and the occlusal surface of the teeth on which it rests.

#### Note:

This procedure is difficult to achieve unless we get primary mobility of the root before it gets fracture with good visibility of the field and the root has visible root canal.

### **5- Irrigation – Suction Technique:**

This technique is useful when the tooth was well luxated and mobile before the root was fractured. The socket is irrigated vigorously and suctioned with fine suction tip. 13





Endodontic file enters the root canal

The root tip is drawn upwards by hand or with a needle holder

# **Surgical Extraction of Tooth / Root**

### **Indications:**

**1-** Teeth with unusual root morphology. Because their removal with the conventional technique will result in complications (e.g. root fracture or alveolar bone fracture).

2- Teeth with hypercementosis of root or with large bulbous roots.

**3-** Teeth with dilaceration of root tips.

**4-** Teeth with ankylosed roots or with abnormalities, e.g., dens in dente.







maxillary premolar with hypercementosis at root tip





Dens in dente of maxillary canine

5- Impacted and semi-impacted teeth.

5-Fused teeth. If extraction done by conventional technique, then part of the alveolar process could be fractured or removed together with the teeth.

7- Maxillary posterior teeth, whose roots are divergent and included in the maxillary sinus or in case of pneumatization of the maxillary sinus ( the maxillary sinus extends as far as the alveolar ridge).





Semi-impacted mandibular third molar



P<sub>1</sub>n<sub>5</sub>eumatization of the maxillary sinus into the alveolar ridge



Roots of upper posterior teeth in close relation with the sinus

**p** fusion of two mandibular premolars



Fusion of maxillary second molar with the third molar in the apical area

8-Broken root tip that have remained in the alveolar bone specially if it is associated with osteolytic lesions, or in such a position that, they could create problems for future prosthesis.

9-Roots with periapical lesions, whose entire removal (for the lesion) through the tooth socket would not be possible with curettage alone.

**10-** Deciduous molars whose roots embrace the crown of the subjacent premolar. If the conventional extraction technique were to be attempted, there is a great risk of extraction of the premolar



16 Deciduous molar, whose roots embrace the crown of the succedaneous premolar



Roots of maxillary and mandibular molars completely covered by bone



**Roots with large periapical lesions** 

# **Contraindications**

Asymptomatic fractured root tips Not associated with periapical pathology Located deep in the socket Small size root tip ( not more than 3-4 mm )

This extraction should not be done especially in cases when: **1** Dealing with older patients.

2There is a risk of serious local complications, such as the dislodging of a root tip into the maxillary sinus or injury of the inferior alveolar nerve, mental nerve, or lingual nerve.

**3–** A large part of the alveolar process needs to be removed.

4– There are serious health problems present.

## **Steps of Surgical Extraction**





### **Three techniques:**

- A- Removal of part of the buccal bone for luxation of the root buccally.
- B- Opening of a window on the buccal bone for removal of the root through the socket or through the window itself.
- C- Creation of a groove on the surface of the root or between the root and bone which allows positioning of the elevator.

### Extraction of the tooth or root with elevators orforceps

#### Note:

Any bone removal by handpiece should be coupled with good irrigation by normal saline.

Postoperative care of wound and suturing



# Surgical Extraction of Teeth

# **Surgical Extraction of Teeth**

**1-** Flap is made ( envelope, 2 or 3 sided ).

**2-** The bone covering the buccal surface of the root (s) is then removed (just below the root bifurcation in multi rooted).

### A- In single rooted teeth:

The extraction is performed easily towards the buccal side that is no longer covered by bone, using forceps or elevators.

### **B- In double rooted teeth:**

The roots are separated, after a vertical groove is created on the crown using a fissure bur, which extends as far as the intraradicular bone, then each root is removed separately.

### C- In multi rooted teeth:

The two buccal roots are sectioned using a fissure bur, the crown together with the palatal root is removed, then the two buccal roots are removed separately, preferably using a straight elevator or root forceps.

**3-** The flap is replaced and sutured. 20



**3- Multi-Rooted Teeth (Maxillary Molars)** 



# Surgical Extraction of Root

## **Steps of Surgical Extraction**



### **Extraction of the root with elevators or forceps**

### Note:

22

Excessive removal of bone should be avoided especially in a patient who is planned for implants.

Postoperative care of wound and suturing



## **Surgical Extraction of Root**

A- After removal of part of the buccal bone

**a- Single rooted1-** Flap is made ( 2 or 3 sided ).



**2-** The mucoperiosteal flap is reflected using periosteal elevator.

**3-** Part of the buccal bone is removed using a round bur until the root is exposed.



**Bone removal** 

**Root exposure** 

**4-** The root is then luxated using a straight elevator and removed by applying a small amount of pressure outwards.





After extraction , the 5surgical site should be inspected, all bony spicules should be removed, any sharp edges should be bony smoothened, sharp areas of bone assessed by are replacing the flap and palpating it with finger. Rongeur or a bone file may be used to smooth these areas.



6- The surgical field is irrigated with saline solution especially at the area of base of the flap as debris tend to collect in this area.

**Root removal** 

**Bone smoothening** 



**7-** Repositioning of the flap and the wound is sutured (simple interrupted suture).





### **B- Multi rooted:**

1-Flap is made ( envelope flap or 2 sided ), extending one or two teeth, beyond the root to be removed.

2-Then part of the buccal bone is removed using a round bur, until the root bifurcation is exposed.

**3-** The roots are sectioned using a fissure bur and are removed with a straight elevator.

**4-** The socket is then cared for and sutures are placed.











## **Steps of Surgical Extraction**



### **Extraction of the root with elevators or forceps**

Postoperative care of wound and suturing



# **Surgical Extraction of Root B- After a window is created on buccal bone**

In this technique the removal of the root is achieved either through the socket itself or through the window which is created.

# a- Root removal through the socket:

**1-** Flap is made -2 or 3 sided flap.







**2-** Reflection of the mucoperiosteal flap to expose the labial or buccal bone.

**3-** A small window is created, using a round bur, on the labia/buccal bone corresponding to the tip of the fractured root.

**4-** The window is then enlarged, and good amount of the root structure is exposed to allow its displacement from the socket using an angled elevator (like apexo or curved warwick james).



This technique is indicated for the removal of roots immediately after their fracture (the socket still open).







b- Root removal through the window which is created:

**1-** A semilunar flap is created.

**2-** The mucoperiosteal flap is reflected to expose the buccal bone.









**3-** A small window is created, using a round bur, on the labial/buccal bone corresponding to the tip of the fractured root.





4-The root is then removed from the window without difficulty, preferably using a an angled elevator.

5-After removal of the root, the socket is cared for and interrupted sutures are placed.

This technique is usually used in cases of fractured small roots, which were not removed during the extraction procedure but remained in the socket for a long time and were eventually totally covered by bone.









## **Steps of Surgical Extraction**



### Extraction of the root with elevators or forceps

Postoperative care of wound and suturing



## **Surgical Extraction of Root**

C- By creation of groove on surface of root or between root and bone, which allows positioning of the elevator.

1-Flap is made - envelope or 2 sided flap.

### 2 Then either:

**a-** A small amount of buccal bone is removed, until part of the root is exposed, a groove is then created on root surface, which allows positioning of the blade of the elevator, to luxate the root outwards.

**b-** A groove is created using a round bur between the buccal bone and the root, which makes enough room to allow for the positioning of the elevator, the blade of the elevator (straight or cryer's elevators) is then seated in the groove, which luxates the root upwards.

This technique is used in the lower posterior teeth, where the buccal bone serves as a fulcrum, is dense and able to withstand applied pressure.



Creation of groove on surface of root



Creation of groove between root and bone

# **Summary of Surgical Extraction of Root**





# **Thank You**

# \*Academic Year 3\*



# Extraction of Teeth (Exodontia) Part 2 Dr. Hamid Hammad Enezei

# Ph.D in Oral & Maxillofacial Surgery


### **Objectives:**

List the indications for teeth extraction.

List the contraindication for teeth extraction.

Identify the main points in clinical evaluation of patients for extraction.

Identify the main points in radiographic evaluation of patients for extraction.



**Further reading** 

Fragiskos D. Fragiskos (2007): Oral Surgery. Springer Hupp (2013), Contemporary Oral and Maxillofacial Surgery. Alsevier 2

### RELATIVE

If the contraindication is provided with additional care one can overcome the complication.

In other words, given the situation, the patient is made fit to undergo extraction once the underlying condition is treated.

### ABSOLUTE

There are few conditions which are absolute contraindications.

These factors will be impediment for extraction even if care is taken.

If extraction is carried out in presence of such absolute contra indications, the outcome may be fatal.

- 1. Local
- Irradiated Tissue
- Extractions performed in an area of radiation may result osteoradionecrosis.
- Adjacent Neoplasm
- Teeth that are located within an area of tumor, especially a malignant tumor, should not be extracted.
- The surgical procedure of extraction could disseminate malignant cells and thereby seed metastasis.

### Pericoronitis

- Patients who have severe pericoronitis around an impacted mandibular third molar should not have the tooth extracted until the pericoronitis has been treated.
- Non surgical treatment should include irrigations, antibiotics and removal of maxillary third molar.

### 2. SYSTEMIC

### □ DIABETES AND HYPERTENSION

• One should investigate the state of these disorders in every patient and extraction should be carried out only after confirming that they are under control.

### PATIENTS ON STEROID THERAPY

- If patient gives history of cortisone therapy, the, dental surgeon has to take certain precautions.
- A physician's opinion must be taken.

# 3. Pregnancy

- The clinician should bear in mind the existence of the possibility of obstetric complications during the first and last trimester.
- Hence if possible, extraction can be carried out after obtaining the obstetrician's expert opinion.

### 4. Bleeding Disorders

- The patients who give a definite history of bleeding episodes need careful evaluation
- Patients with anticoagulant therapy can undergo extraction after obtaining prior advice from the patient's physician/cardiologist.

### 5. Active Infections

- These are relative contra indications.
- For example: Extraction in the presence of active and uncontrolled infection will lead to the regional or systemic spread.
- Hence, it is preferable to control the infection and extraction can be safely carried out under the umbrella of antibiotic therapy.

### 6. Extraction Of Teeth In Recently Irradiated Patients

- These cases deserve special mention.
- Irradiation of jaws reduces blood supply due to fibrosis.

### Osteoradionecrosis

➤ is an extremely serious complication for patients requiring tooth extraction after radiation therapy, and the risk does not appear to diminish with time (for the rest of his/her life the patient should never allow a dentist to extract a tooth after radiation therapy without consulting a radiation

### **Complications after Radiotherapy**



### Dental Extraction after Radiotherapy :

- Delayed wound healing
- ➢ Prolonged alveolar bone exposure
- ➤ Infection
- > Osteoradionecrosis









### Radiation-related caries



# *Mostly temporary* (teeth can be removed after treatment of the pathology)

# **Bleeding disorders**





# immunesuppression





# For a period of 6 months after MI





# Acute leucocytosis or leucopenia







## **Uncontrolled diabetes**





# Bisphophonates and denosumab (RANKL inhibitors)



Local contraindications

Mucositis during chemotherapy





# Local contraindications

# Presence of a pathology in close proximity



Local contraindications

Presence of radiation mucositis



# Local contraindications

# Arteriovenous malformation



# Local contraindications

# Acute pericoronitis (severe)



# Local contraindications

# Acute odontogenic abcess (contraindicaton)





### Pregnancy

- first and last trimester .. NO (take care please)..never do it
- Middle trimester.. YES.. Can
- Obtaining the obstetrician's expert opinion.



Contra-indications for an **extraction** are usually temporary and concern the general medical suitability of the patient for **surgery** (bleeding tendency, immunosuppression, condition following myocardial infarction < 6 months), or a local surgical disorder (abscess, tumor).

Contraindications:

- Absolute

- Relative

These contraindications are less frequent but in base of existence, contraindications are more important than indications.

# **Contraindications Of Extraction:**



### Absolute contraindications:

- Acute (cardiac, pulmonal, cerebral) circulatory catastrophe (e.g. heart attack, coronary thrombosis, stroke, shock)

- Haematological diseases (AML, ALL, agranulocytosis
- Previous radiotherapy of the jaws

### Relative contraindications:

Bad general condition (cachexia, anaemia)

- Increased liability to infection (immunodeficiency, immunosuppresion, steroid therapy)

- Acute infective diseases
- Acute oral infections
- Coagulopathy, anticoagulant therapy (without prep)
- Long lasting, high dose bisphosphonate therapy
- Space reservation
- Orthodontic aspects
- Prosthodontic aspect
- Aesthetic aspects
- Financial aspects





## **Thank You**

# \*Academic Year 3\*



# **Dental Forceps and Elevators Dr. Hamid Hammad Enezei**

### Ph.D in Oral & Maxillofacial Surgery



### **Objectives:**

Describe the basic instruments for exodontia Understand the component parts of the forceps and elevators Identify different types of the extraction forceps Describe the functions of the dental elevator Identify different types of elevators



Further reading Fragiskos D. Fragiskos (2007): Oral Surgery. Springer Hupp (2013), Contemporary Oral and Maxillofacial Surgery. Alsevier 2

### **Basic Instruments for Exodontia**

The conventional extraction is accomplished with the use of extraction forceps and dental elevators.

### **Extraction forceps:**

A- It consists of handle, hinge (joint), and the beaks.

B- The handle has a serrated surface to allow a positive grip & to prevent slippage.

C- The beaks are the functional component of the forceps and grasp the tooth at the cervical region.







### **Cervical grasp**

D- The beaks are designed to be parallel to the long axis of the tooth upon use, accordingly:

**Beaks of the maxillary forceps** are parallel to the handle.

**Beaks of the mandibular forceps** are perpendicular to the handle.







### Beaks parallel to the long axis of tooth



E- Because tooth anatomy varies, forceps with different beaks designs have been manufactured, so that they are used for specific teeth.










**Types of the extraction forceps** 

## **Maxillary Extraction Forceps**

#### A- For the Six Anterior Teeth of the Maxilla: ( Central, Lateral, Canine)

#### Features:

1The beaks are in the same line with the handles (straight forceps).

- 2 The beaks are broad and identical.
- 3 The ends of the beaks are open.



Straight forceps



## B- Maxillary premolars forceps: (First & Second)

#### **Features:**

1The beaks are slightly curved, to compensate for the presence of the lip and cheek and to get parallel application of the beaks with long axis of the tooth (beaks are not in a straight line with the handles).

2The beaks are broad and identical.

3- The ends of the beaks are open.









## C- Maxillary Molar Forceps: ( First and Second Molar )

#### **Features:**

1There are two designs: one for the left and one for the right side.

2 The beaks are not identical.

3The buccal beak has a projection, which fits into the buccal bifurcation of the two buccal roots, while the palatal beak is concave and fits into the convex surface of the palatal root.

**4-** The beaks are curved to compensate for the posterior position of the tooth, to avoid lip & cheek interference and to get parallel application of the beaks with long axis of the tooth









Maxillary right molar forceps

Curved beaks

## D- Maxillary Third Molar Forceps ( Bayonet forceps )

#### **Features:**

1- Longest forceps, the beaks with double curvature to compensate for the posterior position of the tooth, to avoid lip & cheek interference and to get parallel application of the beaks with long axis of the tooth



2- Because third molar tooth varies in shape and size, the beaks of the forceps are broad and identical (without projection), so that these forceps may be used for extraction of both the left and right third molar of the upper jaw.





### **E- Maxillary Anterior Root Forceps**

#### Features:

1The beaks are in the same line with the handles (straight forceps).

2 The beaks are identical.

3The beaks are narrow, pointed and closed.

4They are used for **root removal** of maxillary anterior teeth.





## **F- Maxillary Posterior Root Forceps**

#### Features:

1The beaks are slightly curve.

2 The beaks are identical.

3The beaks are narrow, pointed and closed.

4They are used for removal of **premolar roots and single molar root pieces.** 



Maxillary posterior root forceps





Maxillary premolar forceps



Maxillary anterior teeth forceps **Open- identical-straight** 





**Maxillary premolar forceps Open- identical - curved** 





**Maxillary molar forceps Open- not identical- curved** 



Maxillary third molar forceps **Open-identical-curved** 



Maxillary posterior root forceps **Closed** – identical-curved



Superior view of No. 150A forceps. **B**, No. 150A forceps have parallel beaks that do not touch in contrast to No. 150 forceps beak.

C, Adaptation of No. 150A forceps to the maxillary premolar.





Superior view of No. 1 forceps. **B** and **C**, No. 1 forceps adapted to the incisor

## **Mandibular Extraction Forceps**

They are designed in such a way that the handles and the beaks are at right angle to each other:

To facilitate easy access to mandibular teeth.

To get parallel application of beaks with the long axis of the tooth.

### A- Mandibular Anterior Teeth Forceps

#### Features:

- 1 Identical beaks
- 2 Broad beaks
- 3 Closed beaks







### **B- Mandibular Premolar Teeth Forceps**

#### Features:

- 1 Identical beaks.
- 2 Broad beaks.
- 3 Open beaks.





opened beaks-lower premolar forceps closed beaks-lower ant. forceps



## **C- Mandibular Molar Forceps**

1 Identical beaks.

- 2 Broad beaks.
- 3 Open beaks.

4Both beaks end in a pointed tip (projection), which fit into the bifurcation of the roots buccally and lingually.

**5-** These forceps are used for both sides of the jaw since lower molar has bifurcation on the buccal and lingual sides.

**6-** Two styles available, one for anterior approach, and the other for lateral approach.





**A**, Superior view of cowhorn No. 87 forceps. **B**, Side view of cowhorn forceps. **C** and **D**, Cowhorn forceps adapted to the lower molar tooth





Superior view of No. 53L forceps. **B**, Oblique view of No. 53L forceps. **C**, *Right*, No. 53L; *left*, No. 53R. **D** and **E**, No. 53L adapted to the maxillary molar.



**A**, Superior view of No. 88L forceps. **B**, Side view of No. 88L forceps. **C**, No. 88R adapted to the maxillary molar.



**A**, Superior view of No. 210S forceps. **B**, Side view of No. 210S forceps. **C**, No. 210S adapted to the maxillary molar

#### **D-** Mandibular Root Forceps.

#### Features:

- 1 Identical beaks.
- 2 Long narrow beaks.(longer than anterior or premolar forceps)
- 3 Closed beaks.
- 4 Used for removal of root of all mandibular teeth.
- **Note:** the long beaks to enable the forceps to take a deep grip on the root stamp.







Lower anterior teeth forceps Closed – identical







Lower molar forceps Open –identical with projection





**A**, Superior view of No. 65 forceps. **B**, Side view of No. 65 forceps. **C**, No. 65 adapted to broken root.



The No. 150S forceps (*bottom*) are a smaller version of the No. 150 forceps (*top*) and are used for primary teeth.



# **Dental Elevators**

1 It has three parts:

Handle.

Shank.

Blade.

2-Different designs are available for different clinical applications.







# **Dental Elevators**

#### **Functions of the elevators are:**

A- They are used to luxate teeth or roots from the surrounding bone before the application of the forceps. This Loosening makes the extraction easy, and the incidence of root fracture will be less, and if root fracture occurs during extraction, its removal will be very easy if it loosened before.





**B-** They are used to remove broken roots or surgically sectioned roots from their sockets.



**C-** They are used for roots separation.





# **Types of Dental Elevators**

#### **1- Straight Elevator:**

**A-** It is the most commonly used type of elevator for the loosening of the teeth and roots, in both the upper and lower jaws.

**B-** The handle is pear-shaped, the blade has two surfaces: a convex and a concave one. The concave surface faces the tooth to be luxated. It comes in series of sizes like no.1,2, or 3

C- Two designs available, one with rounded end (usually called straight elevator) and one with sharp, straight end ( Coupland's chisel elevator )



### **2- Cryer's Elevator:**

**A-** It is used only in the lower jaw for removal of molar's roots.

**B-** The blade is triangular in shape and pointed.

**C-** They come in pairs, one for mesial root and other for distal root.

**D-** In certain cases, may be used to remove a whole third molar of the lower jaw.

**E- Winter's elevator**, is similar to cryer's, but has T shape handle, and great force can be generated with such design.



Winter's elevator (T) shape handle





#### **3-** Apexo Elevator:

**A-** It is used to remove root tips in both jaws, also useful for extraction of impacted upper third molars.

**B-** The handle is pear shaped. The shank has a double angle. The blade ends in a sharp point. It comes in pairs for mesial & distal application.

**C-** Some designs have very narrow blades and sharp-tipped ends for very small root fragment.





Very arrow, sharp tip blade





#### 4- Warwick James elevator:

**A-** It comes in three designs straight and curved (one for mesial and other for distal side).

**B-** The handle is flat, the blade is small.

**C-** Used to luxate teeth specially maxillary third molar, and deciduous molar.













Surgical Extraction Of Teeth And Roots - ELEVATORS

# Forms of Elevators



(1)(2)(2)(2)



# **Special Instruments**

1- Instrument to separate the gum from around the tooth during extraction like **Desmotomes and Periotomes** 







**Desmotomes a Straight. b Curved** 

# 2- Instrument for removal of roots called root extractor

It is used to remove broken roots found below the alveolar crest. The spiral end of the instrument is placed inside the extraction socket, and, after screwing the instrument into the root canal of the broken root, traction is used to remove the root from the socket.









# **Thank You**







# **Ph.D in Oral & Maxillofacial Surgery**



1

# **Objectives:**

- 1- To establish a positive professional relationship-patient assessment , evaluation and diagnosis.
- 2- To provide the clinician with information concerning the patient's past dental, medical and personal history.
- 3- To provide the clinician with the information that may be necessary for making a diagnosis.
- 4- To provide information that aids the clinician in making decisions concerning the treatment of the patient.



Further reading

Fragiskos D. Fragiskos (2007): Oral Surgery. Springer Robinson (2007): Tooth Extraction-A Practical Guide. Wright Happ (2013), Contemporary Oral and Maxillofacial Surgery. Alsevier

# INTRODUCTION

To provide best treatment and patient satisfaction, thorough clinical history, examination and diagnostic aids are required Since dental problems are not same in two patients, so thorough examination, evaluation and diagnosis of an individual patient guides the effective treatment plan




- •Diagnosis is defined as utilization of scientific knowledge for identifying a diseased process and to differentiate it from other disease process
- •In other words, literal meaning of diagnosis is determination and judgment of variations from the normal



# Patient Evaluation

The diagnostic process actually consists of four steps;

- 1. First step: assemble all the available facts gathered from chief complaints, medical and dental history, diagnostic test and investigations
- 2. Second step : analyze and interpret assembled clues each and the tentative or provisional diagnosis
- 3. Third step :make differential diagnosis of all possible diseases which are consistent with signs, symptoms and test results gathered
- 4. Fourth step : select the closest possible choice

#### **DEVELOPING A SURGICAL DIAGNOSIS**

-Most of the important decisions concerning a surgical procedure should be made long before the administration of anesthesia. The decision to perform surgery should be the culmination of several diagnostic steps. *The surgeon first identifies;* 

- 1- the various signs and symptoms
- 2- and relevant historical information; then,
- 3- using available patient and scientific data
- 4- and logical reasoning based on experience, the surgeon establishes the relationship between the individual problems.

### THE DIAGNOSTIC PROCESS



For a good analysis, data must be organized into a form that allows for hypothesis testing; that is, the dentist should be able to consider a list of possible diseases and eliminate those unsupported by the patient data or

#### **BASIC NECESSITIES FOR SURGERY**

-Little difference exists between the basic necessities required for oral surgery and those required for the proper performance of other aspects of dentistry. The two principal requirements are;

- adequate visibility and assistance (1)
- (2)

-Although visibility may seem too obvious to mention as a requirement for performing surgery, clinicians often underestimate its importance, especially when the unexpected occurs. Adequate visibility depends on the following three factors:

- (1)
- adequate access, adequate light, and a surgical field free of excess blood and other fluids. (2)(3)



Adequate access not only requires the patient's ability to open the mouth widely but also may require surgically created exposure. Retraction of tissues away from the operative field provides much of the necessary access. (Proper retraction also protects tissues being retracted from being accidentally injured, for example, by cutting instruments.) Improved access is gained by the creation of surgical flaps.

- ✤ Adequate light is another obvious necessity for surgery. However, clinicians often forget that many surgical procedures place the surgeon or assistant in positions that block chair-based light sources. To correct this problem, the light source must continually be repositioned, or the surgeon or assistant must avoid obstructing the light, use more than one overhead light, or use a headlight.
- ✤ A surgical field free of fluids is also necessary for adequate visibility. High-volume suctioning with a relatively small tip can quickly remove blood and other fluids from the field.
- ✤ As in other types of dentistry, a properly trained assistant provides invaluable help during oral surgery. The assistant should be sufficiently familiar with the procedures being performed to anticipate the surgeon's needs. Performance of good surgery is extremely difficult with no or poor assistance.

#### PRESURGICAL MEDICAL ASSESSMENT

It is critical that the surgeon examine the patient's medical status. patients can have a variety of health problems that require treatment modification or medical management before the surgery can be safely performed. Special measures may be needed to control bleeding, lessen the chance of infection, or prevent a medical emergency.

# **Case Sheet-of-History**

## **Definition:**

Case history is defined as; planned, professional conversation between the patient and the clinician in which the patient reveals his/her symptoms fears, or feelings to the clinician so that the nature of the real or suspected illness and mental attitude to it may be determined

### •Purpose:

- 1. To discover whether patient has any general or local condition that might alter the normal course of treatment
- 2. comprehensive medical and previous dental history should be recorded. In addition, a description of the patient's symptoms in his or her own words should be noted
- 3. The purpose of recording patients history and conducting a clinical examination is to arrive at a logical diagnosis to the patients chief compliant and to institute a suitable treatment plan



#### **PRESENT ILLNESS-3**

#### **FAMILY HISTORY-6**

# **Personal data**

Patient name : Age: NAME HEALTH INFORMATION Sex: 7 ADDRESS 2 Marital status: INCOME **Occupation** : 6 LOCALISATION Address 3 **CULTURAL PROFILE** 5 **ONLINE IDENTIFIER** 

4

What is Personal Data?

#### Patient name :

**Better Communication** 

**Documentation** 

## Age:

Related diseases ( Herpetic gingivostomatitis, Bohn nodule LP, Malignancy)

Dose of the drugs

#### Sex:

Mixed names

**Related diseases** 

## Marital status:

SeX transmitted disease

🛛 Pregnancy ( G.enlargment, gingivitis)

**Lactation** 

## Occupation :

Work related hazards ( Gloss blower, Carpenters )

**Address** 

Idea about social background

**Endemic disease (florosis)** 

industrial hazarrds

### **Chief Complain**

Patient's own words

Do not use complicated medical terminology

Pain, Bleeding, Swelling, Esthetic problems, etc

Regular check up

- ✓ It is the description of the problems for which the patient seeks treatment
- ✓ It should be recorded in patients own words and should not be recorded in medical terminology

#### **Importance:**

Overall treatment plan revolves around the chief complaint. It consists of information which promoted patient to visit a clinician. The form of notation should be in patient's own words

<u>Symptoms</u> are phenomenon or signs of deviation from normal and are indicative of illness

#### **HİSTORY OF PRESENT ILLNESS**

#### It is the detailed description of chief complaint

#### More descriptive analysis about this initial information

Signs and symptoms Duration, intensity of pain, Relieving and exaggerating ( triggering )factors

#### The most common toothache may arise either from pulp or from PDL

Mild to moderate type of pain can be of pulpal or periodontal origin. If pain from PDL ,teeth will be sensitive to percussion Pulpal pain will be sharp and depends on the pulpal fibres involved History of present illness

#### Onset.

**Duration.** 

Type (Nature) of pain.

Severity of pain.

Location and site.

Prior occurrence.

**Exacerbating factors.** 

**Relieving factors.** 

Associated phenomenon (Fever, malaise, nausea, etc)

Previous medications.

# **Taking Pain History**

Site / Localization	Where is the pain?							
Type ( Character )	Dull, Throbbing, electrical like, stabbing							
Onset	When did the pain start, and was it sudden or gradual? Include also whether if it is progressive or regressive.							
Duration	Duration of pain or attack							
Severity	How bad is the pain? Does it affect daily activity or affect sleep							
Exacerbating factors and relieving factors.	Hot and cold							
Radiation / Associated phenomenon	Does the pain radiate anywhere? Fever, vomitiing							

- Chief complain of pain
- ✤ I have Pain in the lower right side
- ✤ I have pain on drinking cold water
- History of chief complain

A patient presents with **one week** history of prolonged **throbbing** pain **localized** to the **lower molar region**. Pain is **waken** the patient from sleep, which is **aggravating by cold drinks**, and partially **relived by taking an analegesic** 

## **Taking History of bleeding**

Site / Localization	Where is the bleeding?
Onset	When did the bleeding start ?
Duration of bleeding	Duration
Spontaneous / on tooth brushing/ on eating	Hot and cold
How does it stop?	Afetr tooth brush

# Chief complain

"My gums are bleeding when I brush my teeth."

# History of chief complain

A patient presents with <u>two weeks</u> history of gingival bleeding <u>localized</u> to the <u>lower anterior region</u>. which is <u>initiated by brushing</u>, and last for few minutes

# Past dental history

- Frequency of visiting dentist
- Past caries / restorative experience
- 🏵 L.A
- Causes of loss teeth (perio/ caries)

# Past dental history

- Complications
- ✓ With L.A
- History of difficult extraction

Any post op complications ( bleeding, Pain, delayed healing)

\* Patient compliance

## Past medical history

#### Any medical problems

- ✓ Diseases/ drugs may complicate dental work
- Dental work may adversely affect the disease
- Drug interaction/ adjustment/ adverse reaction
- ✓ Oral manifestation of systemic diseases or drugs
- Hospitalization (Blood transfusion)

Allergy (Drugs / food etc ) (Anaphylactic shock )

# Systemic review

Signs and symptoms

To refer patient with undiagnosed diseases

# Systemic review

- \* CVS (Chest pain, difficulty in breathing, palpitation, clubbing finger)
- \* Respiratory ( Cough, wheeze, difficulty in breathing )
- GIT (Diarreha, constipation, difficulty in swallowing, vomiting, jaundice, bleeding)
- \* Endocrine system ( Polyurea, thirst, polyphagia, weight loss, hair loss, heat intolerance)
- \* Genitourinary system (Burring on urination, blood with urination)
- Hematopietic system (Fatique, brittle nail, eccyhmosis, bruising, gingival enlargment, )
- Central nervous system ( Seizure, numbness, confused, disoriented)

# Social history / Habit

- Smoke, Alcohol,
- Teeth brushing
- Other Para functional habit like thumb sucking, nail biting,

# **Family history**

- Hereditary and familial disorders
- Diabetes mellitus, HTN, Hemophilia, Allergy

# **Clinical Examination**

- A. Extra- oral examination
- B. Intra-oral examination

# Vital signs

- Temperature
- Pulse rate ( 60-100 beats per minute )
- Respiratory rate (16-18 breath per minute)
- Blood pressure (120/80 mm hg)

## **Extra-oral examination**

- Patient build/ Gait : (general appearance)
- Facial symmetry: ( infection, tumor )
- Skin: ( pallor, jaundice, cyanotic )
- Hair and Nail (brittle nail, jaundice, clubbing finger)
- Eyes: (Yellowish color, Pallor, redness)
- Nose: ( deviation, saddle nose )
- Æ Ears:

# **Extra-oral examination**

- Lymph node
   Lymph node
- Cranial nerve (7<sup>th</sup>)
- Salivary glands
- ✤ T.M.J
- Maxillary sinus

# The patient's outward appearance and movement

Examples Lethargic, dirty clothing and hair; body odors extreme thinness or obesity; bent posture; and difficulty breathing



## **General Appearance**

#### 

Cyanosis indicate cardiac or pulmonary insufficiency,

yellowing (jaundice) indicate liver disease,

 petechial or ecchymoses indicate bleeding disorder/ truama.







### **D**Nail

Clubbing seen in cardiopulmonary insufficiency

Yellowish indicate jaundice

 splinter hemorrhages from infective endocarditis

Pale nail bed indicate anemia



The face
Acromegaly
Moon facies in Cushing syndrome
Unilateral paralysis of Bell's palsy



### □The Eyes

- ◆ Exophthalamus indicate
- ◆ Yellow sclera may be caused by liver disease.
- ♦ Pale conjuctiva indicate Anemia
- ◆ Blue sclera indicate Osteogenesis imperfecta





# Intra-oral examination

- Oral hygiene: Poor Fair Good Excellen
   A. Soft Tissue : ( Color, size, shape, texture )
- Lips.
- Labial and Buccal mucosa.
- Palate.
- Gingiva.
- Floor of the mouth
- Tongue:

#### **B. Hard Tissue :**

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

- \* Differential Diagnosis:
- Special Investigation :
  - ✓ Vitality test
  - ✓ X-Ray:
  - Laboratory Investigation.
  - ✓ Biopsy:
- Final Diagnosis:
- & Treatment :

## Complete examination of the oral cavity includes:

- Lips and Labial Mucosa
- Buccal Mucosa
- Mucobuccal Fold
- Hard Palate
- Soft Palate and Uvula
- Oropharynx & Nasopharynx
- •Tongue
- •Floor of Mouth
- •Muscles of Mastication
- •Periodontium
- •Teeth

# THE DIAGNOSTIC PROCESS Examination of Oral Mucous Membrane



# THE DIAGNOSTIC PROCESS Examination of Oral Mucous Membrane



# THE DIAGNOSTIC PROCESS Examination of Oral Mucous Membrane


## Marking the intensity of pain



## Patient is asked to mark the imaginary ruler with grading ranging from 0 to 10 <u>0-No pain 10-Most painful</u>

•Mild to moderate pain can be of pulpal or periodontal origin but acute pain is commonly a reliable sign that pain is of pulpal origin.



## **Thank You**

## \*Academic Year 3\*



## **Extraction of Teeth (Exodontia)**

## Dr. Hamid Hammad Enezei

## Ph.D in Oral & Maxillofacial Surgery



## **Objectives:**

List the indications for teeth extraction.

List the contraindication for teeth extraction.

Identify the main points in clinical evaluation of patients for extraction.

Identify the main points in radiographic evaluation of patients for extraction.



**Further reading** 

Fragiskos D. Fragiskos (2007): Oral Surgery. Springer Hupp (2013), Contemporary Oral and Maxillofacial Surgery. Alsevier 2

## Exodontia

## (Extraction of Teeth)

## **Definition:**

-It is the painless removal of the tooth from the socket.

-It combines the principles of surgery with physical mechanics.

-Accordingly when the principles applied correctly, teeth could be removed from the alveolar process without great strength and without untoward force.



## **Indications for teeth extraction**



### **1- Gross caries:**

Severely carious, nonrestorable tooth.





### 2- Pulpal necrosis/apical pathology:

That could not be treated by conventional endodontics (like curved canal or calcified canal – may need surgery) or failed endodontic treatment and the patient refuse retreatment.



### **3- Periodontal disease:**

Advanced PDD with severe bone loss and great tooth mobility.



### 4 Orthodontic reason:

Correction of a crowded dentition frequently requires tooth extraction to create a space for teeth alignment.

### 5 Malposed teeth:

A common example is an upper third molar which erupt in severe buccal position causing ulceration and trauma to the cheek.

### 6- Prosthodontic reason:

Like over erupted tooth due to loss of tooth in the apposing arch, this will interfere with the construction of adequate prosthesis.

### 7- Cracked teeth:

Tooth that has cracked crown with root fracture that is unmanageable by  $res_5$  torative procedure should be considered for extraction.













### 8- Impacted teeth:

Fully impacted or partially erupted teeth should be considered for removal.

### 9- Supernumerary teeth:

Usually they are impacted and should be removed since it may interfere with the eruption of the permanent teeth or it may cause their resorption or displacement.

## **10-** Teeth associated with pathologic lesions:

Such a teeth should be removed, but sometimes could be retained and root filled. If maintaining the tooth compromises the complete surgical removal of the lesion, then the tooth should be removed.





## **11** For prophylactic reasons:

Some patients with a medical problems may require teeth removal prophylactically like patients need organ transplant, radiation therapy, cardiac surgery.





### **12** Teeth involved with jaw fracture:

Patients with fracture of the mandible or alveolar process may require teeth removal that are located in the line of the fracture (especially if it is severely damaged, infected or luxated ) since it may interfere with proper management of the fracture.



### 13-Patient attitude & Financial issues:

All the indications for the extractions may become stronger if the patient is unwilling or financially unable to support the7decision to maintain the tooth.











## Orthodontic extraction





## Supernumerary teeth



## retained deciduous teeth

## **Unrestorable teeth**







## Tooth interfering with prosthetic crowns



## **Periodontally weak**



## **Prior to radiation**



## **Contraindications For Teeth Extraction**

A-Systemic Contraindication
1 Uncontrolled metabolic disease:
ex: Diabetes
End stage renal disease
2 Uncontrolled leukemia and lymphoma:
<b>Potential complications: Infection</b>
Bleeding
3 Cardiac diseases:
ex: Unstable angina pectoris
<b>Recent</b> myocardial infarction
4- Pregnancy- relative contraindication.
5- Coagulation disorders:
ex: Hemophilia
Platelet disorders
6- Medications:
ex: Corticosteroid
Immunosuppressive agents

8 Cancer chemotherapeutic agents

## **B-** Local Contraindications

- 1 History of radiotherapy for head & neck cancer: potential complication: Osteoradionecrosis
- 2 Teeth that are located within an area of tumor: potential complication: Disseminate malignant

cells (metastases)

3Acute pericoronitis around partially erupted third molar:

More postoperative complications

4- Acute dentoalveolar abscess

Patient may be unable to open widely

Difficult to get profound anesthesia

### **Clinical Note:**

The most rapid way for resolution of dental infection obtained when the tooth is removed as early as possible, therefore acute infection is not a contraindication to extraction once the access and anesthesia can be ensured.

## **Clinical Assessment of Teeth for Removal**

## A- Access to the tooth (mouth opening)

Limitation of mouth opening (trismus) may compromise the extraction. Trismus may be due to:

(1)Infection around or in the muscles of mastication.

(2) TMJ dysfunction or ankylosis.

(3) Muscle fibrosis.

## **B-** Location & position of the tooth

Anteriorly located tooth better access than posteriorly located.

Well aligned tooth in the dental arch has normal access for instrument placement than crowded or malposed teeth.

## **C- Mobility of the tooth**

Greater mobility is seen in severe PD disease, accordingly extraction is simple.













### **D-** Condition of the crown

1Presence of big carious lesion or large restoration cause weakening of the crown and the possibility of crown or filling fracture during extraction is high.



2Endodontically treated teeth become dry and brittle so possibility of the fracture is high.

3Accumulation of large calculus around a tooth interferes with the proper placement of the forceps and may settle into the socket after extraction so it should be remover before the extraction.





### **E-** Condition of the adjacent teeth

If the adjacent teeth have a crown or large restoration or endodontically treated, great care should be given during the extraction to avoid fracture or displacement of the restoration





### F- Mobility of the tooth

Greater mobility = simple extraction.

## **G-** Examine if there is pain by apical palpation (pressure):

Palpate the apical area of the tooth by fingertip with slight pressure to explore if there is pain upon palpation. Presence of pain indicates there is possibility of apical pathosis while no pain does not exclude apical pathology unless confirmed by x-ray.



### H- Vitality of the tooth:

By hot / cold application or electric pulp testing

## I- Tooth Tenderness:

By slight pressure with end of the mirror to detect if there is any apical involvement of the tooth ( beginning of spread of infection from the pulp or actual infection in the apical area)









### J- Presence of apical swelling or sinus tract:

Indicates apical pathology, at the beginning there will be vestibular swelling, later on it will open and changed into sinus tract for drainage. Sometime gutta percha point inserted into the sinus tract and x-ray taken to verify the source of infection.



## **Radiographic Assessment of Teeth for Removal**

### A- Tooth morphology

### **1- Number of the roots:**

Majority of teeth have typical no. of roots, but sometimes extra root present making the extraction more difficult, in such case modification of the treatment plan can be made to prevent fracture of the additional root.

### 2- Root curvature & divergence:

More curvature –more possibility of the fracture

More divergence makes the total root width so wide that makes the extraction more difficult.

### **3- Size and shape of the root:**

Short: simple extraction

Long : difficult extraction

Conical shape: simple extraction

Bulbous (hypercementosis): difficult extr1a3ction













### 4- Root caries or resorption:

Caries extension into the root or if there is internal or external root resorption weakens the root and makes it more liable to fracture.

## 5- Presence or absence of RCT:

Presence of RCT for many years makes the tooth more brittle or ankylosed and this makes the extraction more difficult.

### 6-PDL space:

The more the space, the simple the extraction will be.









## B- Relationship to vital structure

## **1- Maxillary sinus:**

The roots of the maxillary premolars and molars are closely related to the floor of the maxillary sinus, if thin layer of bone exists between the roots and sinus or there is periapical pathology, the potential for sinus perforation during extraction increases.

## 2 Inferior dental canal:

This point is more important when dealing with impacted mandibular molar than other erupted teeth.

## 3 Mental foramen:

This foramen is closely related to the apices of the lower premolars, and should be taken into consideration when dealing with surgical extraction of mandibular premolars.











## **C-** Condition of the surrounding bone:

### **1- Bone density:**

**A-** Bone that is more radiolucent is likely to be less dense which make the extraction more easy.

**B-** Bone that is more opaque is likely to be more dens which makes the extraction more difficult.



## 2- Presence of apical pathology:

Apical pathology like cyst or granuloma should not be left after extraction, but should be removed at the same time of teeth removal.







## **Thank You**

## \*Academic Year 3\*



## Dental Forceps, Elevators and Surgical Instruments

## Dr. Hamid Hammad Enezei

Ph.D in Oral & Maxillofacial Surgery



## **Objectives:**

Describe the basic instruments for exodontia Identify different types of the extraction forceps Describe the types and functions of the dental elevator Identify different types surgical instruments in dentistry



**Further reading** 

Fragiskos D. Fragiskos (2007): Oral Surgery. Springer Hupp (2013), Contemporary Oral and Maxillofacial Surgery. Alsevier 2



EXTRACTION FORCEPSES

ELEVATORS

SURGICAL INSRUMENTS

## EXTRACTION FORCEPS

## **Extraction Forcepses**



## **EXTRACTION FORCEPSES**



# UPPER Forcepses

## 1 - FORCEPS FOR UPPER ANTERIOR





Use: Extraction of upper anterior teeth

## FORCEPS FOR UPPER ANTERIOR



## 2- FORCEPS FOR UPPER PREMOLARS





Use: Extraction of upper premolars

## FORCEPS FOR UPPER PREMOLARS



## **3- FORCEPS FOR UPPER MOLARS**









Use: Extraction of upper molars (right side Or left side)

## FORCEPS FOR UPPER MOLARS



## FORCEPS FOR UPPER MOLARS




# 4- FORCEPS FOR UPPER REMANING ROOT BAYONET



S 2-

#### FORCEPS FOR UPPER REMANING ROOT



## **REED'S FORCEPS**

### **5- FORCEPS FOR UPPERTHIRD MOLAR**



## FORCEPSES FOR LOWER TEETH





#### **1- LOWER ANTERIOR FORCEPS**



#### LOWER ANTERIOR FORCEPS

## ERIOR FORCEPS



#### **2- FORCEPS FOR LOWER PREMOLAR**



## FORCEPS FOR LO WER PREMOLAR



## 3- FORCEPS FOR LOWER MOLAR



#### Use: Extraction of lower molar teeth

## FORCEPS FOR LOWER MOLAR



## 4- FORCEPS FOR LOWER REMAINING ROOTS



**Use: Extraction of lower remaining roots** 

## **ELEVATORS**



## **STRAIGHT ELEVATORS**





## **1-STRAIGHT ELEVATOR**



## **STRAIGHT ELEVATOR**

*Trinciple:* Wedging & lever



#### USES:

- 1 luxate a mesio-angular
  - impacted lower third molar with distally curved roots. fulcrum is interdental bone
- 2 mult. ext. of neighboring teeth





Use : luxation of R.R in accessible areas

## **CURVED APEXO**



#### Use : luxation of R.R in inaccessible areas





#### Use :removal of R.R in multi-rooted teeth





#### Use : luxation of upper third molars



#### CROSS BAR SOCKET APPLICATOR



#### Use :removal of R.R in mandibular multi-rooted teeth

#### **CROSS BAR** BUCCAL APPLICATOR



Use :Luxation of vertically impacted lower third molar





## SURGICAL INSTRUMENTS



#### BARD-BARKER BLADE HANDLE AND BARD BARKER BLADES



## PERIOSTEAL ELEVATOR

## Use: reflection of flap



Use: forceful mouth opening



## **TONGUE DEPRESSOR**



## **TONGUE HOLDER**







Use: retraction of flaps or cheek









## **MALLET AND CHISELS**

- × BEVELED CHISEL
  - + UNIBEVELED removal of bone in flaks
  - + BIBEVELED (OSTEOTOME):
    - tooth division
    - Condylotomy
      - Re-fracturing malunion bone

3 H2K 566-03 PH

## **BONE RONGEURS**

### Use: bone cutting forceps









## **ARTERY FORCEPS**

#### CURVED

× MOSQUITO HEMOSTAT

## Use: ligation of blood vessels and blunt dissection during I&D
# **TISSUE FORCEPS**





Use: handling normal soft tissues

# **ALLIS FORCEPS**



Use: handling pathologic all soft tissues

# **BONE CURRETTE**

# Use: 1-bone curett 2-cyst remova







Use: grasping of firm tissues(bone-ligamentmuscles)

# **BONE FILE** Use: smoothening of sharp bony edges







# SUTURING INSTRUMENTS



# **SUTURING INSTRUMENTS**

### × NEEDLE HOLDER

# SUTURING INSTRUMENTS

### × NEEDLE









### × INTERRUPTED SUTURE



# × CONTINOUS SUTURES



# × CONTINOUS LOCKED SUTURE BLANKET SUTURE



× MATTRESS SUTURE VERTICAL





# × MATTRESS SUTURE HORIZONTAL



# **Thank You**

\*Academic Year 3\*

# **Principles of Exodontia Dr. Hamid Hammad Enezei** Ph.D in Oral & Maxillofacial Surgery





### **Objectives:**

- **1-** Understand mechanical principles involved in tooth extraction **2-** Outline the principles of forceps use
- **3-Outline the basic rules for using elevators**
- **4-Understand fundamental requirements for a perfect extraction**
- **5-** Identify the steps of tooth extraction



Further reading

Fragiskos D. Fragiskos (2007): Oral Surgery. Springer Robinson (2007): Tooth Extraction- A Practical Guide. Wright Hupp (2013), Contemporary Oral and Maxillofacial Surgery. Alsevier Methods of extraction

#### Elevators/or Forceps extraction or Closed extraction



Trans alveolar extraction or Open or surgical extraction

### Types of exodontia





# "You have to see well what you do in order to do well what you see" G.C Ingham



### **Mechanical Principles Involved in Tooth Extraction using elevators**

Three principles involved:

- 1- Lever action.
- 2 Wedge action.
- **3** Wheel & axle action.

### **1- Lever Action:**

Elevators are used primarily as lever.





# **Lever Principle**



**Example - Straight elevator (coupland elevator)** 

# Wedge Principle



E.g. : Apexo Elevator , Warwick James I E



**Example : Cross bar elevator , Cryer elevator** 

### 2- Wedge Action:

**A-** With the forceps, when the beaks of the forceps forced into the periodontal ligament space, act as a wedge causing expansion of the bony crest and force the tooth out of the socket.



Wedge expands, splits, and displaces portion of substance that receive it.

**B-** With the straight elevator when used to luxate a tooth, we force it into the periodontal ligament space, so it displaces the root out of the socket.





### **Mechanical Principles Involved in Tooth Extraction**

#### **3- Wheel & AxleAction:**

**A-** This action is clearly identified with the Cryer elevator, when one root of the multi rooted teeth is left in the alveolar process.

**B-** The handle of the Cryer serves as an axle while the tip of the blade act as a wheel so engages and elevates the root from the socket.









# **Dental Elevators**

□ Elevators are single bladed instruments designed to loosen or remove roots, root fragments, or teeth , which they do by moving them out of the sockets along a path determined by natural curvature of the roots

### Rules of using an elevator

- □ Never use adjacent tooth as fulcrum
- □ Never use lingual plate as fulcrum
- Never use buccal plate as fulcrum except where odontectomy has been performed
- □ Always use finger guards to protect the patient







# Hazard of using dental elevators

- Damaging or extracting adjacent teeth
- □ Fracturing maxilla or mandible
- □ Fracturing the alveolar process
- □ Accidental penetration of maxillary antrum and soft tissues (may perforate greater blood vessels)
- □ Forcing root or a third molar into antrum
- □ Forcing apical third of root of man.molar into man.canal,or lingual plate or pterygomandibular space







### **Principles of Forceps Use**

The forceps apply five major motions to luxate the teeth and expand the bony socket:

#### **1-** Apical pressure:

By inserting the beaks deep into the PDL space, two goals will be achieved:

A- Bony expansion

(wedging action).

**B-** Center of rotation of the tooth (fulcrum) displaced apically, and this decreases the chance of apical root fracture.



center of rotation

Strong apical pressure by the forceps – bony expansion & center of rotation more apically



Center of rotation more occlusal results in fracture of root apex



Center of rotation more apical results in less apical pressure, but great expansion of bony cortex

### **2-** Buccal pressure:

By moving the forceps buccally, it results in

A- Expansion of the buccal plate especially at the crest of the ridge.

**B-** Leads to apical pressure (toward lingual side).

C- Excessive buccal pressure can fracture buccal bone or cause fracture of the apical portion of the root.

#### Lingual or palatal pressure: 3

A- Expansion of lingual plate especially at the crest of the ridge.

**B-** Leads to apical pressure (toward buccal side).

C- Excessive lingual pressure can fracture lingual bone or cause fracture of the apical portion of the root.

#### **4** Rotational pressure:

It results in internal expansion of the tooth socket. It is suitable for a single rooted or conical shape root but multi rooted teeth or curved root are more likely to fracture by this technique.

#### **Tractional force:**

Useful for delivering the tooth from the socket.



**Rotational force** 



lingual force: lingual plate expansion, apical pressure



**Tractional force** 

7

#### **Clinical Notes:**

**A-** Majority of the teeth are removed by a combination of buccal & lingual forces.

**B-** In the maxilla, buccal bone is thin while palatal bone is thick, accordingly maxillary teeth are removed with more buccal forces and less palatal forces.

**C-** In the mandible, buccal bone is thin anteriorly but thick posteriorly (molar region), accordingly incisors, canines, premolars are removed with more buccal force and less lingual force, while the molars teeth removed with more lingual pressure.



#### **D-** Support is mandatory during forceps use.









### **Basic rules for using elevators**

1-The elevator must be held in the dominant hand and the index finger placed along the blade.

2-This instrument must always be used buccally, and never on the lingual or palatal side.

3-The concave surface of the blade must be in contact with the mesial or distal surface of the tooth to be extracted, and be seated between the tooth and alveolar bone.

4-During luxation, the adjacent tooth should not be used as a fulcrum, but only the alveolar bone. Otherwise, there is a risk of damaging the periodontal ligament fibers of the adjacent tooth. Unless multiple adjacent teeth to be removed in this case it can be inserted in between the two teeth since movement of the adjacent tooth will be desirable.





#### **5-** Support is mandatory during elevator use.









### **Fundamental Requirements for a Perfect Extraction**

1- Adequate access and visualization of the field of surgery.

2- An unimpeded (unobstructed) pathway for the removal of the tooth.

3- The use of a controlled force to luxate and remove the tooth.



Path of removal is obstructed

### **Methods of Tooth Extraction**

Erupted teeth could be extracted by two ways:

#### 1- The closed (routine) technique.

This is the most frequently used technique for every extraction.

#### 2- The open (flap or surgical) technique

This technique is indicated when the routine method fails to extract the tooth, it involves incision making, flap reflection, bone removal, tooth extraction then suturing.

#### Note:

The correct technique for any case should lead to an atraumatic extraction.

The wrong technique may result in traumatic and lengthy extraction.



### **Clinical Steps of the Closed Extraction Technique**

Practically speaking to remove a tooth we should:

1Tear PDL fibers that hold the tooth in the socket.

2 Expand the alveolar bony walls to provide unimpeded pathway.

These objectives are achieved by the use of the elevators and forceps as levers and wedges with increasing force.

### **Steps:**

A- loosening of soft tissue attachment from the cervical portion of the tooth (gingival separation) by the use of sharp instrument like periosteal elevator, desmotome or periotome.

#### **Objectives:**

1To ensure that profound anesthesia has been achieved.

2To allow the elevator & tooth forceps to be positioned more apically.









Soft tissue loosening (gingival separation)

**B-** Luxation of the tooth with a dental elevator.

#### Objectives:

- 1 Tearing of PDL fibers.
- 2 Expansion of the alveolar bone walls





**C-** Adaptation of the forceps to the tooth in which the beaks should grasp the root apically as far as possible (apical pressure).

#### Objectives:

1 Bony expansion (wedging action).

2Center of rotation of the tooth (fulcrum) displaced apically, and this decrease the chance of apical root fracture.







Crown grasp
**D-** Luxation of the tooth with the forceps, by buccal and lingual pressure ( the force applied should be slow steady pressure rather than series of rapid force) with frequent apical repositioning of the forceps. If anatomy of the root permit, rotational force applied.

# **Objective:**

- 1 Tear all PDL fibers.
- 2 Expansion of the bony socket.

E- Removal of the tooth from the socket by tractional force buccally (avoid damaging opposing teeth).



# **Order of Extraction**

- Anaesthesia---- more earlier in the maxilla
- Usually maxillary teeth are extracted first (debris and any fragments cannot be lost in open mandibular Socket
- Posterior teeth should be removed first for better vision --- blood can collect in posterior region

# **Extraction movement**

- Primary movement: Along longitudinal axis of root
- Secondary movement: Main extracting movement
  - Rotatory
  - Buccolingual or labiolingual
  - Mesiodistal
  - Lifting the tooth



- Upper central and lateral incisor
  - Rotation only
- Upper canine

Rotation initially, some labiolingual movement may be needed

Upper premolar and molar
 Bucco-palatal movement



 Lower central and lateral incisor

Rotatory movement

Lower canine

Rotatory and labio-lingual

Lower premolar

Rotatory

Lower molar

Bucco-lingual movement



# Dr. Hamid Hammad Enezei PhD

Oral & Maxillofacial Surgery





# **Thank You**

\*Academic Year 3\*





# Dr. Hamid Hammad Enezei

# **Ph.D in Oral & Maxillofacial Surgery**





# **Objectives:**

- 1- Describe chair & dentist position during the extraction.
- **2-** Understand the technique of extraction for each tooth.
- **3-** Outline the principles of extraction of deciduous teeth
- 4- Outline post extraction care for tooth socket.
- **5** List post extraction instructions
- **6** Identify the order and number of teeth to be extracted in one dental visit.



Fragiskos D. Fragiskos (2007): Oral Surgery. Springer Robinson (2007): Tooth Extraction- A Practical Guide. Wright Hupp (2013), Contemporary Oral and Maxillofacial Surgery. Alsevier





# **Chair – Dentist Position During Extraction**

# **A- Chair position:**

# **1- For maxillary teeth:**

1The chair should be tipped backward so that when the patient open his mouth the occlusal plane is at an angle of about 60 degrees to the floor.

2The height of the chair should be in such a way that patient's mouth is at or slightly below the operator's elbow level.





# 2- For mandibular teeth:

1The chair should be in more upright position so that when the patient open his mouth, the occlusal plane is parallel to the floor

2The height of the chair should be lower than for upper teeth which provide stable comfortable position.



Chair in upright position Height – less than upper



# **Chair – Dentist Position During Extraction**

# **B- Dentist position:**

# 1- For maxillary teeth:

For all maxillary teeth (left & right) the dentist should stand in front of the patient and to the right side during the extraction.

# 2- For mandibular teeth:

For left quadrant, the dentist should stand in front of the patient to the right side.

For right quadrant, the dentist should stand behind the patient in the right side.



# Note:

1-These positions are for a standing, right handed dentist, while for seated position or left handed dentist modifications required.

2-During extraction the left hand plays an important role in

A- Providing support and stability for the alveolar process and the jaw.

B- Provide tactile information to the surgeon regarding the progress of extraction.

C- Reflection of the soft tissues like lip, cheek, or tongue.





Behind

Role of the left hand during the extraction

# **Patient and surgeon Position**



For a maxillary extraction the chair should be tipped backward and maxillary occlusal plane is at 60 degrees to the floor. The height of the chair should be patient's mouth is at or below the operator's elbow level



For the extraction of mandibular teeth, the patient should be positioned in a more upright position. the occlusal plane is parallel to the floor. The chair should be lower than for extraction of maxillary teeth.

# **CHAIR AND PATIENT POSITIONS**

- Efficient patient and operator positions are beneficial for the welfare of both individuals
- A patient who is in a comfortable position is more relaxed, has less muscular tension and is more capable of cooperating with the dentist
- By using proper operating positions and good posture the operator experiences less physical strain
- The dental chair should support the patient's head, back and arms
- Modern dental chairs are the contoured to provide total body support and comfort for the patient in any chair position
- The patient's head is supported by the headrest cushion which is attached to the chair bach and is adjustable or articulated

# **CHAIR AND PATIENT POSITIONS**

- ✓ Chair and patient positions are important considerations. modern dental chairs are designed to provide total body support in any chair position
- ✓ Chair- position is a very important aspect in the success of a dental treatment
- The correct positioning helps the operator to have a good visibility and accessibility of the oral cavity
- Proper positioning of the patient and the operator, illumination and retraction for optimal visibility are the fundamental prerequisites to proper dental treatment
- If operator maintains proper position and posture during treatment, the operator is less likely to get strain, be more efficient and less chances of getting musculoskeletal disorders



















### ✤ FOR A RIGHT HANDED OPERATOR:

- i. Right front or 7'o clock position
- 2. Right or 9' o clock position
- 3. Right rear or 11'o clock position
- 4. Direct rear or 12' oclock position

#### LEFT HANDED OPERATOR'S POSITIONS ,

- 1- 5 O'CLOCK
- 2- 3 O'CLOCK
- 3- 1 O'CLOCK

# 4 zones of activity

# **DENTIST'S ZONE**

8-12 o'clock for righthanded dentist
12-4 o'clock for left handeddentist
STATIC ZONE
Instrumentarium
12-2 o'clock for righthanded dentist
10-12 o'clock for left-handed dentist









# **CHAIR POSITION**

- $\checkmark$  It should be able to provide comfort to the patient
- $\checkmark$  It should be able to provide total body support
- Headrest of chair should be attached for supporting patient's chin and reducing strain on chin muscles
- $\checkmark$  It should be able to provide maximum working area to the operator
- $\checkmark$  It should be placed at the convenient location with adjustable control switches
- $\checkmark$  Foot switches are preferred to improve infection control

# **CHAIR POSITION**

- The headrest supports the patient's head and slightly elevates the chin thus reducing strain on the neck muscles
- Some chairs are also equipped with programmable operating positions
- The patient should have direct access to the chair
- While making the patient sit on the dental chair, the chair height is kept low and the back rest is upright so that the patient can sit with ease
  - During examination or treatment procedures, the chair is tilted back
  - The switches for chair adjustment should be easily accessible to the dentist
  - Foot switches are preferred to use it, so the dentist can avoid hand contamination during treatment

# COMMON PATIENT POSITIONS



✓ 1- Upright Position this is the initial position of chair from which further adjustments are made

# Patient sitting in upright position of dental chair



And the second s

# ✓ 2- Almost Supine

# Patient is almost in a lying down posture

- The patient's head, knee and feet are approximately at the same level; however, the head should not be positioned lower than the feet except when the patient is in syncope.

• Patient's head should not be lower than feet except in case of syncope attack

# ✓ 3- Reclined 45 Degrees :

- In this position, the chair is reclined at 45 degree so that when the patient is seated, the mandibular occlusal surfaces are almost 45 degree of the floor.

# **Supine position**

-lying on back, facing upward-for Maxillary tr-dental chair back parallel to the floor

 The choice of chair position can be decided by the dentist based on the type of procedure and the arch being worked on



Distance terms and percent moves it 2008, 2008 by Standard, an improve prillegenet ter-

# Semi-supine position

For mandibular treatment areas -the back of the dental chair is slightly upright



# **Reclined 45 degrees**





Schematic presentation of correct dental chair positioning for pregnant patients

# **Specific Technique For Removal of Each Tooth**

# **Extraction of Maxillary Central Incisors**

#### **Instruments:**

Extraction forceps for six anterior maxillary teeth.

#### **Position:**

The dentist should stand in front of and to the right of the patient (right handed).

# Support:

By the index and the thumb finger.

# **Technique:**

The initial movements are in a labial direction, and then palatal. Since the root is conical in shape, rotational forces are applied and the final extraction force is applied labially.



#### **Extraction of Maxillary Lateral Incisors**

The same technique as for central incisors, the difference is that the extraction movements are labial and palatal only without rotational force because it has curvature of the root tip palato-distally.





#### **Extraction of Maxillary Canines**

The extraction technique is the same for lateral incisors with some degree of difficulty due to:

- 1 Their firm anchorage in alveolar bone.
- 2 The roots are long with frequent curvature of the root tip.

**3-** The labial surface of the root is covered by thin alveolar bone, so there is a risk of fracturing the alveolar process.





## **Extraction of Maxillary Premolars**

**Instruments:** Maxillary premolar forceps.

#### **Position:**

The dentist should stand in front of and to the right of the patient (right handed).

## Support:

By the index and the thumb finger.

## **Technique:**

1For the first premolar, because it usually has two roots, buccal and palatal pressure should be gentle and slight. If movements are vigorous and abrupt, there is a risk of fracturing the root tips. Rotational motions are not allowed due to the tooth's anatomy.

2For the second premolar the extraction is easier, because the tooth mostly has one root.

**3-** The final movement for both teeth is buccal





## **Extraction of Maxillary First Molars**

**Instruments:** Maxillary right & left molar forceps.

#### **Position:**

The dentist should stand in front of and to the right of the patient (right handed).

### Support:

By the index and the thumb finger.

# Note:

The maxillary first molar has three diverging roots, and the tooth is firmly anchored in the alveolar bone, therefore this tooth requires the application of strong force during its extraction, which may cause fracture of the crown or root tips.

## **Technique:**

To avoid tooth fracture, initial movements must be gentle, with buccopalatal pressure and an increasing range of motion, especially buccally, where resistance is less. The final extraction movement is a buccal curved motion.











#### **Extraction of Maxillary Second Molars**

**Instruments, Position, Support & technique:** The same as for maxillary first molars because the teeth have similar anatomy, however, it is easier because of little divergence of the roots and quite often are fused together in a conical shape.

#### **Extraction of Maxillary Third Molar**

**Instrument:** maxillary third molar forceps.

**Position and Support:** same as for upper molars. **Note:** The maxillary third molar is the smallest of all molars and varies greatly in size, number of roots, and root morphology. The tooth is usually easily removed because the buccal bone is thin and the roots are usually conical and fused.

**Technique:** If the third molar has erupted completely its extraction is accomplished by applying buccal pressure and very gentle palatal pressure. Root anatomy of the third molar may permit extraction using elevator only. The elevator is positioned in the mesio-buccal corner.









#### **Extraction of Mandibular Anterior Teeth**

**Instruments:** Mandibular anterior teeth forceps. **Position:** 

1For right side -- stand behind the patient in the right side.

2For left side-- stand in front of the patient in the right side.

### Support:

By index and thumb finger – right side. By index and middle finger – left side.

## Note:

**Mandibular incisors:** Their extraction is easy, due to their morphology (narrow, single roots) and the thin labial alveolar bone surrounding the root.

**Mandibular canines:** They are more difficult to extract compared to incisors, due to the long root and frequent curvature of the root tip.

**Technique:** Pressure is applied labially and lingually, gradually increasing in intensity, only slight rotational force is permitted. The final extraction movement is outwards and downwards (to avoid damaging opposing teeth).



**Buccal force** 

Lingual force

**Tractional force** 

#### **Extraction of Mandibular Premolars**

**Instruments:** Mandibular premolar forceps.

**Position and Support:** Same for anterior teeth.





#### Note:

Premolar extraction is considered quite easy because their roots are straight and conical.

#### **Technique:**

Buccolingual force is applied for extraction of these teeth. Gentle rotational force may also be applied. The final extraction movement is outwards and downwards.



**Buccal force** 

Lingual force

**Tractional force**
# **Extraction of Mandibular Molars** (first & second molars)

Instruments: Mandibular molar forceps.

**Position and Support:** Same for lower anterior and premolars.

#### Note:

The mandibular second molar has a morphology similar to that of the first molar, but its removal is more easily than the first molar, because its roots are smaller and less divergent.

**Technique:** The extraction technique is the same for both molars. Initially the movements are gentle with buccal and lingual pressure. After the tooth is slightly mobilized, force is gradually increased and the final extraction movement is buccal (outward & downward).





#### **Extraction of Mandibular Third Molar**

**Instruments:** Mandibular molar forceps

**Position and Support:** Same for molars.





#### Note:

The mandibular third molar usually has two roots, which are smaller and usually fused in a conical shape.



#### **Technique:**

**1-** Buccolingual pressure is applied, but since the lingual alveolar bone is very thin compared to the buccal alveolar bone, therefore, the force that mobilizes the tooth must be more in the lingual direction.

2- If the third molar has one root or if the roots converge, the extraction may be accomplished using the straight elevator alone.

The alveolar ridge is used as a fulcrum (correct placement)



Using the adjacent tooth (second molar) as a fulcrum (incorrect placement) creates great tension around the tooth, with a risk of injury to tissues surrounding the root.

Extraction of the mandibular third molar using a straight elevator

#### **Extraction of Deciduous Teeth**

1-The forceps used to extract deciduous teeth are small and have narrow beaks.

2-The extraction technique is similar to that used for permanent teeth and it is much easier, especially if their roots are resorbed.

**3-** The dentist must pay particular attention when extracting deciduous molars because of the risk of simultaneously extracting the bud of the permanent tooth.

#### **Preventive measure:**

The beaks of the forceps must be positioned on the mesial or the distal area of the tooth and not the center (root bifurcation), underneath which is the permanent tooth. Sometimes sectioning of the tooth or surgical extraction is indicated.





Deciduous mandibular molar with subjacent permanent tooth



## **Post Extraction Care of Tooth Socket**

1-The extracted tooth should be examined to be sure it is complete and not fractured (fractures site will be sharp while resorption site will be smooth).

2-The socket should be checked for the presence of any debris such as piece of bone, fragment of broken filling, root fragment or any sharp bone edges, if so, the alveolar margin is smoothed using rongeur forceps or a bone file also if there is any gum tear should be evaluated for the need of suturing.

3-The bottom of the socket is curetted (as long as the tooth is nonvital with periapical radiolucency) with a periapical curette, to remove any periapical lesion from the area, because if kept, there is a chance they will develop into a cyst (residual cyst). Sometimes the lesion is firmly attached to the root tip and is extracted together with the tooth. When the lesion is large and the entire lesion cannot be removed through the socket alone, then surgery is required.

4-In case of upper posterior teeth extraction, the socket should be checked for the presence of OAF specially with difficult extraction

5The lingual and buccal plates are compressed (squeezed) slightly using finger pressure to restore (oppose) the expansion of the socket caused by the extraction, and also for initial control of hemorrhage.



Nonvital tooth with periapical radiolucency



Curettage of the socket after tooth extraction for removal of the periapical lesion



Residual cyst

Root

Root with firmly attached lesion at tip

## **Post Extraction Instructions**

**1**-Bleeding: The patient must bite firmly on gauze placed over the wound for 30–45 min.

2-Analgesia: The patient should start taking painkiller as soon as possible before the action of local anesthesia get finished (paracetamol is adequate and safe for simple extraction).

**3**-Antibiotics: These are prescribed only if the patient has certain medical conditions or infection.

4-Rest: Avoidance of physical work or exercise is recommended for the remainder of the day since such activity may restart bleeding.

5-Diet: The patient's diet on the day of the extraction must consist of cold, liquid foods (yogurt, milk, cold soup, orange juice, etc.). Avoid smoking and alcohol intake.

6-Oral hygiene: Rinsing the mouth and brushing of the teeth are not allowed for the first 24 h. After this, the mouth may be rinsed with warm salt water, 4 times a day for 3–4 days as well as teeth brushing resumed.

















## **Order of Extraction**

1- Upper teeth should be removed before lower teeth due to:
A- Anesthesia of upper teeth achieved by infiltration technique which has rapid onset of action and disappears rapidly.

**B-** Debris such as portion of amalgams and fractured crowns may fall into the empty socket of the lower teeth if lower extraction is performed first.

**Disadvantage:** hemorrhage from upper teeth extraction may interferes with visualization during mandibular extraction.

2- The sequence of extraction should start with the most posterior then anterior one to avoid any bleeding that interferes with the visualization of the field.

### Number of teeth to be extracted in single visit

**1-** Depend on the general medical condition of the patient and the difficulty of the extraction.

**2-** Extraction of a great number of teeth at an out-patient one visit is contraindicated as it may upset the patient and the blood loss can be considerable.

**3-** Practically speaking, between four and eight teeth could be extracted in one visit but where it is necessary to do more, the patient should be admitted to hospital.



# **Intra Alveolar Extraction**

- $\checkmark$  It means direct access is gained on the tooth to be extracted.
- $\checkmark$  It is carried out with the help of dental forceps and elevators.
- Forceps Technique

## Indications

- Fairly mobile teeth,
- Extraction of single tooth,
- Extraction of multiple teeth which are scattered.

## \* Contraindications

- In complicated extractions,
- Deformed roots,
- Hard tissue pathology of roots like hyper-cementosis etc.
- Badly destroyed teeth

## ✤ Advantages –

- It causes the least amount of trauma.
- Promotes retention of a satisfactory blood clot.
- Favours rapid healing.

# Various movement for extraction of individual tooth

Upper central, lateral,	First apical →slight labial/buccal →
canine, first pre-molar,	slight palatal →rotation with
second pre-molar	traction
Upper molars	First apical $\rightarrow$ buccal $\rightarrow$ palatal $\rightarrow$ deliver the tooth buccaly
Lower central, lateral,	First apical → slight labial/buccal
canine, first pre-molar,	→ slight lingual → rotation with
second pre-molar	traction
Lower molars	First apical → buccal → lingual → deliver the tooth buccaly

## Elevator technique

Indications

• To luxate teeth which cannot be engaged by beaks of forceps eg: impacted, malposed or grossly destructed teeth.

• To remove roots

# **ROLE OF DENTIST'S LEFT HAND**



Hand Support During Extraction of Maxillary Central Incisor



## Hand Support During Extraction of Maxillary Premolar



## Hand Support During Extraction of Maxillary Molar



















# **Thank You**

\*Academic Year 3\*



# ARMAMENTARIUM OF ORAL SURGERY Part 1

# Dr. Hamid Hammad Enezei

## **Ph.D in Oral & Maxillofacial Surgery**



## **Objectives ( Part 1, 2):**

1-Identify the equipments, instruments and materials used in oral surgery.

2-Identify the instruments used to make:

Incision, Flap reflection, Bone cutting, Grasping, Curetting, Irrigation, Suturing.

**3-Identify the materials that used for:** 

Suturing , Hemostasis , Wound covering , Tissue regeneration.

4-Understand the preparation, action, and clinical indication of different types of materials used in oral surgery.



Further reading Fragiskos D. Fragiskos (2007): Oral Surgery. Springer Hupp (2013), Contemporary Oral and Maxillofacial Surgery. Alsevier

## **Armamentarium includes:**



## A-Instruments

## **B-** Materials

C- Equipments





## **A-Instruments**

**1- Scalpel (Handle and Blade)** ( They are used for incision making in the tissue)

#### A- Handle:

\* It is reusable (disposable also available).

\* The most commonly used handle in oral surgery is the Bard–Parker no. 3.

\*Its tip may receive different types of blades.

#### **B-Blade:**

\*Blades are disposable knives.

\*Three different types that used in oral surgery: (no. 11, 12, & 15).





Is the most common type of blade used in oral surgery, which is used for making an incision around teeth and on the alveolar ridges and alveolar mucosa.







The most common type. Incision around teeth and on the alveolar ridges and alveolar mucosa.











It is used for making small incisions, such as those used for incising an abscess.







### Gingival abscess

Vestibular collection of an abscess

It is indicated for making: 1 Incisions in the gingival sulcus. 2Incisions posterior to the teeth, especially in the maxillary tuberosity area.





Surgical procedure to remove fibrous tuberosity (thick)

#### Placement of the blade on the handle

By the help of a needle holder or hemostat, the blade is grasped and it slides into the slotted receiver with the beveled end parallel to that of the handle.

#### Note:

The scalpel is held in a pen grasp and its cutting edge faces the surface of the skin or mucosa that is to be incised







## 2- Periosteal Elevator



#### (Used for reflection of the gingiva or mucoperiosteum)

#### A- Molt no.9 :

Has two different ends: a pointed end, used for elevating the interdental papillae of the gingiva, and a broad end, which facilitates elevating the mucoperiosteum from the bone.

#### **B-** The Freer elevator:

It has thin ends, used for reflecting the gingiva surrounding the tooth before extraction.

#### **C-** The Seldin elevator:

It has broad ends, used for reflection as well as retraction of the flap during the surgical procedure.



### **3- Retractors**

#### (Used for retracting soft tissues)

A variety of retractors have been designed to retract the cheeks, tongue, and mucoperiosteal flap during the surgical procedure to provide:

1- Access for better visibility during surgery.

2- Protection for soft tissue from sharp cutting instruments.

Note: mouth mirror commonly used for cheek & tongue retraction.







## **4- Tissue Forceps**

Tooth like projection

Receptor

(Used for grasping soft tissue)



#### There are two designs:

1- One has wedge shaped projection at the tip ( tooth on one side, and a receptor on the other), which fit into each other when the handles are locked. This mechanism allows the forceps to grasp the soft tissues very tightly.

#### Uses:

This design used during suturing to grasp the tissues firmly while the needle is passed as well as in biopsy taking.

## **4- Tissue Forceps**

Tooth like projection

Receptor

(Used for grasping soft tissue)

**2-** Without wedge-shaped projection, but grooves at the tip.

#### Uses:

This design used during suturing to grasp the tissue as well as grasping small instruments during the surgical procedure.



#### **5- Bone Burs**

#### (Used for removing bone)

\* They are available in different shapes and sizes like round bur and fissure bur.







\*A large bone bur similar to an acrylic bur may be used when the surgical procedure involves greater bone surface area (like torus palatinus) or smoothing of bone edges of the wound.

**Note:** Good irrigation system should accompany the use of these burs to overcome the problem of heat generation.









6- Rongeur Forceps (Bone cutter) (Bone nibbler)

(Used for removing bone)

There are 2 designs :

- 1 Side cutting forceps
- 2 Side & end cutting forceps

Side & end cutting forceps is the most practical one in oral surgery since they are end cutting so they can be inserted into the sockets for removal of inter-radicular bone after extraction.











#### **7- Bone File**

This instrument is used to smooth bone, not to remove large pieces of bone.

It has two ends: one small end and another with a large surface .

The cutting surface is made up of many small parallel blades, which are set (directed) in such a way that only pulling action is effective (pull stroke).









Steps of smoothing irregular alveolar ridge by bone file



## 8- Bone Chisel and Mallet

These instruments are used together for bone removal or sectioning (bone or tooth).

\*Mallets are instruments with heavy-weighted ends, the surfaces of the ends are made of lead or of plastic so that some of the shock is absorbed when the mallet strikes the chisel.

\*Bone chisels available in different shapes and sizes, their cutting edges are concave, monobeveled or bibeveled.

\*The bibevel chisel is used for sectioning multi-rooted teeth.

\*They are often used in lingual tori removal









## 9- Periapical Curettes

(Used for removing soft tissue from bony cavities)

1-They are angled double-ended, spoon-shaped instruments.

2-Their shape facilitates its entry into bone defects and extraction sockets.

**3-** The main use of this instrument is the removal of granulation tissue, small cysts, bone chips or foreign bodies.



Curettage of the socket after tooth extraction for removal of the periapical lesion



#### **10- Mouth Props & Bite Blocks**

**Mouth props**: facilitate opening and keeping the mouth open.

**Bite blocks:** keep the mouth open when the surgical procedure requires mouth opening for long period and when patients can not fully cooperate with the dentist.







## **Summary**

1- Scalpel (Handle and Blade)( They are used for incising tissue)

**2- Periosteal Elevator** (Used for reflection of the gingiva or mucoperiosteum)

**3- Retractors** (Used for retracting soft tissue)

4 Tissue Forceps (Used for grasping soft tissue)

5 Bone Burs (Used for removing bone)



6- Rongeur Forceps (Used for removing bone)



7- Bone File (Used to smooth bone)







8- Bone Chisel and Mallet (Used for bone removal or sectioning (bone or tooth)









10MouthProps& BiteBlocks(Used for opening or keepingthe mouth open)







# **Thank You**

\*Academic Year 3\*



# ARMAMENTARIUM OF ORAL SURGERY Part 2

# Dr. Hamid Hammad Enezei

## **Ph.D in Oral & Maxillofacial Surgery**


### **Objectives ( Part 1, 2):**

1-Identify the equipments, instruments and materials used in oral surgery.

2-Identify the instruments used to make:

Incision , Flap reflection , Bone cutting , Grasping , Curetting , Irrigation , Suturing.

**3-Identify the materials that used for:** 

Suturing , Hemostasis , Wound covering , Tissue regeneration.

4-Understand the preparation, action, and clinical indication of different types of materials used in oral surgery.





Further reading Fragiskos D. Fragiskos (2007): Oral Surgery. Springer Hupp (2013), Contemporary Oral and Maxillofacial Surgery. Alsevier

## **A-Instruments**

1- Scalpel (Handle and Blade)( They are used for incising tissue)

2- Periosteal Elevator (Used for reflection of the gingiva or mucoperiosteum)

**3- Retractors** (Used for retracting soft tissue)

4 Tissue Forceps (Used for grasping soft tissue)

5 Bone Burs (Used for removing bone)



6- Rongeur Forceps (Used for removing bone)



7- Bone File (Used to smooth bone)







8- Bone Chisel and Mallet (Used for bone removal or sectioning (bone or tooth)









10MouthProps& BiteBlocks(Used for opening or keepingthe mouth open)





# **11- Hemostats** (Artery forceps)

(Used for bleeding control by clamping the bleeding vessel)

1- It is either straight or curved .

2-The curved type (mosquito) is most commonly used one.

3-The beaks of the hemostat have parallel grooves.

**4-** Hemostats may also be used for holding soft tissue during biopsy taking, and in drainage an abscess cavity .





## **12- Needle Holders**

### ( Used for holding the needle during suturing )



**1-** It looks similar to a hemostat.

2- In the needle holder, the grooves in the internal surface of the beaks is crosshatched, permitting a firm and stable grasp of the needle, while the beaks of the hemostat have parallel

grooves.





## **12- Needle Holders**

### **Used for Holding the Needle During Suturing**



## 13- Scissors ( Used for cutting )

- Two types
  1 Suture scissors
- 2 Soft tissue scissors.





**1-** Suture scissors have sharp, **short cutting** handle since their main use is to cut sutures.

blades and long

2- Soft tissue scissors have sharp, long cutting blades with sharp or rounded tips and are used for removing excess gingival tissue, dissecting and undermining the mucosa from the underlying soft tissues.





## **13- Scissors Used for cutting**

Two types 1 Suture scissors

2 Soft tissue scissors.









#### **Castroviejo Micro Scissors:**

Have a unique design that allow extremely smooth cutting of fine and coarse tissue with reduced trauma.

Beaks are straight or curved with serrated handle (non slip grip)

### **14- Irrigation Instruments**

Irrigation with saline solution during bone removal is necessary and achieved by using either plastic syringe with a blunt needle or by using special irrigation system that gives a steady stream of saline solution.

#### **Advantages:**

1 It cools the bone burs & prevents heat generation in the bone.

2It increases the efficiency of the bone burs by washing away bone chips from the flutes of the burs.

3Irrigation of the under surface of the flap before suturing is important to remove any debris that interfere with the wound healing.







## **Instruments - Summary**

**1- For incision making: Scalpel** 



9- For curetting cavity: **Periapical Curette** 



#### **2- For flap reflection: Periosteal Elevator**

**3- For flap retraction: Retractor** 

**4For grasping tissue: Tissue Forceps** 

**5For bone removal: Bone Burs Rongeur forceps Bone chisel & mallet** Bone1fOile( smoothing)







**11For bleeding control: Hemostat** 

12 For holding needle in suturing: **Needle Holder** 

**13For cutting: Scissors** 

**14For irrigation: Plastic syringe Irrigation system** 















## **B-** Materials

#### **1- Suture Material (stitch)**



#### **3- Local Hemostatic Agents**



5- Materials for Bone Regeneration



#### **2- Suture Needles**



### 4- Materials for covering wound



## **B- Materials**

## **1- Suture Material (stitch)**

**Used** to reapproximate the wound edges & protect underlying tissues from infection or other irritating factors.

**Different types** of suture material are available & classified according to their:

Diameter. Resorbability. Whether they are monofilament or polyfilament.







## 2- Suture Needles

A variety of needles are available, they differin:

Shape (straight, curved).

Diameter.

**Cross-sectional view.** 

Size (length).



### **3- Local Hemostatic Agents**

These agents are suitable for **local use** only and can stop bleeding which is due to injury of small blood vessels. They are fully absorbed by the tissues within weeks.

#### The main hemostatic agents are:

#### **A - Natural Collagen Sponge:**

**preparation:** It is a white sponge material, nonantigenic.

#### **B- Gelatin Sponge (gelfoam):**

Preparation: sponge material, nonantigenic.

**C- Oxidized Cellulose ( surgicel ): preparation:** It is available in gauze form or pellet form.

Indication: Post extraction bleeding socket



Surgicel - gauze





#### **D- Bone Wax:**

**Preparation:** It is available as a sterilized, nonabsorbable solid plate of wax

**Indication:** It is used to control bleeding that originates in bone or chipped edges of bone.

Action: Its hemostatic action is brought by mechanical obstruction of the osseous cavity, which contains the bleeding vessels.









4- Materials for covering wound or filling a surgical cavity

#### **A- Vaseline Gauze:**

**Preparation:** Gauze in sterilized packages

#### Indication:

For covering exposed wounds, bone cavities (ex: after cysts removal).





Surgical cavity

#### **B- Iodoform paste & Gauze:** Preparation:

Available as a ribbon gauze or paste that has antiseptic, analgesic and hemostatic properties.

#### Indication:

For covering exposed wounds, bone cavities as well as for the treatment of dry socket.

16



Iodoform paste



#### Iodoform Ribbon Gauze

#### **C- Surgical Dressing:**

#### **Preparation:**

This is an autopolymerized putty like paste, available in sterilized package.

#### Indication:

It is used in periodontal and oral surgery as a temporary protective covering of wounds after surgical procedures until healing occurs.





### **5- Materials for Bone Regeneration**

These materials are used whenever there is a surgical procedure that create a bony defect (removal of cysts, extraction of impacted teeth, etc.) to aid in bone regeneration and eliminates the defect or limits its size.

#### **Indication:**

1- Periodontal surgery: useful in the regeneration of periodontal tissues.

2 Implant surgery: for filling of bone defects around an implant.

3Oral surgery: for augmentation of a deficient alveolar ridge for better prosthesis.

## **Types:** 1- Membranes:

These may be: Absorbable: like collagen membrane. Nonabsorbable: like titanium reinforced membrane.



#### Absorbable collagen membrane



membrane in an area of bone deficit after surgical extraction

## 2- Bone Grafts:

Types of bone graft:

- 1- Autografts, which are composed of tissues from the same patient.
- 2- Allografts, which are composed of tissues from another individual.
- 3- Heterografts (Xenografts) which are composed of tissues from various animals.
- 4- Alloplastic grafts, which are composed of synthetic bone substitutes, e.g., hydroxylapatite, and calcium hydroxide in cream form











hydroxylapatite granules

## **C-Equipments**

## **1- Sugical Handpiece**

Advantages : 1-It functions at high speeds and has great cutting ability.

2-It does not emit air into the surgical field.

### 2- Electrosurgical Unit

This is an electrical device, providing high-frequency electrical current used for:

A- Cauterization of the vessels (hemostasis).

B- Making incision of tissues (electrosurgery).













### **3- The Piezoelectric Bone Surgery** ( **Piezosurgery**)

It is a new technology for osteotomy (bone cutting), which utilizes the ultrasonic microvibrations of the blade .

It was invented to overcome the limits of precision and security of traditional instruments in bone surgery.





### **4- LASER MACHINE**

### Advantages:

1-Certain procedures performed using dental lasers may not require anesthesia or sutures.

2-Laser dentistry minimizes bleeding because the high-energy light beam aids in the clotting of exposed blood vessels, thus inhibiting blood loss.

3-Bacterial infections are minimized because the high-energy beam sterilizes the area being worked on.

4-Damage to surrounding tissue is minimized.

5-Wounds heal faster and tissues can be regenerated.







### **Dental applications:**

**1-** Oral surgery: ex: frenectomy procedure or removal of a benign lesions.

2- Periodontal surgery; ex: crown lengthening procedure.

3- Conservative dentistry: Ex : cavity preparation, endo disinfection.

- 4 -Teeth sensitivity.
- 5 Teeth whitening procedure.
- 6 -TMJ disorder.











## **Thank You**

# Methods of administration of local anesthesia by

# Dr. Elham Hazeim Abdulkareem University of Anbar College of Dentistry

The methods of administration of local anesthesia are sub-divided into three methods: topical, infiltration, and regional techniques.

1. Topical or surface anesthesia: it is obtained by the application of a suitable anesthetic agent to an area of either skin or mucous membrane in which it penetrates to anesthetize superficial nerve endings. It is commonly used to obtain anesthesia of mucosa prior to injection.

2. Sprays containing an appropriate local anesthetic agent are particularly suitable for this purpose because of their rapidity of action.

The active ingredient is 10% lidocaine hydrochloride. The onset time of anesthesia is approximately 1 minute and the duration round about 10 minutes.

**3.** Ointments containing 5% lidocaine hydrochloride can be used for a similar purpose but take 3-4 minutes to produce surface anesthesia.

Ethyl chloride spray when sprayed on to either skin or mucosa volatilizes so rapidly that it quickly produces anesthesia by refrigeration

This technique is of limited value but is occasionally used to produce surface anesthesia prior to incision of a fluctuant abscesses

## Maxilla

The maxilla has thin labial/buccal cortical plate; and moreover shows areas of porosity, and the compact bone presents numerous foramina which aid in absorption of local anesthetic solution. These factors, therefore, make the maxilla more favorable for infiltration anesthesia techniques.

## Mandible

The bone is generally dense and has thicker cortical plates than maxilla, particularly in posterior region, more so in the region of external oblique ridge. Only the anterior part of mandible presents sufficient porosity, which is favorable for infiltration techniques.

## 2. Infiltration anesthesia:

In this method the anesthetic solution deposited near the terminal fibers of any nerve it will infiltrate through the tissues to reach the nerve fibers and thus produce anesthesia of the localized area served by them, this technique is subdivided into:

## A. Sub-mucous injection:

in this technique the solution is deposited just beneath the mucous membrane. This technique is unlikely to produce anesthesia of the dental pulp, it is often employed to anesthetize the long buccal nerve prior to the extraction of mandibular molars or for soft tissue surgery



Fig.: Infiltration—submucosal injection the position of the point of the needle is in the submucosal layers.

## **B.** Supra-periosteal injection:

- in some sites, such as the maxilla, the outer cortical plate of alveolar bone is thin and perforated by tiny vascular canals.
- In these areas when anesthetic solution is deposited outside the periosteum, it will infiltrate through the periosteum, cortical plate, and medullary bone to the nerve fibers. By this means, anesthesia of the dental pulp can be obtained by injecting alongside the approximate position of the tooth apex.
- The Supra-periosteal injection is the technique most frequently used in dentistry.





Fig. 2: Infiltration—supraperiosteal injection in anterior maxilla— the position of the point of the needle is at an angle of 45° to the long axis of the tooth.

# C. Sub-periosteal injection:

in this technique the solution is deposited between the periosteum and the cortical plate. This technique is painful since the periosteum is firmly bound to the cortical plate



Fig.: Infiltration subperiosteal injection in anterior maxilla—the position of the point of the needle is at an angle of 90° to the long axis of the tooth and the alveolar bone as seen from the side.

## **D. Intra-osseous injection:**

in this technique, the solution is deposited within the medullary bone. The procedure is carried out by the use of bone drills and needles especially designed for this purpose.

In practice, the effectiveness of the anesthetic solution available has reduced considerably the need for intraosseous injection and the technique is seldom used.



## E. Intraligamentary (Periodontal or Peridental) Injection

## Nerve anesthetized:

Terminal nerve endings at the site of injection at the apex of the tooth.

## Area anesthetized:

bone, soft tissue & apical & pulpal tissue in the area of injection.

## **Indications:**

- 1. Pulpal anesthesia of one or two teeth.
- 2. Situations in which regional block are contraindicated (e.g.: bleeding disorders).
- 3. Aid in the diagnosis of pulpal discomfort.
- 4. As an adjunctive technique following nerve block if partial anesthesia is present.



Fig.: Infiltration intraligamentary injection—the position of the point of the needle is in the periodontal space Contraindications:

- **1. Infection or acute inflammation in the area of injection.**
- 2. Patient who requires (numb) sensation for psychological comfort.

#### Advantages:

- 1. Avoid anesthesia of the lip, tongue & other soft tissues.
- 2. Minimum dose of local anesthesia is required.
- 3. Rapid onset of profound pulpal & soft tissue anesthesia.
- 4. Less traumatic than conventional block technique.

#### **Disadvantages:**

- 1. Excessive pressure or rapid injection may break the glass cartridge.
- 2. A special syringe may be required.
- **3.** Excessive pressure can produce focal tissue damage.

#### **Technique:**

- 1. A 27 gauge short needle recommended.
- 2. Area of insertion; the long axis of the tooth to be treated on its mesial or distal of the root.
- **3.** Target area; is the depth of the gingival sulcus.

## **Intrapulpal Anesthesia**

This technique is indicated for obtaining anesthesia for procedures which require direct instrumentation of the pulpal tissue.

A 25 or 27-gauge needle is inserted directly into the pulp chamber.

The needle should be held firmly or wedged into the pulp chamber or the root canal.



Initially, slight discomfort is felt by the patient which subsequently gets subsided. Sometimes the needle is bent to get a proper angle for good approach Fig.: Infiltration—intrapulpal injection—the point of the needle is in the pulp chamber (*Courtesy:* Dr Vibha Hegde)

## 3. Regional (block) anesthesia:

In this technique, the anesthetic solution deposited near a nerve trunk will, by blocking all impulses, produce anesthesia of the area supplied by that nerve.

Although this technique may be used in the maxilla, it is of particular value in dentistry for producing anesthesia in the mandible.

The use of infiltration technique in the mandible is unreliable due to the density of the outer cortical plate of bone.

By placing the anesthetic solution in the pterygomandibular space near the mandibular foramen, regional anesthesia over the whole distribution of the inferior dental nerve on that side is obtained.

- Local anesthesia in the maxilla:
- Anesthesia of the upper molar teeth:
- The pulp of upper third, second, and first molar (with the exception of the mesiobuccal root) are innervated by branches of posterior superior alveolar nerve.
- This nerve is also responsible for providing sensory innervation to the buccal gingiva in the molar region and the mucoperiosteal attached to them.
- Deposition of the anesthetic solution close to the nerve (posterior superior alveolar nerve) after it leaves its bony canal produces regional anesthesia of the structures it supplies.
- This technique is termed posterior superior dental block.
- This technique carries the risk of damage to the pterygoid venous plexus so it is of limited use; instead infiltration technique is commonly used. <sup>13</sup>
Technique of posterior superior alveolar nerve block (fig: I):

- 1. Partially open the patient's mouth, pulling the mandible to the side of injection
- 2. Retract the patient's cheek with your finger or mirror (for visibility)
- 3. Insert the needle into the height of the mucobuccal fold over the second molar.

4. Advance the needle slowly in an upward, inward, and backward direction in one movement

- 5. Deposit 1 -1.8 ml of the anesthetic solution
- 6. Slowly withdraw the syringe.





# **Infiltration technique:**

- **1.** Hold the syringe parallel with the long axis of the tooth.
- 2. Insert the needle into the height of the mucobuccal fold over the target tooth.
- **3.** Advance the needle for a few millimeters.
- 4. Inject slowly about 0.6 ml (third of a cartridge)
- 5. Slowly withdraw the syringe.

<u>Note:</u> in the block technique we get anesthesia to the whole area supplied by the nerve (posterior superior dental nerve), while in the infiltration technique, the anesthesia is limited to the specific target tooth.

# Anesthesia of the upper premolar teeth

The mesio-buccal root of the upper first molar, both premolars, and buccal supporting tissue and mucoperiosteum related to them are innervated via the middle superior alveolar nerve. Infiltration technique is usually employed to anesthetize these structures (same procedure mentioned above).





## Anesthesia of the upper anterior teeth

The upper central, lateral incisor, and canine teeth and labial supporting tissue and muco-periosteum related to them are innervated via anterior superior dental nerve.

Infiltrationtechniqueisusuallyemployedtoanesthetizethesestructures(sameprocedurementioned above).

**Note:** in conservative treatment of a tooth we require anesthesia of the pulp only, while during surgical procedure such as tooth extraction we need anesthesia of the pulp, buccal and lingual gingiva and the associated bone and periosteum.



Palatal anesthesia is necessary for dental procedures involving manipulation of palatal soft or hard tissues (ex: tooth extraction).

The palatal mucopriosteum is firm in consistency and is closely adapted to the bone.

These features make it necessary to inject local anesthetic solutions under greater pressure than is required in other sites.

For his reason, patients should be informed prior to injection that palatal injections cause some discomfort or pain. This may be minimized by inserting the needle with the bevel facing the bone and as near as possible **at right angel to the vault of the palate.** 

Palatal anesthesia could be achieved by greater palatine nerve block, which provides anesthesia of the posterior portion of the hard palate,

or nasopalatine nerve block, which provides anesthesia of the anterior portion of the hard palate.

Or by infiltration technique in which the solution is deposited in the palatal tissue adjacent to the target tooth.

Note: In the greater palatine nerve block (fig.4), the solution is deposited near the greater palatine foramen, which is located just distal to the maxillary second molar about 1 cm toward the midline.





In nasopalatine nerve block (fig:5), the solution is deposited near the incisive foramen, which is located in the midline of the palate about 1cm posterior to the maxillary central incisors.

In the infiltration technique the solution is deposited about half way between the midline and the gingival margin of the target tooth.





### **Infra- Orbital injection:**

This technique is rarely used since the infiltration techniques are so effective in the maxilla, however it may be of value if numerous extractions or extensive surgery are to be undertaken in the maxillary incisor and canine regions.

It may also be employed for anesthetizing an anterior tooth where the use of infiltration technique is precluded by the presence of infection at the site of injection.

This technique is based upon the fact that the solution deposited at the orifice of the infra-orbital foramen passes along the canal to involve both the anterior and middle superior alveolar nerves thus producing anesthesia of the incisor, canine, and premolar teeth and their supporting structures. This block technique could be achieved either by an intra oral or an extra oral approach.

### Intra oral approach

This approach is more popular and allows the needle to be kept out of the patient's sight.

### **Technique:**

The infra-orbital ridge is palpated and the infra-orbital notch located with the tip of the first finger, which is then moved slightly downward to lie directly over the infraorbital foramen.

With the finger tip maintained in that position the thumb is used to reflect the upper lip and expose the site of injection.

Then the tip of a long needle is inserted into the mucous membrane over the apex of the second premolar. The needle is advanced in line with the long axis of this tooth to a depth of 1.5-2 cm. Then 1 ml of anesthetic solution is given which is enough in most instances.





### Extra-oral approach

The infra-orbital foramen is located in the manner previously described, the patient told to close his eyes, then the tip of a short needle is inserted until it lies at the foramen followed by deposition of 1 ml of the solution.

## Anesthesia of the upper deciduous teeth:

In children, multiple vascular canals perforate the thin labio-buccal alveolar plate. For this reason, infiltration techniques are highly effective in producing anesthesia of upper deciduous teeth.

Care should be taken to avoid the mistake of misjudging the length of the roots and inserting the needle too deeply into the tissues.

#### Local anesthesia in the mandible

Due to the density of the buccal bone plate, infiltration techniques are of limited value in the mandible and block anesthesia is most frequently employed. This is achieved by the deposition of solution around the inferior dental nerve in the pterygomandibular space.

#### Boundaries of the pterygomandibular space

- The space is bounded anteriorly by the pterygomandibular raphe and the fibers of superior constrictor and buccinator muscles that are inserted into it.
- The posterior boundary is formed by the parotid gland.
- The ramus of the mandible forms the lateral wall while the medial pterygoid muscle forms the medial wall; the lateral pterygoid muscle forms the roof of the space.
- The lingual nerve ascend diagonally through the space passing just infront of the inferior alveolar nerve as the latter emerges from the mandibular foramen.
- A shallow bony depression just above the foramen is the site in which the solution should be deposited for an inferior dental block.



# The inferior dental block (IDB):

The success of this technique is entirely dependent upon the accurate *deposition* of *the* solution; this technique will anesthetize the inferior alveolar nerve and its terminal branches (mental and incisive nerve).

It is accomplished in the following manner: The patient is seated in the chair and the head rest adjusted so that his mandibular occlusal plane is horizontal when the mouth is open.

The dentist should stand in front of his patient for the right IDB and behind the chair for the left IDB (this is for the right handed dentist). Upon intra-oral inspection, two important landmarks should be identified: the retromolar triangle and the pterygomandibular raphe (this structure passes upward and inward from the posterior end of the mylohyoid line of the mandible to the Hamulus of the medial pterygoid plate, the point of insertion of the needle should be lateral to and infront of it),



The thumb of the left hand is passed along the buccal surfaces of the lower molar teeth until the external oblique ridge is felt. The tip of the thumb is then passed inward to lie in the retro-molar fossa. The mid point of the nail should lie in the deepest part of the coronoid notch; this position usually coincides with the internal oblique line.

A long needle is inserted at this point. With the barrel of the syringe held parallel to the mandibular occlusal plane and over the second premolar tooth of the opposite side of the mouth, the tip of the needle is inserted for about 2 - 2.5 cm until its tip lightly contact the bone above the mandibular foramen.

When bone is contacted withdraw approximately 1mm to prevent subperiosteal injection and about 1.5 ml of the solution deposited. Then slowly withdraw the syringe and when approximately half its length remains within tissues deposit few drops of the solution to anesthetize the lingual nerve. The onset of anesthesia is signalled by a change of sensation in the lower lip and the tongue when compared with the other side; this may be described as pins and needles, frozen, wooden like, etc...





### Note

1. The dimensions and shape of the mandible may vary in patients of differing race, size, and age. Thus the width of the ascending ramus and hence the position of the mandibular foramen may vary between individuals. For this reason it is often helpful to palpate both the anterior and the posterior borders of the ascending ramus by the thumb and the index finger, the needle may then be inserted as previously described and directed midway between the two fingers.

2. Bilateral IDB should be avoided whenever possible since it produces considerable discomfort primarily from the lingual soft tissue anesthesia which usually persist for several hours after injection, in addition, the patient feels unable to swallow and because of lack of sensation he\she is more likely to self injury the anesthetized soft tissue.

3. Lingual nerve also could be anesthetized by infiltration technique, by deposition of about 0.5 ml of the solution in the lingual sulcus adjacent to the target tooth.

### Long buccal nerve block (LBB):

The long buccal nerve provides sensory innervation to the buccal soft tissues adjacent to the mandibular molar only (buccal gingiva). The main indication for LBB is during any procedure that involves manipulation of these tissues (ex: during extraction of the molar teeth). LBB is achieved by means of a submucous injection in which the solution is deposited (few drops) just posterior and buccal to the last molar tooth in the arch.

Infiltration technique is achieved by deposition of the solution in the muco-buccal fold adjacent to the target tooth.

#### Note:

1. When the long buccal nerve anesthetized, the patient rarely experiences any symptoms due to the small size of the anesthetized area.

2. The depth of penetration of the needle is seldom more then 2-4 mm, and usually only 1-2 mm.





# **Anesthesia of the lower anterior teeth:**

- The lower anterior teeth are supplied by the terminal branch of the inferior dental nerve (incisive nerve).
- In the mid-line of the mandible there is a considerable overlap (anastomosis) between the left and right incisive nerve, and it is this anastomosis that renders an inferior dental block sometimes ineffective in the incisor region.



• Fortunately the labial alveolar plate in this region is thinner and more porous than elsewhere; accordingly the infiltration technique could be used in this area. This is achieved by deposition of about 1 ml of the solution in the labial sulcus adjacent to the target tooth.

### The mental nerve block (MNB):

When the anesthetic solution given near the mental foramen, it will enter the inferior dental canal to produce anesthesia of the premolar, canine, and incisor teeth of that side the mental foramen, which is usually situated between the apices of the premolars, could be determined by an imaginary line passing vertically from the supraorbital notch through the infra-orbital foramen will usually cross the mental foramen when the mouth is closed.



About I ml of the solution should be injected over the foramen.

# Anesthesia of the lower deciduous teeth:

In children, multiple vascular canals perforate the labiobuccal alveolar plate, for this reason, infiltration techniques are highly effective in producing anesthesia of lower deciduous teeth. However, inferior alveolar block technique has the advantage of one needle puncture.

# **Complications of local anesthesia**

# Introduction:

Administration of local anesthesia is daily routine for most dental practitioners normally no adverse effects are seen. However complications even very serious one can occur in daily practice.

# Complications related to local anesthesia can be divided into tow categories:

- 1. Preoperative (during giving anesthesia) complications.
- 2. Postoperative complications.

Both can usually be avoided buy using the correct technique and dosage however if complications occur the dentist should know how to manage them.

### Each dentist dealing with local anesthesia should know:

- 1. The possible complications of local anesthesia.
- 2. The preventive measures of these complications.
- 3. The treatment possibilities of these complications.

# **Preoperative complication**

# 1. Pain at injection

Pain during administration of LA solution can be caused by many reasons.

### **A- Factor depending on the solution:**

1-Low PH value; this could irritate the tissue.

2-Temperature of the solution: warmer solution feeling more comfortable than cold. The cartridge can be warmed in the dentist hand or in warm water before the injection.

### **B-** The practitioner related factors:

This is related to the technique used can be avoided by:

1- Fast injection and high injection pressure cause rapid swelling of the tissue & pain. This a slower injection technique.

2- Aggressive insertion of the needle can tear soft tissue, blood vessel, nerve & periosteum & cause more pain & other.

### Complications

An inaccurate injection site can lead to an intramuscular or intraneural inj. when the needle penetrates a nerve; the patient feels a sudden (Electrical shock) in the distal area of the nerve. Pain after intramuscular injection is due to fibrosis or inflammation inside the muscle.

# 2. Lack of effect:

Some time the dentist face a problem, where the patient despite a conventional anesthesia still feels pain during treatment.

The problem is most common with block anesthesia especially in lower jaw. Reasons behind his failure in LA can be classified as:

- 1- Anatomical.
- 2- Pathological.
- 3- Psychological.
- 4- Poor inj technique

# **1-Anatomical reasons include :**

- accessory nerve supply
- Abnormal course of he nerves.
- Thick cortical plate of the mandibular alveolus precludes infiltration of anesthesia & therefore infiltration anesthesia is insufficient in the lower jaw.
- Variation in the foramen location.
- Some times the teeth can be innervated by more than one nerve.

### 2- pathological reasons are :

A- Trismus (limited mouth opening) in these cases it is impossible to use conventional technique of inferior alveolar nerve block. Therefore the so called Akinosi (closed mouth) technique is useful. In these cases there will be an increased possibility of failure of achieving adequate anesthesia.

**B-** Infection & inflammation: if the pulp is inflamed, the low tissue PH may cause lack of effective anesthesia in that area. However this does not explain failure in block anesthesia, where the solution is inj 4-5 cm from the inflamed area. A possible cause is hyperalgesia; the inflammation makes the nerve more sensitive. A minimal stimulation can cause pain perception. In those patient to obtain sufficient anesthesia more solution has to be injected, for example by combining a block & infiltration anesthesia. Supplemental intra-ligamentary or intra osseous inj can be used if necessary.

C- Previous surgery or trauma.

### **3- Psychological reasons :**

Ex fear & anxiety. They can cause failure in LA, to enable successful anesthesia, relaxation of the patient is some time needed. For this the use of a sedative like benzodiazepine may be helpful

**4- Poor technique:** Is the most common cause for insufficient anesthesia in inferior alveolar nerve block anesthesia common mistakes are:

A- Injection of anesthesia too soon on the anterior ascending ramus as the needle point touches the lingual cortical bone anterior to lingula

B- Another mistake is to injection inferior to mandibular foramen

C- The solution can directed away from a nerve if it injection too rapidly & forcefully. So when ever possible it is advisable to follow accurate technique to prevent this complication.

#### **3- Fainting & vasovagal attack**

Collapse of the patient in the dental clinic may or may not be loss of consciousness. Vasovagal attack is a reflex of nervous system that cause the heart to slow down (bradycardia ), at the same time affect he nerves to blood vessel in the legs permitting these vessel to dilate (widen ). As a result the heart ejects less blood so the blood pressure drops. Consequently the circulating blood goes to the legs rather than to the head, so the brain is deprived of 02 & the fainting episode occurs. In an attempt to redistribute the blood to the vital organs.

*Note:* pain & anxiety are the most important predisposing factor to vasovagal attacks.

#### Signs and symptoms:

The patients often complain of the following symptoms:

1-Dizziness 2-weakness 3-nausea

& following signs:

I-The skin is pale immediately before the collapse.

2- Cold skin 3- clammy skin 4- weak & slow pulse.

#### First aide should be taken immediately:

The head should be lowered quickly by adjusting the back of the dental chair so that the patient assume the supine position with legs elevated. Tight collar & belts should be loosened and respiration is stimulated. Spontaneous recovery is usual, but if sign of recovery are not apparent within 30-45 sec of the first aide measure, the collapse probably is not a vasovagal attack & in this case medical emergency team should be asked immediately for help. Meanwhile, we need to maintain a patent air way & supplementary 02 & consider the need for CPR (cardio-pulmonary resuscitation).

# 4. Hypersensitivity or allergy to LA :

This phenomenon occurs due to sensitivity of some individual to certain substance known as (allergens).

Any LA agent may evoke such a response, but it is more commonly seen with ester type agent (e.g. Procaine) than amide type (e.g. lidocaine) .true allergy to amide type is extremely rare.

#### Hypersensitivity reaction could be due to:

1- LA agent.

- 2- Vasoconstrictor.
- 3- Additives: e.g. bisulphite which used as preservative.

In general hypersensitivity reaction to LA is very rare & represent less than 1% of all complications of LA.

Many of complications suspected to be allergic are actually psychogenic reactions caused by fear of dental treatment.

Other reasons may also be the presence of adrenaline in LA solution which can cause several general symptoms including palpitation, restlessness & nausea.

True allergic response to LA may be localized or generalized & immediate or delayed in onset .they may vary from mild skin irritation or rashes to an anaphylactic reaction. Local reactions are seen more frequently than systemic & usually resolve without active treatment. If any degree of allergic reaction is observed, it is very important to determine the actual cause (allergen). Inadequate diagnosis & treatment can be life threatening to the patient 41

Anaphylactic shock: It rarely occurs but it is life threatening it is characterized by:

- 1- Profound fall in blood pressure
- 2- Dyspnea & respiratory embarrassment
- 3- Facial & laryngeal edema
- 4- Loss of consciousness.
- 5- Urticaria.

It is life threatening condition since it causes air way obstruction in association with laryngeal edema .unless treatment is initiated immediately; the condition may progress to fatal termination.

#### Management of anaphylaxis:

-If you suspect the anaphylactic reaction occurring, immediately seek medical help. The treatment must begin before BP & breathing problem become life threatening. - epinephrine is the most imp medication for the treatment of anaphylaxis; it is inj into a muscle. Epinephrine works rapidly to make blood vessel contract, preventing them from leaking more fluid. It also relax air way, helping the individual breathe easier, relieves cramping in the GIT & stop itching, rapidly reverse the life threatening symptom of anaphylactic shock, if given in time.

- Even if the individual responds to the epinephrine, it is vitally imp to go to an emergency

room immediately.

- 02 & medication may be given to improve breathing.
- IV fluid may be necessary to restore adequate BP.

-additional medication (anti histamine) to contract the effect of histamine & to help in prevention of delayed allergic reaction. -If the victim stops breathing, perform cardio pulmonary resuscitation (CPR) immediately. Continue CPR until the person begins breathing again.

### 5- Over dosage & toxicity

It is relatively rare ,a toxicity reaction can occur when the conc. of LA in circulation increases too rapidly within a short period of time e.g. inj too rapidly into highly vascular area or IV inj .The toxic effect is primarily directed to the CNS & CVS because these tissues are composed of excitable membrane.

Mild over dose sign & symptom:

Restlessness, retention of consciousness talkativeness, & agitation, which may end into convulsion along with increased BP, heart rate, respiratory rate & classified according to the onset.

1- Slow onset  $\leq 5$  min following administration of LA (the possible cause rapid absorption & too large total dose).

2- Slower onset >= 15 min following administration of LA (the possible causes are abnormal biotransformation & renal dis.).

We use the following protocol in management of such cases:

Step 1 reassure the patient that every thing will be all right.

Step 2 administer 02 via nasal cannula or nasal hood.

*Step 3* monitor vital signs.

*Step 4* administer anti convulsants. IV infusion (for diazepam 5mg/ min, for midazolam 1 mg/min).

*Step 5* permits the patient to recover as long as necessary; the patient can leave the dental office escorted or unescorted if we believe that full recovery has occurred, if the doubt exist regarding the level of recovery then .

*Step 6* seeks medical help.

*Step* **7** after termination of reaction be sure that the patient be examined by physician to determine the cause of this reaction.

### Severe overdose reaction:

Unconsciousness with or without convulsion, vascular collapse, coma, cardiac arrest & respiratory arrest & classified into:

1-Rapid onset (within 1 min) the possible cause is IV inj.

2-slow onset (5 to 15 min) (the possible causes are too large dose, rapid absorption & abnormal biotransformation. We use the following protocol in management of such cases:

Step 1 terminate dental treatment.

Step 2 if convulsion is present, protect the patient arm, legs & head & loosen the tight clothing and immediately seek emergency medical assistance.

*Step 3* provide basic life support (maintenance of adequate air way & oxygenation are of utmost importance during management of LA induced seizures .

*Note:* seizures cause increase 02 utilization & hyper metabolism, with increased production of C02 & lactic acid & lead to acidosis this in turn lower the seizer threshold & prolongation of reaction. Cerebral blood flow during the seizure also increase this lead to elevation of anesthetic blood level within the CNS.

*Step 4* administer anticonvulsant

# **BIBLIOGRAPHY**

1. Archer WH. A Manual of Dental Anesthesia. An Illustrated Guide for Students and Practitioners. 2nd ed. WB Saunders Company 1958.

2. Howe GL, Whitehead FIH. Local Anesthesia in Dentistry. A Dental Practitioner Handbook. 3rd ed. Varghese Publishing House 1997.

- 3. Malamed SF. Handbook of Local Anesthesia. 4th ed. New Delhi: Harcourt India Private Limited (Mosby) Indian Reprint 2001.
- 4. Nevin, Mendel, Nevin HR. Problems in Dental Local Anesthesia. Brooklyn, NY: Dental Items of Interest Publishing Company Inc 1954.
- 5. Nevin, Mendel, Puterbaugh PG. Conduction, Infiltration and General Anesthesia in Dentistry. Brooklyn, NY: Dental Items of Interest Publishing Company Inc 1948.
- 6. Richard Bennett C. Monheim's Local Anesthesia and Pain Control in Dental Practice. 7th ed. New Delhi: CBS Publishers and Distributors 1990.



# Extraoral and Intraoral Examination

Oral Surgery Third Year College of Dentistry University of Anbar

Dr. Elham H. Abdulkareem

### History Taking and Clinical Examination of Patients on a Dental Clinic

#### History taking: This should include the following:

- Demographic data
- Presenting complaint (the presenting problem, ideally recorded in the patient's own words)
- History of presenting complaint (a detailed history of the complaint and the symptoms associated with it recorded)
- Dental history
- Medical history (current and past)
- Social and family history

#### **Obtaining information**

The initial step in any patient contact is to confirm their identity. In addition, the clinician should introduce themselves, in order that the patient knows to whom they are talking, which may help to lessen the patient's anxiety. It is also important to ensure that the person with the presenting problem is allowed to speak for themselves.

#### **Demographic data**

Demographic data are necessary to identify the correct patient and corresponding clinical record. The minimum data required include the following:

- Title (Mr, Mrs, Ms, Miss, etc.)
- Name (both forename and surname)
- Marital status
- Date of birth
- Sex
- Occupation
- Current address and contact telephone numbers

Name and contact details of the patient's general medical/dental practitioner. The data obtained must be recorded accurately. Some conditions are correlated with

age, sex, ethnicity or occupation, and demographic data may help to diagnose a presenting condition more easily, for example a heavy metal worker with dark blue staining at the gingival margins.

#### **Preparation for Examination**

- Review the patient's histories
- Examine radiographs
- Patient understanding
- Cultural sensitivity

#### **History taking**

History taking is a skill that requires practice. Patients respond in different ways to similar lines of questioning, and it may be necessary to modify questioning style or to ask the same question several times but in different ways in order to optimise the information obtained.

#### **Components of the history**

#### Presenting complaint and history of presenting complaint

Patients attending a DC often present complaining of pain. In order to reach a reliable differential diagnosis or diagnosis, it is important to obtain a clear description of the pain. Where possible, use open questions and avoid prompting the patient; for example, 'What does the pain feel like?' is preferable to 'Is the pain sharp or dull?' which restricts and influences the patient response. Once a description of the pain obtained, more specific questions can be used to develop the history further. Basic questions that should be asked are summarised below:

- Site of pain (ask the patient to point with one finger to the place of maximum pain).
- Ask the patient if the pain radiates anywhere.
- Get the patient to describe the pain, e.g. dull ache, sharp, throbbing or shooting.
- Is the pain intermittent or constant? How frequent is it?
- Onset gradual or sudden?
- Is there anything that makes the pain better or worse?
- What treatments has the patient tried and were they effective?
- Does the pain keep the patient awake at night or wake them from sleep?

- Is the pain affected or initiated by hot or cold stimuli?
- Are there any associated symptoms such as swelling, numbness or pain?

#### Past and current medical history

The past and current medical history is a description of previous and current medical issues. A systematic approach to data collection is required. It is important to go through the collected information carefully with the patient in order to identify any areas of confusion. The medical history should be carefully recorded.

- Cardiovascular: Myocardial infarction, angina, hypertension
- Respiratory: Asthma, chronic obstructive pulmonary disease.
- Gastrointestinal: Peptic ulceration,
- Hepatic: hepatitis
- Haematological: Blood borne viruses, e.g. hepatitis A/B/C, HIV, clotting disorders, leukaemia, anaemia
- Neurological: Epilepsy, cerebrovascular disease, psychological/psychiatric disorders
- Musculoskeletal: Muscular dystrophy, joint replacements
- Genitourinary and renal: Genitourinary infection, renal disease or failure, renal transplant, etc.
- Drug history: Prescribed, non-prescribed,
- Allergy: Drugs, latex.
- Past hospitalisation: Medical/surgical
- Social history: Occupation
- Smoking habits: duration, frequency and type
- Alcohol consumption: Quantity
- Home and family circumstances

The degree of severity of a medical problem can be judged by asking further questions. For example, in patients with epilepsy, the frequency and severity of the problem should be ascertained using key questions such as: What medications do you take? What symptoms do you get prior to a seizure? When was your last seizure? Have you been treated in an accident and emergency unit or admitted to a hospital following a seizure? If the clinician has this information, a simple risk assessment can be made, and in the event of a seizure, emergency care is facilitated and a specialist's help more easily obtained if required.
A thorough medical history may uncover conditions that are relevant to diagnosis of the presenting complaint, for example oral ulceration in a patient taking the potassium channel activator, or to the subsequent management of the patient, such as alcoholic cirrhosis in a patient requiring extractions. Clearly, patients taking the anticoagulant warfarin require special consideration. It is important to remember that warfarin interacts with patients who have a high alcohol intake may also have problems with coagulation. In such cases, if surgical treatment is considered, blood should be taken for a full blood count (principally to check the platelet count) and a clotting screen.

#### **Patient examination**

The examination should start from the moment the patient walks into the dental clinic. The clinician should remember to examine both the normal and affected sides. The examination is divided into an extra-oral examination, followed by intra-oral examination.

#### **Extra-oral examination**

The extra-oral examination starts with a visual examination of the head and neck with particular note made of swellings or deformity, asymmetry of the face, abnormal colour or scars on the skin or lips. A clinical photograph is a useful way of recording information.

#### **Sequence of examination**

- 1. Face, Lips, Skin and Eyes
- 2. Lymph Nodes: a. Occipital b. Auricular c. Superficial Cervical d. Deep Cervical e. Submental f. Submandibular
- 3. Glands: a. Parotid b. Submandibular c. Sublingual d. Thyroid (Figures 3 and 4)
- 4. Larynx
- 5. Hyoid bone
- 6. Muscles: a. Masseter b. Temporalis c. Sternocleidomastoid d. Mylohyoid

#### 7. Temporomandibular Joint: a. Right b. Left: Figure 2

Inspection:

- Any swelling over the joint region
- Symmetry and extent of movement
- Clicking sound

Palpation: Pre-auricular and intra-auricular

- Extent and symmetry of movement
- Temperature of the overlying skin
- Tenderness Crepitus

Auscultation: Clicking

**Inspection:** It does detect any unusual changes in the oral cavities, as it's based on vision.

- Color changes "pigmentation or caries"
- ✤ Tooth Fracture and different Lesions.

**Palpation:** This depends on our sense of touch to feel any abnormalities and to differentiate it from the normal. The types are Bimanual, Bilateral We can detect the different consistency, temperature, Mobility, in duration.

**Percussion:** We examine this by striking an object on the tooth and evaluate the produced sound. This technique also helps in detection of tooth mobility.

**Probing:** This is important technique as it can detect caries and any periodontal diseases.

Auscultation: listen to the normal sounds produced by the Patient

- Wheezing: Respiratory disease
- TMJ clicking: TMJ disorder

**Odor:** smelling the patient oral odor can help in the differential diagnosis. Such as Acetone odor= Uncontrolled DM

#### **Intraoral examination**

Lips & intraoral mucosa

- View/palpate lips, labial and buccal mucosa, and mucobuccal folds.
- Examine and palpate the tongue
- Mucosa of the floor of the mouth.
- Hard and soft palates, tonsillar areas, and pharynx
- Use a mirror

Oropharynx, nasopharynx, and larynx.

Note: amount and consistency of the saliva and evidence of dry mouth (xerostomia).



Figure 1: Bimanual Palpation. A: Examination of the buccal mucosa by simultaneous palpation on extraorally and intraorally. B: Examination of the floor of the mouth by simultaneous palpation with fingers of each hand in apposition.





Figure 2: Assessment of the Temporomandibular Joint.



Figure 3: Lymph Nodes.



Figure 4: Lymph Node examination.



#### References:

- 1. https://www.DrmanaliR/oral-diagnostic-process-part-2
- 2. https://www. indiandentalacademy/extra-oral-examination

3. https://www.JaceySheckler/extraoral-and-intraoral-examination

4. https://www.slideshare.net/ozident/intra-oral-examination



# Definition

Vasoconstrictors are the chemical agents or adjuncts added to local anesthetic solutions

- (a) to oppose vasodilatation caused by these agents and
- (b) to achieve hemostasis.

### ACTIONS

The addition of a vasoconstrictor to a local anesthetic agent causes constriction of blood vessels and thereby controls tissue perfusion. The effects caused by addition of vasoconstrictors to local anesthetic agents are:

1. It decreases the blood flow to the site of injection, because of vasoconstriction.

2. It decreases the rate of absorption of local anesthetic agent into cardiovascular system.

3. It lowers the plasma level of local anesthetic agent (Cannall et al, 1975) and Wildsmith et al, 1977), thereby, decreasing the risk of systemic toxicity of local anesthetic agent.

4. Higher volumes of local anesthetic agent remain in and around the nerve for longer periods; thereby increasing the duration of action of most local anesthetic agents (Brown, 1968).

5. It decreases bleeding at the site of injection because of decreased perfusion. This is useful when increased bleeding is expected during a surgical procedure (Carpenter et al, 1989; and Myers and Heckman, (1989).

# **CLASSIFICATIONS**

These drugs can be classified on the basis of mode of action, into the following three categories.

1. Direct acting drugs: These drugs stimulate or exert their action directly on the adrenergic receptors. For example, epinephrine, norepinephrine, dopamine, levonordefrine and isoproterenol, etc.

2. Indirect acting drugs: These drugs act by releasing norepinephrine from the adrenergic nerve terminals (from their intraneuronal storage sites), e.g. tyramine, amphetamine, methamphetamine, hydroxyamphetamine.

**3. Mixed acting drugs: These drugs have both direct and indirect actions. For example, metaraminol and ephedrine.** 

# **ADRENERGIC RECEPTORS**

### Types

We have 2 types of adrenergic receptors based on their action on smooth muscles.

### $\alpha$ Receptors

Activation of receptors results in vasoconstriction.

## β*Receptors*

 $\beta$  receptors are subcategorised further into  $\beta$ 1,  $\beta$ 2 and  $\beta$ 3 receptors.

1. β1 receptors are found in the heart and small intestine; and are responsible for: (i) cardiac stimulation (increase in the HR and increase in the strength of contraction) and (ii) renin release.

2.  $\beta$ 2 receptors are found in bronchi, vascular beds, and uterus; and are responsible for:

i. Relaxation of bronchial muscles resulting in bronchodilatation

ii. Relaxation of muscles in the walls of blood vessels, resulting in vasodilatation, and iii. Relaxation of uterus (Lands et al, 1967).

3.  $\beta$ 3 receptors are found in brown and white adipose tissue; and are responsible for lipolysis.

### **DILUTION OF VASOCONSTRICTORS**

The dilution of vasoconstrictors is commonly referred to as a ratio (e.g. 1 to 1000; and is written as 1:1000).

• 1:1000 means 1 g or 1000 mg of solute (drug) contained in 1000 ml of solution. Therefore, a 1:1000 dilution contains 1000 mg in 1000 ml or 1.0 mg/ml of solution.

Vasoconstrictors used in local anesthetic solutions in dentistry are more diluted than 1:1000. Hence, a solution of a dilution 1:10,000 will contain 0.1 mg/ml of solution and 1:100,000 dilution will contain 0.01 mg/ml of solution.

### Epinephrine

remains the most commonly used and the most effective vasoconstrictor used in medicine and dentistry.

- **Proprietary Name: Adrenaline.**
- Source: It is secreted primarily by adrenal medulla. It is available as a synthetic and is also obtained from the adrenal medulla of animals.

Mode of action: It acts directly on both  $\alpha$  and  $\beta$  adrenergic receptors;  $\beta$  effects predominate.

It must be borne in mind that the endogenous release of adrenaline, caused by the stress of going to a Dental Surgeon, sight of dental instruments, and the apprehension of pain caused by the needle-sticks produces effects similar to those produced by excessive exogenous administration of adrenaline.

# Maximum Recommended Doses (MRD) of Epinephrine

This drug is potent; and can produce undesirable results

- (a) if used in large volumes; or
- (b) (b) if inadvertently injected intravascularly. Hence, these drugs should be used with due consideration to their benefits and risks. It is recommended that the least concentrated solution that produces effective pain control should be used.
- (c) Lidocaine is available with three dilutions of epinephrine (1:80,000, 1:100,000 and 1:200,000), in many other countries.
- (d) recommended that the maximum dose of epinephrine should be limited to 0.2 mg per appointment. The American Heart Association has recommended restriction of epinephrine in local anesthetics when administered to patients with ischemic heart diseases.

A. Normal healthy adult patients (0.2 mg per appointment)

- 10 ml of a 1: 50,000 dilution (5 cartridges)
- 16 ml of a 1: 80,000 dilution (8 cartridges)
- 20 ml of a 1: 100,000 dilution (10 cartridges)
- 40 ml of a 1: 200,000 dilution (20 cartridges)

B. Patients with clinically significant cardiovascular disease (0.04 mg per appointment) (approximately 1/5th of the dose for normal patients)

2 ml of a 1: 50,000 dilution (1 cartridge)

3.2 ml of a 1: 80,000 dilution (1.6 cartridges)

4 ml of a 1: 100,000 dilution (2 cartridges)

8 ml of a 1: 200,000 dilution (4 cartridges)

### **Medically Compromised Individuals**

The Maximum Recommended Doses (MRD) is 0.04 mg/appointment.

#### **For Hemostasis**

Epinephrine is used in local anesthetic solution to minimise bleeding by depositing at the site of surgery via infiltration anesthesia during various surgical procedures. A local anesthetic solution containing 1:50,000 dilution of epinephrine is more effective in minimising bleeding than more dilute solutions such as 1:100,000 or 1:200,000.

**Systemic Actions** 

#### **Cardiovascular System**

The overall actions of epinephrine on the heart and cardiovascular system are the result of direct stimulation. If the stimulation of the heart persists for a long time, there will subsequently, be overall decrease in cardiac efficiency. These actions are as follows:

- i. Increased stroke volume
- ii. Increased heart rate
- iii. Increased cardiac output
- iv. Increased strength of contraction
- v. Increased systolic and diastolic pressures
- vi. Increased myocardial oxygen consume

#### **Blood Vessels**

- 1. The primary action of epinephrine is on microcirculation (small arterioles and precapillary sphincters).
- 2. The blood vessels which supply skin, mucous membranes and kidneys contain primarily  $\alpha$  receptors.
- 3. Epinephrine produces constriction of these blood vessels. The blood vessels supplying skeletal muscles contain both  $\alpha$  and  $\beta$ 2 receptors, with  $\beta$ 2 receptors predominating.
- 4. Epinephrine in small doses produces dilatation of these vessels as a result of β2 effect.
- 5.  $\beta$ 2 receptors are more sensitive to epinephrine than  $\alpha$  receptors.
- 6. Large doses of epinephrine produce vasoconstriction because of stimulation of  $\alpha$  receptors.

### **Respiratory System**

It has a  $\langle 2 \text{ effect on the smooth muscle of the bronchioles, and hence is a potent dilator. It is a drug of choice in cases of bronchospasm as in bronchial asthma.$ 

### Central Nervous System

In therapeutic doses, epinephrine does not stimulate central nervous system. It stimulates central nervous system only if excessive doses are administered.

### Metabolism

The actions of epinephrine are terminated by one of the following methods:

i. Primarily, by its reuptake by adrenergic nerves.

ii. Inactivated by the enzymes catechol-O-aminotransferase (COMT),

and monoamine oxidase (MAO), both of which are present in liver.

iii. Very small amounts are excreted unchanged in urine.

#### Overdose

- In small amounts commonly used in dentistry, only the arterioles in the
- immediate area of injection are affected by vasoconstrictors.
- The factors that may produce toxic manifestations are:
- (i) large volumes, (ii) high concentration of vasoconstrictors;
- (ii) and (iii) inadvertent vascular injection of comparatively small amounts of vasoconstrictors. The manifestations can be in the form of tachycardia, hypertension, palpitation, headache, tremors, pallor, and in rare cases, ventricular fibrillation.
- The systemic effects of vasoconstrictors usually last as long as the blood levels of the drug remain elevated.

### **Clinical Applications in Dentistry**

It is used along with a local anesthetic agent, as a vasoconstrictor:

- (i) for achieving hemostasis,
- (ii) to decrease the absorption of local anesthetic agent into cardiovascular system,
- (iii) to prolong duration of action.

### **Availability in Dentistry**

Epinephrine is available in the following dilutions: 1:80,000, 1:100,000 and 1:200,000 for use in dentistry.

### **Norepinephrine (Nor-adrenaline)**

It is not commonly used in dentistry.

### Felypressin

It is available as a vasoconstrictor, in combination with prilocaine, in dental

- local anesthetic cartridges, in many countries (Citanest forte). *Proprietory name:* Octapressin
- *Chemical structure:* It is a non-sympathomimetic amine; and a synthetic analogue of vasopressin.
- *Mode of action:* It acts by directly stimulating vascular smooth muscle. It has little direct effect on the heart or on adrenergic nerve transmission. Its actions are more pronounced on venous rather than on arteriolar microcirculation (Altura, 1965).

### Systemic actions:

i. *Myocardium:* No direct effects.

ii. *Coronary arteries:* Doses greater than therapeutic doses, may impair blood flow through coronary arteries.

iii. *Vasculature:* Doses greater than therapeutic doses, cause constriction of cutaneous blood vessels producing facial pallor.

iv. *Uterus:* It has both antidiuretic and oxytocic actions; hence the latter action contraindicates its use in pregnant patients.

v. *Maximum doses:* Patients with clinically significant cardiovascular impairment: Maximum Recommended Dose: 0.27 IU (international units) (that is 9 ml of 0.03 IU/ml or 2 cartridges).

**vi.** *Side effects and overdose:* Studies have shown a wide margin of safety with felypressin (Klingenstrom et al, 1967).

**vii.** *Clinical applications:* Used as vasoconstrictor with local anesthetic to decrease their absorption and increase their duration of action.

**viii.** *Availability:* It is used in dentistry in a dilution of 0.03 IU (international units)/ml with 3% prilocaine.

**ix.** *Disadvantage:* It acts primarily on venous circulation, hence felypressin is not as effective as conventional vasoconstrictors in providing hemostasis during surgical procedures (McClymont and Crowther, 1988).

### x. Indications:

 a. It may be safely used in patients with medical problems which contraindicate the administration of epinephrine (Anderson and Reagan 1993), which include:

(i) mild to moderate cardiovascular diseases, including hypertension, and (ii) other non-cardiovascular diseases; such as hyperthyroidism.

b. It can be used in patients who are on antidepressant drugs such as tri- or tetracyclic acid antidepressants or monoamine-oxidase inhibitors.

### xi. Contraindications:

a. Local anesthetic containing felypressin is not recommended for use when hemostasis is required because of their predominant effect on venous rather than arteriolar circulation (Newcomb and Wait, 1972).

b. Pregnancy: It is not recommended for use during pregnancy because of its oxytocic action on Uterus.

### SELECTION OF A VASOCONSTRICTOR

The selection of an appropriate vasoconstrictor is based on following factors:

- 1. Length of the surgical or dental procedure
- 2. Requirement for hemostasis during the surgical procedure
- 3. Requirement for postoperative pain control
- 4. Medical or physical status of the patient and concurrent medications taken.

### Length of the Surgical or Dental Procedure

The duration of pulpal and hard tissue anesthesia with 2% lidocaine only lasts for about 10 min.

The addition of epinephrine of 1:50,000 or 1:100,000 dilution prolongs the duration to about 60 min. Hence, for any oral surgical or dental restorative procedure requiring 40-50 minutes, it is difficult to achieve consistent pulpal anesthesia without inclusion of a vasoconstrictor.

#### **Requirement of Hemostasis**

Some of the vasoconstrictors are effective in minimising blood loss during surgical procedures. However, most of vasoconstrictors also produce vasodilatation as a result of decline in the tissue levels of epinephrine, which is known as rebound phenomenon.

While using epinephrine as a vasoconstrictor for hemostasis, we must consider that epinephrine is both  $\langle$  and  $\mathbb{R}$  agonist. It possesses both  $\alpha$  and  $\beta$  actions; and produces vasoconstriction through it's effect.

However, felypressin stimulates venous circulation more than the arteriolar circulation, and therefore, is of minimal value in achieving hemostasis (Altura et al, 1965).

### **Requirement for Postoperative Pain Control**

Profound pain control of adequate duration by a local anesthetic agent with a vasopressor is used. Plain local anesthetic agent produces pulpal anesthesia of shorter duration that a local anesthetic agent with a vasopressor and is likely to produce stress response.

#### Medical or Physical Status of the Patient and Medications Concurrently Taken

The benefits and risks of including a vasoconstrictor in a local anesthetic solution in patients who are medically compromised, must be weighed against benefits and risks of using plain local anesthetic solution (Goulet et al, 1992).

Once the medical status of the patient is improved or corrected, dental or surgical procedures requiring administration of local anesthetic agents with vasoconstrictors are indicated.

### The groups where inclusion of vasoconstrictor is contraindicated are

### given below.

1. Patients with significant cardiovascular disease such as (a) ischemic heart disease, (b) hypertension, and (c) cerebral strokes.

2. Patients with certain uncontrolled non-cardiovascular diseases, such as: thyrotoxicosis or hyperthyroid states, and diabetes mellitus.

3. Patients receiving non-specific beta-blockers, monoamine-oxidase inhibitors (MAOIs), tricyclic antidepressants (TCAs) and phenothiazine's.

- 4. Patients with sulphite sensitivity.
- 5. Patients who are undergoing general anesthesia with halogenated agents

6. Pregnancy.

1. Patients with significant cardiovascular disease such as (a) ischemic heart disease, (b) hypertension, and (c) cerebral strokes. Epinephrine and other vasoconstrictors can be used in moderate amounts in patients with mild to moderate cardiovascular disease.

Felypressin has minimal cardiothoracic stimulatory actions and is nondysrhythmogenic; it is recommended for patients with significant cardiovascular disease. Patients with severe cardiovascular diseases are at too great a risk for elective dental therapy, such as:

a. Ischemic heart disease

i. Patients with history of acute myocardial infarction within last six months.

ii. Patients with history of acute anginal episodes on daily basis or where signs and symptoms are increasing in severity (angina pectoris) (pre-infarction stage or unstable angina).

iii. Patients with cardiac dysrhythmias despite appropriate antiarrhythmic drug therapy (Goulet et al, 1992).

iv. Post-coronary artery bypass surgery, less than six months.

b. Hypertension: Patients with systolic BP greater than 200 mm of Hg and diastolic BP greater than 110 mm of Hg should not receive any dental care unless BP is corrected.

c. Cerebral strokes: Patients with history of less than six months after cerebrovascular accident.

Epinephrine and other vasoconstrictors can be used in moderate amount in patients with mild to moderate cardiovascular disease. Felypressin has minimal cardiovascular stimulatory actions and is non-dysrhythmogenic; it is recommended for significant cardiovascular risk patients.

2. Patients with certain uncontrolled non-cardiovascular diseases, such as: thyrotoxicosis or hyperthyroid states and diabetes mellitus. Medical treatment of the conditions and a consent from the treating physician is

necessary to use vasoconstrictor in such conditions.

a. Thyrotoxicosis or hyperthyroid states: Epinephrine is contraindicated in patients with clinical evidence of hyperthyroid states (Goulet et al, 1992). The signs and symptoms include: Exophthalmos, tremors, hyperhydrosis, increased basal metabolic rate, irritability and nervousness, increased body temperature, inability to tolerate heat, increased HR, and increased BP.

b. Diabetes mellitus: Diabetes is not only a metabolic, and endocrinal disease, but also a vascular one. Microcirculatory changes will result in impaired blood flow to the tissues. The inclusion of a vasoconstrictor in local anesthetic solution may further compromise the inadequate blood supply, and result in local ischemia and tissue sloughing.  Patients receiving non-specific beta-blockers, monoamine-oxidase inhibitors (MAOI), tricyclic and tetracyclic antidepressants and phenothiazines for psychiatric ailments.

a. Patients taking monoamine-oxidase inhibitors (MAOI) are not at an increased risk provided the doses used are within the recommended range (Goulet et al, 1992; and Verrill, 1975).

b. Patients receiving tricyclic and tetracyclic antidepressants: Such patients are at a risk of developing dysrhythmias with administration of epinephrine. Hence, its dose should be minimal. The administration of levonordefrin or norepinephrine in patients receiving tricyclic

antidepressants is absolutely contraindicated. Large doses may induce severe (exaggerated) response. Felypressin has minimal cardiothoracic stimulatory actions and is nondysrhythmogenic, and hence can be used. In all the abovementioned conditions felypressin can safely be used.

4. Patients with sulfite sensitivity: Local anesthetic solutions contain an antioxidant, which retards oxidation of the vasoconstrictor. The most commonly used antioxidant in dental cartridges is sodium bisulfite.

5. Patients who are undergoing general anesthesia with halogenated agents: Epinephrine should not be used as a vasoconstrictor during GA when a patient is receiving an inhalational halogenated anesthetic agents such as halothane, methoxyflurane, and ethrane. These vasoconstrictors cause cardiac dysrhythmias. In such situations, felypressin is recommended.

6. Pregnancy: Felypressin is not indicated in pregnant patients because of its potential oxytocic actions.

#### LOCAL ANESTHETIC CARTRIDGES AND VIALS

It contains primarily the local anesthetic drug, and also the other ingredients, which are as follows:

- 1. Local anesthetic drug
- 2. Vasopressor/vasoconstrictor drug
- 3. Preservative for vasopressor
- 4. Sodium chloride (NaCl) or Ringer's solution
- 5. Distilled water
- 6. General preservatives.

### PRESERVATIVE FOR VASOPRESSOR

Local anesthetic solutions containing vasoconstrictors also contain a specific agent, an antioxidant, that acts as a preservative for vasoconstrictors.

- The most frequently used antioxidant is sodium-bisulfite or sodium metabisulfite. It prevents biodegradation of vasopressor by O<sub>2</sub> which might be present in the cartridge either introduced during manufacture, or which got diffused through semipermeable membrane or the rubber diaphragm after filling at the time of storage of the cartridge.
- Sodium-bisulfite reacts with O<sub>2</sub> before O<sub>2</sub> can destroy vasopressor. Sodium-bisulfite is oxidised to sodium-bisulphate, a chemical with even lower pH (acidic), than before oxidation.
- Local anesthetic solutions without vasoconstrictors have a shelf-life of about 48 months.
- Local anesthetic solutions containing vasoconstrictors have their shelf-life reduced to 18 and 12 months for epinephrine and phenylephrine; and norepinephrine and levonordefrine, respectively. This is so because of the instability of the vasoconstrictors. Hence fresh solutions produce better analgesia and cause less irritation to the tissues.
- Allergy to bisulfites must be considered in pre-anesthetic medical evaluation.

### SODIUM CHLORIDE (NaCl) OR RINGER'S SOLUTION

- 1. It is added to the contents of dental cartridge to make the solution isotonic with the tissues of the body.
- 2. Hypertonic solution produces tissue edema, paresthesia, sometime lasting for several months following drug administration.

### **DISTILLED WATER**

It is used as a diluent to provide the volume of solution in the dental cartridge.

### **GENERAL PRESERVATIVES**

A number of chemicals are used as general preservatives.

These are added to increase the shelf-life; and include:

- (i) Methylparaben, (ii) Thymol,
- (ii) i. Methylparaben: It is a bacteristatic and fungistatic agent. It has been excluded from cartridges in USA, from 1st January 1984, following reports of allergy. However, it is still used in multidose vials in some countries including India.
- ii. Thymol: It is antiseptic, fungistatic, and antihelminthic.

# Surgical Anatomy in Relation to Local Anaesthesia by Dr. Elham Hazeim Abdulkareem College of Dentistry University of Anbar

# The trigeminal nerve contains a large number of sensory (afferent) and motor (efferent) neurons.

• The trigeminal nerve has a wide innervation area.

- The nerve provides the sensitivity of the dentition, the mucosa of the mouth, nose and paranasal sinuses and the facial skin.
- The nerve also contains motor fibres that innervate, among others, the masticatory muscles.



- The trigeminal ganglion is formed by the aggregation of cell bodies of sensory neurons.
- After the ganglion, three branches of the trigeminal nerve can be distinguished: the ophthalmic nerve (n. V1), the maxillary nerve (n. V2) and the mandibular nerve (n. V3).
- The motor root joins the mandibular nerve only, once it has exited the skull via the foramen ovale.

- The sensory areas covered by the three main branches are generally as follows:
- The ophthalmic nerve carries sensory information from the skin of the forehead, the upper eyelids and the nose ridge and the mucosa of the nasal septum and some paranasal sinuses.
- The maxillary nerve innervates the skin of the middle facial area, the side of the nose and the lower eyelids, the maxillary dentition, the mucosa of the upper lip, the palate, the nasal conchae and the maxillary sinus.
- The mandibular nerve innervates the skin of the lower facial area, the mandibular dentition, the mucosa of the lower lip, cheeks and floor of the mouth, part of the tongue and part of the external ear.

#### **Ophthalmic Nerve**

The ophthalmic nerve (n. V1) enters the cranial part of the orbit.

This nerve carries only sensory fibres, and, just before leaving the cranial cavity through the superior orbital fissure, it branches off into three:

the nasociliary, frontal and lacrimal nerves.

These nerves run in the roof of the orbit and are involved in the sensory innervation of a large number of structures, such as the mucosa of the frontal,

sphenoid and ethmoid sinus,

the mucosa of the nasal cavity and

nasal septum,

the skin of the nose ridge,

the upper eyelid and forehead till the hairline and the mucosa that covers the eyeball and inside of the eyelids.


- The maxillary nerve (n. V2), too, is solely sensory.
- It enters the pterygopalatine fossa via the foramen rotundum Through the inferior orbital fissure, it reaches the floor of the orbit and proceeds there as the infraorbital nerve, first in the infraorbital sulcus and then in the infraorbital canal. It then reaches the face via the infraorbital foramen.
- Within the pterygopalatine fossa, the maxillary nerve is connected via a number of branches to the upper side of the parasympathetic pterygopalatine ganglion.
- Sensory fibres run through these branches which exit on the lower side of the ganglion and form, among others, the following nerves:





- The nasal nerves and nasopalatine nerve that run through the sphenopalatine foramen to the nasal mucosa.
- The nasal nerves innervate the back part of the nasal mucosa.
- The nasopalatine nerve, which runs forwards along the nasal septum and reaches the oral cavity through the incisive canal, innervates the mucosa and bone of the nasal septum, frontal two-thirds of the palate and the palatal gingiva of the maxillary teeth.
- The greater palatine nerve that runs via the greater palatine canal to the mucosa of the hard palate where it subsequently innervates the palatal gingiva of the maxillary alveolar process and the pulp of the palatal first molar and premolar.

- The lesser palatine nerves that run to the mucosa of the soft palate via the lesser palatine canals Together with the palatine nerves, there are also parasympathetic and orthosympathetic fibres that run from the pterygopalatine ganglion to the salivary glands in the palatal mucosa
- In the pterygopalatine fossa, the maxillary nerve also branches into the posterior superior alveolar nerve and the zygomatic nerve
- The posterior superior alveolar nerve exits the pterygopalatine fossa through the pterygomaxillary fissure and runs over the maxillary tuberosity.
- The nerve divides into a large number of little branches, the posterior superior alveolar rami, which enter the wall of the maxilla through small openings and innervate the maxillary molars and corresponding buccal gingiva.



- The zygomatic nerve arrives in the orbit via the inferior orbital fissure and branches into the zygomaticotemporal and zygomaticofacial nerves.
- These exit the lateral orbital wall through small canals in the zygomatic bone and innervate the skin above.
- The zygomatic nerve also contains postganglionic parasympathetic fibres that come from the pterygopalatine ganglion and that join the lacrimal nerve for the lacrimal gland.

- As it runs along the orbital floor, the infraorbital nerve branches in two:
- the middle superior alveolar nerve, for the innervation of the maxillary premolars and the corresponding buccal gingiva,
- and the anterior superior alveolar nerve, for the maxillary canine and incisors and the corresponding buccal gingiva.
- These nerves usually run between the mucosa and outer wall of the maxillary sinus. There they divide into a number of small branches, the medial and anterior superior alveolar that penetrate into the maxillary alveolar process via small openings.
- Inside the bone, they form together with the posterior superior alveolar, right above the apices, an extensive nervous network – the superior alveolar plexus – from which short little branches are sent to the dentition and gingiva.

- Once the infraorbital nerve reaches the face via the infraorbital foramen, it splits into a large number of branches for the sensory innervation of
- the skin of the lower eyelid,
- the infraorbital region,
- the side of the nose and the skin and
- mucosa of the upper lip

- The mandibular nerve (n. V3) contains both sensory and motor fibres. This nerve exits the skull through the foramen ovale and ends in the infratemporal fossa
- The mandibular nerve is located just below this foramen between the lateral pterygoid muscle and the tensor veli palatini muscle. The nerve sends a motor branch to the latter muscle
- The mandibular nerve splits into two main branches:

# The anterior and posterior trunks

• from the anterior trunk, a sensory nerve emerges, the buccal nerve and a number of motor nerves, i.e. the pterygoid nerves, the deep temporal nerves and the masseteric nerve.

# Three branches emerge from the posterior trunk:

- the auriculotemporal nerve (sensory)
- the lingual nerve (sensory)
- the inferior alveolar nerve (mixed sensory and motor)

deep temporal nerves mandibular nerve posterior superior alveolar nerve auriculotemporal nerve maxillary artery middle meningeal artery lateral pterygoid chorda tympani muscle facial nerve buccal nerve inferior alveolar nerve medial pterygoid muscle mylohyoid nerve lingual nerve submandibular ganglion sublingual gland submandibular gland mental nerve hypoglossal nerve

- The buccal nerve runs along the medial surface of the upper head of the lateral pterygoid muscle and moves laterally between the muscle. The nerve contains fibres for the salivary glands in the buccal mucosa. The nerve innervates the skin and mucosa of the cheek and the buccal gingiva of the mandibular alveolar process at the level of the molars and premolars.
- The pterygoid nerves are short motor branches for the medial and lateral pterygoid muscles.
- The masseteric nerve runs laterally along the top of the upper head of the lateral pterygoid muscle and reaches the deep surface of the masseter muscle via the mandibular notch.
- The deep temporal nerves also run high along the lateral pterygoid muscle and penetrate the medial side of the temporal muscle.
- The auriculotemporal nerve arises as two roots encircling the middle meningeal artery. After the roots have merged to a single nerve, this nerve first runs laterally behind the mandibular neck

 the inferior alveolar nerve contains motor and sensory fibres. It runs deep to the lateral pterygoid muscle. Emerging from beneath this muscle, it directs to the mandibular foramen. Just before it enters the mandibular canal, it gives off its motor mylohyoid branch for the mylohyoid muscle and for the anterior belly of the digastric muscle.



- The nerve, artery and vein travel anteriorly in mandibular canal, as far forward as mental foramen, which is located at a point below and between the roots of the premolars, where the nerve divides into its terminal branches, (i) mental nerve, and (ii) incisive nerve.
- i. Mental nerve: It emerges from the mandibular canal through the mental foramen in the form of a major bulk and divides into three branches that innervate (i) skin of chin, (ii) skin and mucous membrane of lower lip, and (iii) buccal mucosa from the incisor to the premolars. It carries a few secretomotor fibers from chorda tympani to labial minor salivary glands.
- **ii.** Incisive nerve: It is the smaller terminal branch and the continuation of inferior alveolar nerve, within the substance of the body of the mandible, anterior to the mental foramen. It supplies the pulps of anterior teeth, central and lateral incisors, and canine, and sometimes the first bicuspid, supporting alveolar bone, periodontal ligament, and the overlying soft tissues anterior to the mental foramen.

# **Innervation of the Tongue**

- Various nerves are involved in the sensory and motor innervation of the tongue.
- The general sensitivity (pain, touch, temperature) of the anterior two-thirds of the tongue is supplied by the lingual nerve (branch of the n. V3).
- The specific sensitivity (taste) of the anterior two-thirds is supplied by the chorda tympani (branch of the n. VII) that has joined with the lingual nerve.
- The sensitivity, general and specific, of the posterior third of the tongue is supplied by the glossopharyngeal nerve (n. IX). The motor innervation of the tongue takes place through the hypoglossal nerve (n. XII).



- The lingual nerve is joined, directly after its separation of n. V3, by the chorda tympani.
- This is a branch of the facial nerve with preganglionic parasympathetic fibres from the brainstem and sensory nerves for the taste of the anterior two-thirds of the tongue.
- The lingual nerve runs deep to the lateral pterygoid muscle and forwards over the lateral surface of the medial pterygoid muscle. At the level of the apices of the third mandibular molar, it lies immediately beneath the mucosa against the inner side of the mandible.



Fig. 2.9 The sensory innervation of the oral cavity. a The palate, the superior alveolar process, the cheek and the upper lip. b The tongue, the inferior alveolar process, the cheek and the lower lip

- It continues superiorly to the mylohyoid muscle, passing under the submandibular duct, and then ascends in the tongue
- The section of the lingual nerve that comes from n.
  V3 supplies the general sensitivity (pain, touch, temperature) of the anterior two-thirds of the tongue, the mucosa of the floor of the mouth and the lingual gingiva of the inferior alveolar process
- The postganglionic parasympathetic fibres run to the submandibular and sublingual glands

# **METHODS OF LOCAL ANESTHESIA**

- There are several methods of achieving pain control with local anesthetic agents. The various types of techniques used for deposition of these agents, in dentistry are as follows:
- (1) Surface or topical anesthesia,
- (2) Infiltration anesthesia,
- (3) Field block, and
- (4) Nerve block or conduction anesthesia.

The selection of the type of anesthesia depends upon the area and the type of surgery. In general, infiltration anesthesia is adequate for a small isolated area, field block is indicated when two or more teeth are being treated, while a nerve block is indicated for dental or surgical procedure in a quadrant of a jaw.

# **Surface or Topical Anesthesia**

 By this method small terminal nerves in the surface area of the intact mucosa or the skin up to the depth of about 2 mm are anesthetised by application of a local anesthetic agent directly to the area.

# **Nerves Anesthetised**

• Superficial nerve endings.

# Indications

- i. Prior to the infiltration injection techniques or nerve blocks for making the insertion of the needle painless
- ii. Prior to carrying out incision and drainage of abscesses
- iii. Prior to removal of sutures.

# Forms

#### I. Spray

- I. The active ingredient is a suitable local anesthetic agent, such as 10% or 15% lignocaine hydrochloride in water base. The agent is expelled in small quantities from an aerosol container
- II. Advantage: Rapidity of onset. The onset of time of anesthesia is approximately 1 minute and the duration of anesthesia is approximately 10 minutes.
- III. Disadvantage: When used as a spray, the solution is spread over more extensive area than desired.
- IV. Method of application: It is used as a spray on the area in which the needle penetration is proposed to be made; or it can also be sprayed on a small cotton pellet or roll and then placed on the proposed site of injection for about one minute.

**II. Ethyl chloride spray:** It produces anesthesia by refrigeration. When sprayed onto either mucous membrane or skin, it gets volatilised rapidly, and produces rapid anesthesia.

- •Method of application: The spray is directed over a limited area until "snow" appears.
- Care: Undue inhalation of vapours by the patient should be avoided.
- Use: Occasionally used to produce surface anesthesia prior to taking an incision for fluctuating abscesses.

#### Ointment

It is used for similar purposes as spray. The active ingredient is a suitable local anesthetic agent, such as 5% lignocaine hydrochloride.

• Time of onset: 3-4 minutes.

• Application: Ointments are used for application over tender and inflamed gingivae prior to deep scaling.

#### Emulsion

The active ingredient is a suitable local anesthetic agent, such as 2% lignocaine hydrochloride.

• Indications:

i. When full mouth impressions are to be taken in patients who are prone to retching.

ii. Relief of postoperative pain or tenderness following mucogingival surgical procedures such as gingivectomy.

• Method of application: One teaspoonful of the emulsion is swished around in the mouth and oropharynx for 1-2 minutes; and later is spat out immediately prior to taking impressions.

#### Infiltration Anesthesia or Local Infiltration

 This method is also known as terminal or peripheral anesthesia, as the induction of anesthesia is by the action of anesthetic agents on the terminal nerve fibers.

#### **Maxilla and Mandible**

#### Maxilla

 The maxilla has thin labial/buccal cortical plate; and moreover shows areas of porosity, and the compact bone presents numerous foramina which aid in absorption of local anesthetic solution. These factors, therefore, make the maxilla more favorable for infiltration anesthesia techniques.

#### Mandible

The bone is generally dense and has thicker cortical plates than maxilla, particularly in posterior region, more so in the region of external oblique ridge. Only the anterior part of mandible presents sufficient porosity, which is favorable for infiltration techniques.

DI	Dense cortical bone
D2	Dense to porous cortical bone surrounding dense trabecular bone
D3	Thin porous cortical bone surrounding fine trabecular bone

D4 Fine trabecular bone

Anterior mandible Posterior mandible Anterior mandible Posterior mandible Anterior maxilla Anterior maxilla Posterior maxilla Posterior maxilla





 Small peripheral terminal branches or free nerve endings in a certain area of oral surgical procedure are anesthetised by deposition of local anesthetic solution in the area. Solution is injected beneath the mucous membrane, or along the periosteum, or beneath the skin.

# Complications of local anesthesia by Dr. Elham Hazeim Abdulkareem

### Introduction:

Administration of local anesthesia is daily routine for most dental practitioners normally no adverse effects are seen. However complications even very serious one can occur in daily practice. Complications related to local anesthesia can *be divided into two categories:* 

- 1. Preoperative (during giving anesthesia) complications.
- 2. Postoperative complications.

Both can usually be avoided by the correct technique and dosage however if complications occur the dentist should know how to manage them. *Each dentist dealing with local anesthesia should know:* 

- 1. The possible complications of local anesthesia.
- 2. The preventive measures of these complications.
- 3. The treatment possibilities of these complications.

## I-Peroperative complication

## 1. Pain at injection

Pain during administration of LA solution can be caused by many reasons.

- A- Factor depending on the solution:
- 1-Low PH value; this could irritate the tissue.

2-Temperature of the solution: warmer solution feeling more comfortable than cold. The cartridge can be warmed in the dentist hand or in warm water before the injection.

B- The practitioner related factors: This is related to the technique used can be avoided by:

1 - Fast injection cause rapid swelling of the tissue & pain.

2- Aggressive insertion of the needle can tear soft tissue, blood vessel, nerve & periosteum & cause more pain & others

# **Complications**

An inaccurate injection site can lead to an intramuscular or intraneural inj. when the needle penetrates a nerve; the patient feels a sudden (Electrical shock) in the distal area of the nerve. Pain after intramuscular injection is due to fibrosis or inflammation inside the muscle.

#### 2. Lack of effect.

Sometimes, the dentist face a problem, where the patient despite a conventional anesthesia still feels pain during treatment. The problem is most common with block anesthesia especially in lower jaw. Reasons behind his failure in LA can be classified as:

- 1- Anatomical.
- 2- Pathological.
- 3- Psychological.
- 4- Poor injection technique.

### 1- Anatomical reasons include :

- accessory nerve supply
- Abnormal course of the nerves.
- Thick cortical plate of the mandibular alveolus precludes infiltration of anesthesia & therefore infiltration anesthesia is insufficient in the lower jaw.
- Variation in the foramen location.
- Some times the teeth can be innervated by more than one nerve.

# 2- Pathological reasons are:

A- Trismus (limited mouth opening) in these cases it is impossible to use conventional technique of inferior alveolar nerve block. Therefore the so called Akinosi (closed mouth) technique is useful. In these cases there will be an increased possibility of failure of achieving adequate anesthesia.

**B-** Infection & inflammation: if the pulp is inflamed, the low tissue PH may cause lack of effective anesthesia in that area. However this does not explain failure in block anesthesia, where the solution is injection 4-5 cm from the inflamed area. A possible cause is hyperalgesia; the inflammation makes the nerve more sensitive. A minimal stimulation can cause pain perception. In those patient to obtain sufficient anesthesia more solution has to be injected, for example by combining a block & infiltration anesthesia. Supplemental intra-ligamentary or intra osseous injection can be used if necessary.

C- Previous surgery or trauma.

#### **3-** Psychological reasons:

Fear & anxiety. They can cause failure in LA, to enable successful anesthesia, relaxation of the patient is some time needed. For this the use of a sedative like benzodiazepine may be helpful.

*4- Poor technique:* Is the most common cause for insufficient anesthesia in inferior alveolar nerve block anesthesia .common mistakes are:

A- Injection of anesthesia too soon on the anterior ascending ramus as the needle point touches the lingual cortical bone anterior to lingula .

B- Another mistake is to injection inferior to mandibular foramen.

C- The solution can directed away from a nerve if it injection too rapidly & forcefully. So when ever possible it is advisable to follow accurate technique to prevent this complication.

# 3- Fainting & vasovagal attack

Collapse of the patient in the dental clinic may or may not be loss of consciousness.

Vasovagal attack is a reflex of nervous system that cause the heart to slow down (bradycardia), at the same time affect the nerves to blood vessel in the legs permitting these vessel to dilate (widen). As a result the heart ejects less blood so the blood pressure drops. Consequently the circulating blood goes to the legs rather than to the head & the fainting episode occurs. In an attempt to redistribute the blood to the vital organs.

Note: pain & anxiety are the most important predisposing factor to vasovagal attacks.

#### Signs and symptoms:

The patients often complain of the following symptoms:

**1-Dizziness** 

2-weakness

3-nausea

& following signs:

I-The skin is pale immediately before the collapse.

2- Cold skin

3- clammy skin 4- weak & slow pulse.

#### First aid should be taken immediately:

- The head should be lowered quickly by adjusting the back of the dental chair so that the patient assume the supine position with legs elevated.
- Tight collar & belts should be loosened and respiration is stimulated. Spontaneous recovery is usual, but if sign of recovery are not apparent within 30-45 sec of the first aide measure, the collapse probably is not a vasovagal attack & in this case medical emergency team should be asked immediately for help.
- Meanwhile, we need to maintain a patent air way & supplementary & consider the need for CPR (cardio-pulmonary resuscitation).

#### 4. Hypersensitivity or allergy to LA :

This phenomenon occurs due to sensitivity of some individual to certain substance known as (allergens).

Any LA agent may evoke such a response, but it is more commonly seen with ester type agent (e.g. Procaine) than amide type (e.g. lidocaine).true allergy to amide type is extremely rare.

Hypersensitivity reaction could be due to:

- 1- LA agent.
- 2- Vasoconstrictor.
- 3- Additives: e.g. bisulphite which used as preservative.
- In general hypersensitivity reaction to LA is very rare & represent less than 1% of all complications of LA. Many of complications suspected to be allergic are actually psychogenic reactions caused by fear of dental treatment.
- Other reasons may also be the presence of adrenaline in LA solution which can cause several general symptoms including palpitation, restlessness & nausea. True allergic response to LA may be localized or generalized & immediate or delayed in onset .they may vary from mild skin irritation or rashes to an anaphylactic reaction. Local reactions are seen more frequently than systemic & usually resolve without active treatment. If any degree of allergic reaction is observed, it is very important to determine the actual cause (allergen). Inadequate diagnosis & treatment can be life threatening to the patient.

**Anaphylactic shock:** It rarely occurs but it is life threatening it is characterized by:

- 1- Profound fall in blood pressure
- 2- Dyspnea & respiratory embarrassment
- 3- Facial & laryngeal edema
- 4- Loss of consciousness.

# 5- Urticaria.

It is life threatening condition since it causes air way obstruction in association with laryngeal edema unless treatment is initiated immediately; the condition may progress to fatal termination.

#### Management of anaphylaxis:

- If you suspect the anaphylactic reaction occurring, immediately seek medical help. The treatment must begin before BP & breathing problem become life threatening.
- epinephrine is the most imp medication for the treatment of anaphylaxis; it is inj into a muscle. Epinephrine works rapidly to make blood vessel contract, preventing them from leaking more fluid.
- It also relax air way, helping the individual breathe easier, relieves cramping in the GIT & stop itching, rapidly reverse the life threatening symptom of anaphylactic shock, if given in time. Even if the individual responds to the epinephrine, it is vitally imp to go to an emergency room immediately & medication may be given to improve breathing.
- IV fluid may be necessary to restore adequate BP. Additional medication (anti histamine) to contract the effect of histamine & to help in prevention of delayed allergic reaction. -If the victim stops breathing, perform cardio pulmonary resuscitation (CPR) immediately. Continue CPR until the person begins breathing again.
### **5- Over dosage & toxicity**

It is relatively rare ,a toxicity reaction can occur when the conc. of LA in circulation increases too rapidly within a short period of time e.g. inj too rapidly into highly vascular area or IV injection. The toxic effect is primarily directed to the CNS & CVS because these tissues are composed of excitable membrane.

### Mild over dose sign & symptom:

Restlessness, retention of consciousness talkativeness, & agitation, which may end into convulsion along with increased BP, heart rate, respiratory rate & classified according to the onset.

1- Slow onset <= 5 min following administration of LA (the possible cause rapid absorption & too large total dose).

2- Slower onset  $\geq 15$  min following administration of LA (the possible causes are abnormal biotransformation & renal disease).

### We use the following protocol in management of such cases:

Step 1 reassure the patient that every thing will be all right.

*Step 2 administer* via nasal cannula or nasal hood.

*Step 3* monitor vital signs.

Step 4 administer anti convulsant. IV infusion (for diazepam 5mg/min, for midazolam 1mg/min).

*Step 5* permits the patient to recover as long as necessary; the patient can leave the dental office escorted or unescorted if we believe that full recovery has occurred, if the doubt exist regarding the level of recovery then . *Step 6* seeks medical help.

*Step* **7** after termination of reaction be sure that the patient be examined by physician to determine the cause of this reaction.

## Severe overdose reaction:

Unconsciousness with or without convulsion, vascular collapse, coma, cardiac arrest & respiratory arrest & classified into:

1-Rapid onset (within 1 min) the possible cause is IV inj.

2-slow onset (5 to 15 min) (the possible causes are too large dose, rapid absorption & abnormal biotransformation.

## We use the following protocol in management of such cases:

Step 1 terminate dental treatment.

*Step 2* if convulsion is present, protect the patient arm, legs & head & loosen the tight clothing and immediately seek emergency medical assistance.

*Step 3* provide basic life support (maintenance of adequate air way & oxygenation are of utmost importance during management of LA induced seizures.

*Note:* seizures cause increase O2 utilization & hyper metabolism, with increased production of C02 & lactic acid & lead to acidosis this in turn lower the seizer threshold & prolongation of reaction. Cerebral blood flow during the seizure also increase this lead to elevation of anesthetic blood level within the CNS.

*Step 4* administer anticonvulsant with good oxygenation & acidosis is not present within 2-3 min seizure will cease. If the seizure is protracted (4-5min with no indication of termination then consider administration of anticonvulsant.

*Step 5* post seizure CNS depression is usually present & It s intensity equal to that of excitation phase (the patient is drowsy ,or unconscious, breathing may be shallow or absent BP ,HR are depressed so maintenance of patent air way & oxygenation are necessary ,beside administration of IV fluid is initial management of hypotension.

*Step 6* allow the patient to rest until recovery

**Note:** toxicity can also be produced by vasoconstrictor, such as adrenaline in this case the possible symptoms are increase in fear & anxiety, tremor, head ache & palpitation.

## To avoid toxicity:

1- Keep in mind the max safe doses of anesthetic. In children & elderly patient, safe level of LA are lower than the remaining population.

2- Use aspirating technique before & during inj. In this manner an IV inj can be avoided.

3- Injection the LA solution slowly.

4- Monitor the patient for possible S.E during injection.

### **6-Drug interaction:**

In some patient the administration of 2 drugs will counteract each other, while in other potentiation occurs. In patient under tricyclic antidepressant (TCA) ,variable degrees of potentiation of BP response to catecholamine (adrenaline & noradrenalin )will occur ,even to small doses therefore percussion should be taken during the use of these types of vasoconstrictor with patient taking the TCA. If LA contains felypressin vasoconstrictor, no effect will occur.

### 7-needle breakage

It is a rare complication in modern dentistry because of the larger awareness of he possible causes & the method for avoiding this complication. The reasons for needle breakage may be:

1- Weakness of the metal of the needle: Metal alloy used nowadays in the needles are flexible, more durable than previously, this lower the number of needle Breakage, but has not entirely solved the problem.

2- re-usage of the needle:

It is not accepted in dentistry nowadays; yet re-usage may happen during the same appointment when giving additional dosage of LA to the same patient. Repeated injection with the same needle cause fatigue of the structure & the risk of needle fracture increases.

3-incorrect technique: this includes

- Aggressive insertion of the needle in to the tissue.
- Sudden change in the direction inside the tissue.
- Too deep penetration, the needle goes up to it s hub inside the tissue & might fracture at this point. The hub is considered the most common point of needle fracture.
- 4- Sudden movement of the patient or practitioner.
- 5. Manufacturing defect

### Prevention of needle breakage:

- The dentist should check the inj needles before using them. If there is any suspicion of inadequate product quality a new one should be used.
- Avoid repeated usage of the needle even for the same patient.
- Adequate learning of inj technique for different areas in the mouth.
- Avoid aggressive movement & change of the direction during inj.
- Aggressive insertion can lead to sudden movement of the patient making the dentist unable to control the movement of the needle.

### What to do when the needle breaks:

- 1 -Stay calm & try to localize broken part in the tissue.
- 2- Tell the patient what has happened & try to relax & comfort him/her.
- 3- Stabilize the patient jaws in order that the needle stays in place (if the patient move his jaws the tension from the muscle of masticatory system cause the needle to penetrate the tissue &pain).
- 4- If a portion of the needle is visible, grasp it firmly with a pair of locking forceps & remove it.
- 5- If you can not remove the broken part yourself, refer the patient to oral & maxillofacial surgeon. In the department of oral & maxillofacial surgery the patient is examined well & the needle is removed surgically UGA ,this will provide complete immobility & muscle relaxation with the aid of some method that are used to localized the needle (ex CT scan , ultrasonography's).

## 8. Facial nerve paralysis:

This complication arises if the tip of the needle is inserted too far back & behind the ascending ramus of the mandible during inferior alveolar nerve block. The anesthetic agent then deposited in the substance of the parotid gland where it anesthetize the branches of the facial nerve causing paralysis of facial muscles.

Facial nerve paralysis secondary to LA injection is temporary & will last the expected duration of anesthesia of soft tissue of the oral cavity.

Clinically, the patient will immediately complain of transient paralysis of the muscle of the chin, lower lip, upper lip, of eyelid (in ability to close the eye) & in ability to raise the eye brow of the affected side.

## Management:

1-reassure the patient of the transient nature of the event & will last for few hours.

2-advice the patient to use an eye patch until motor function return.

**3-if contact lenses are worn, they should be removed.** 

# 2-post operative complication

## I- hematoma

A hematoma is localized mass of extravasated blood that may become clinically noticeable following the injection. It is caused by penetration of the BV with the needle during penetration or withdrawal of the needle in to the tissue leading to bleeding inside the tissue & the formation of a hematoma. The patient will notice development of swelling & discoloration of a bruise lasting IO days.

Intra orally the vessels most commonly associated with hematoma are:

- 1- The pterygoid plexus of veins.
- 2- Posterior superior alveolar vessels.
- 3- The inferior alveolar vessels in the pterygomandibular space.
- 4- The mental vessels.
- 5- The infraorbital vessels.

### Prevention:

- Learn anatomical landmark & injection technique.
- Avoid relocating the needle to different sites inside the tissue.

## Management:

1- If it is visible immediately following injection, apply direct pressure if possible.

2- Once bleeding has stopped inform the patient of what happened & re-evaluate the possibilities of continuing the treatment.

3- Instruct the patient to avoid application of heat over the area for at least 6 hours.

4- Tell the patient to expect discoloration of the area.

5- Prescribe analgesics & if infection is suspected, suitable antibiotic is given.

6- If it is invisible like in case of pterygomandibular space hematoma, the patient will normally come in the second or third day complaining of trismus. In this case treat the patient as the treatment of trismus.

## 2- Trismus:

Trismus is defined as limited jaw opening (locked jaw) mainly due to spasm of the muscle of mastication.

### The causes of spasm:

1- Penetration of the muscle by needle during administration of LA which might cause bleeding in to the muscle & spasm.

2- Hematoma as a result of damage of blood vessel in the area near one of muscle of mastication.

3- Infection in an area close to one of the muscle of mastication .The most common muscle to be the source of trismus is the medial pterygoid muscle which can be penetrated by the needle during inferior alveolar nerve block.

The main symptom is the limitation of movement of mandible which is often associated with pain, symptoms arise within one to four days of injection, their duration & severity are variable .if properly managed improvement should be noticed within 2-3 days if no improvement is noted, consider other possible cause such as infection & treat accordingly.

## **Management of trismus:**

1- Apply hot moist towel to the site for approximately 20 minute every hour.

2- Use analgesia as required.

3- Instruct the patient to gradually open &close the mouth & use chewing gum as mean of physiotherapy.

# **3- Infection:**

Infection following LA administration is extremely rare nowadays due to the use of disposable needles.

It may occur if the **needle has been contaminated** prior to insertion or if the patient significantly immunocompromised. If an infection does occur, it will likely manifest initially as pain & trismus one day post inj. If these symptoms persist for days & continue to worsen, the possibility of infection should be considered. At this stage, this patient should be examined for other sign & symptom of infection such as swelling, lymphadenopathy & fever & drainage of pus collection should be considered.

When there is an active site of infection such as an abscess, needles should not be inserted because there is a potential for spreading the infection.

## 4- Prolonged anesthesia & paraesthesia:

Anesthesia, paresthesia, dysesthesia, hyperesthesia; may persist beyond the expected duration of a local anesthesia affecting the tongue or lip.

## Anesthesia: is a total loss of sensation

Paraesthesia: is an altered sensation, characterized by burning, tingling, pricking sensation experienced by the patient.

Dysesthesia: is an Abnormal, unpleasant sensation experienced by the patient in the absence of stimuli.

Hyperesthesia: is an increased sensitivity to noxious stimuli.

### Causes:

1- Direct trauma from the bevel of the needle.

2- injection of the anesthetic solution that is contaminated with a neurotoxic substance such as alcohol.

**3-** Hemorrhage & infection in close proximity to a nerve: Most of these are transient & resolve within 8 weeks -2 years, but they become irreversible in rare occasions.

## Management:

- 1- Reassurance of the patient that the condition is transient.
- 2-Note the sign & symptom & follow up of the patient
- 3- Refer the patient to maxillofacial surgeon if the symptoms persist.

# **Abbreviations:**

Lymphadenopathy: any disease process that cause enlargement of lymph node.

- The vital signs (BP, PR, RR, temperature)
- BP = blood pressure PR = pulse rate
- RR= respiratory rate Temp= temperature
- S.E= side effect UGA= under general anesthesia
- // = fracture Inj. = injection
- Imp.= important LA = local anesthesia
- Dis = disease

# **Classification of nerve fibers:**

The fibers of peripheral nerves classified according to the basis of electrophysiological and morphological differences:

**Type A ; fiber:** largest fibers further divided into four groups :-1) A- alpha ( $\alpha$ ): responsible for motor action & muscle properioception. 2) A-beta ( $\beta$ ): responsible for motor action & muscle properioception. 3) A-gamma ( $\gamma$ ): responsible for motor action & muscle properioception. 4) A- delta ( $\delta$ ): responsible for fast sharp pain, temperatures, touch, & pressure

**Type B fibers :** preganglionic responsible for sympathetic activity. **Type C fibers :** unmyelinated , the most numerous in the peripheral nervous system. Responsible for conduction of dull or burning pain. Noxious stimuli are transmitted to the CNS by way of A $\delta$  &C fibers. The lightly myelinated A $\delta$  fibers are responsible for conduction of

sharp, bright pain while unmyelinated C fibers conduct dull or burning pain.

## **Requirements of local anesthetic agents:-**

- It should not be irritating to the tissue to which it is applied
- It should not, cause any permanent alteration of nerve structure
- Its systemic toxicity should by low
- It must be effective regardless of whether it is injected into the tissue or applied locally to mucous membranes.
- The time of onset of anesthesia should be as short as possible
- The duration of action, must be long enough to permit completion of the procedure.
- It should have potency sufficient to give anesthesia without using harmful concentration
- It should be relatively free from producing allergic reactions
- It should be stable in solution and readily undergo biotransformation in the body
- It should either be sterile or be capable of being sterilized by the heat without deterioration.

# Nerve membrane:-

The nerve membrane consists of two layers of lipid molecules (phospholipids) and associated proteins, lipids and carbohydrates (fig: 2). All biologic membranes are organized to

- (1) block the diffusion of water soluble molecules.
- (2) be selectively permeable to certain molecules via specialized channels and
- (3) transduce information by protein receptors responsive to chemical or physical stimulation. Since the nerve membrane exhibits selective permeability, therefore significant differences exist for ions between the intracellular and the extracellular concentrations. Accordingly high concentration of K inside while high concentration of Na and CI outside the nerve membrane.

In some nerves an insulating lipid rich layer of myelin covers this membrane (fig: 3). The outer most layer of myelin consists of the schwann cell cytoplasm and its nucleus. A gap between two adjoining Schwann cells and their myelin spirals called node of Ranvier at these nodes the nerve membrane is exposed directly to the extracellular medium.

1. Ester linked local anesthetic (ex: procaine) are readily hydrolyzed in aqueous solution, while amide linked types (ex: lidocaine) are relatively resistant to hydrolysis.

2. a greater percentage of an amide linked drug is excreted unchanged in the urine than of an ester linked drug.

# The ester type includes the following:-

- 1. procaine
- 2. chloroprocaine
- 3. propoxycaine
- 4. butacaine
- 5. cocaine
- 6. benzocaine
- 7. hexylcaine
- 8. piperocaine
- 9. tetracaine

The amide type include the following:-

- 1. lidocaine
- 2. prilocaine
- 3. articaine
- 4. bupivacaine
- 5. dibucaine
- 6. etidocaine
- 7. mepivacaine

## Pharmacokinetic of local anesthetics:-

**Distribution:-**

Once absorbed into the blood local anesthetics are distributed throughout the body to all tissues. The blood level of local anesthetic is influenced by the following factors:-

1. rate at which the drug is absorbed into the cardiovascular system.

2. rate of distribution of the drug from the vacular compartment to the tissue (more rapid in healthy patients than in those who are medically compromised e.g; congestive heart failure patients).

3. elimination of the drug through the metabolic or excretory pathways.

The last two factors act to decrease the blood level of local anesthetic. **Note:-**

All local anesthetics readily cross the blood brain barrier they also readily cross the placenta and enter the circulatory system of the developing fetus.

# Dilution of vasoconstrictors'

The di l u t ion of vasoconstrictor is commonly referred to as ratio (1 to 1000, written 1: 1000). This 1:1000 mean that there is I gm (1000 mg) of anesthetic drug contained in 1000 ml of solution.

Therefore a 1:1000 dilution contains 1000 mg in 1000 ml or 1.0 mg/ml of solution. Accordingly 1:10000 dilutions contain 1000 mg in 10000ml or 0.1 mg/ml of solution. & 1:100 000 dilution

contain 1000mg in 100 000 ml or 0.01 mg/ml of solution.

Туре	mg/ml	dilution
Adrenalin	0.02	1:50 000
Adrenalin	0.01	25 1:80 000
Adrenalin	0.01	1:100 000
Adrenalin	0.005	1:200 000
Levonorde frin	0.05	1:20 000
Noradrenalin	0.033	1:30 000

## The mg/ml value of the various vasoconstrictors are :

We should remember that the max dose of the vasoconstrictor is 0.2 mg per appointment for adult healthy patient. And 0.04 mg per appointment for patient with clinically significant cardiovascular dis, & should be considered when calculating the max number of cartridge that are allowed to be given to the patient per appointment & according to the type & cone. of the vasoconstrictor.

Туре	Pulpal	Soft tissue anesthesia
Lidocaine 2%	5-10 min.	60 min
Prilocaine 4%	5-10 min	90 min
Mepivacaine 3%	20-40 min	120 min
Lidocaine 2%	60 min	180-240 min
Prilocaine 4%	60 min	120 min
Bupivacaine 0.5%	> 90 min	240-540 min



Fig (2): dental syringe (top loaded cartridge)



Fig(**3**):pressure syringe (origenal design)









## Bent needle



## Fig (7):centric and eccentric perforation













#### **Third Year**

#### **Oral Surgery**

#### **Dental Management of the Medically Compromised Patients**

#### Medical history

A medical history must be taken for every patient who is to receive dental treatment. There are two basic techniques were used to obtain a medical history. The first technique consists of an interview of the patient, in which the interviewer questions the patient and then records a narrative of the patient's verbal responses on a form sheet.

The second technique was used a printed questionnaire that the patient fills out.

#### 1. Cardiovascular Disease

Patients with cardiovascular disease are especially vulnerable to physical or emotional challenges that may be encountered during dental treatment.

**2. Heart failure** results from an underlying cardiovascular problem such as coronary heart disease or hypertension. Patients with untreated or symptomatic heart failure are at increased risk for myocardial infarction (MI), arrhythmias, acute heart failure, or sudden death, and generally are not candidates for elective dental treatment. <u>Chair position may influence ability to breathe, with some patients unable to tolerate a supine position.</u> <u>Vasoconstrictors should be avoided, if possible, in patients taking digitalis glycosides (digoxin) because the combination can precipitate arrhythmias.</u>

#### 3. Heart Attack

A history of a heart attack (myocardial infarction) within the very recent past may prevent elective dental care, because during the immediate post infarction period, patients are at increased risk for reinfarctions, arrhythmias, and heart failure. Patients may be taking medications such as antianginals, anticoagulants, adrenergic blocking agents, calcium channel blockers, antiarrhythmic agents, or digitalis. Some of these drugs may alter the dental management of patients because of potential interactions with vasoconstrictors in the local anesthetic, adverse side effects, or other Considerations. Stress and anxiety reduction measures may be advisable.

#### 4. Angina Pectoris

is a brief substernal pain results from myocardial ischemia, commonly provoked by physical activity or emotional stress, is a common and significant symptom of coronary heart disease. Patients with angina, especially unstable or severe angina, are at increased risk for

#### Third Year

#### **Oral Surgery**

arrhythmias, MI, and sudden death. A variety of vasoactive medications, such as nitroglycerin,  $\beta$ -adrenergic blocking agents, and calcium channel blockers, are used to treat angina. Caution is advised with the use of vasoconstrictors. Stress and anxiety reduction measures may be appropriate. Patients with unstable or progressive angina are not candidates for elective dental care.

### 5. High Blood Pressure

Patients with hypertension (blood pressure greater than 140/90 mm Hg) should be identified by history and the diagnosis confirmed by blood pressure measurement. Patients with a history of hypertension should be asked if they are taking or are supposed to be taking antihypertensive medication. Failure to take medication often is the cause of elevated blood pressure in a patient who reports being under treatment for hypertension. Current blood pressure readings and any clinical signs and symptoms that may be associated with severe, uncontrolled hypertension, such as visual changes, dizziness, spontaneous nosebleeds, and headaches, should be noted.

Some antihypertensive medications, such as the nonselective ßadreneraic blocking agents, require caution may in the use of vasoconstrictors. The coadministration of calcium channel blockers with macrolide antibiotics (e.g., erythromycin, clarithromycin) can result in excessive hypotension. Stress and anxiety reduction measures also may be appropriate. Elective dental care should be postponed for patients with severe, uncontrolled hypertension (blood pressure of 180/110 mm Hq or higher) until the condition can be brought under control.

### 6. Artificial Heart Valve

A diseased valve may be replaced with artificial or prosthetic valves. Such replacement valves are associated with a high risk for development of infective endocarditis, with significant morbidity and mortality. Accordingly, the AHA recommends that all patients with a prosthetic heart valve be given prophylactic antibiotics before most dental procedures. Patients with an artificial heart valve also may be on anticoagulant medication to prevent blood clots associated with the valve. In such patients, excessive bleeding may be encountered with surgical procedures; it is therefore necessary to determine the level of anticoagulation before any invasive procedure.

### 7. Hemophilia or inherited bleeding disorder

Patients with an inherited bleeding disorder such as hemophilia A or B, or von Willebrand's disease, are at risk for severe bleeding after any type of dental treatment that causes bleeding, including scaling and root planing. These patients must be identified and managed in cooperation with their physician or haematologist. Patients with severe factor deficiency may require factor replacement before invasive treatment, as well as aggressive postoperative measures to maintain hemostasis.
### **Oral Surgery**

### 8. Blood Transfusion

Patients with a history of blood transfusions are of concern from two aspects. The underlying problem that necessitated a blood transfusion, such as an inherited or acquired bleeding disorder, must be identified, and alterations in the delivery of dental treatment may have to be made. These patients also may be carriers of hepatitis B or C or may have become infected with the human immunodeficiency virus (HIV) and must be identified. Laboratory screening or medical consultation may be appropriate to determine the status of liver function, and, as always, standard infection control procedures are mandatory.

### 9. Anemia

A significant reduction in the oxygen-carrying capacity of the red blood cells may result from an underlying pathologic process such as acute or chronic blood loss, decreased production of red blood cells, or haemolysis. Some anaemias, such as glucose-6-phosphate dehvdrogenase (G6PD) deficiencv and sickle cell disease, reauire dental management Oral lesions, infections, modifications. delayed wound healing, and adverse responses to hypoxia all are potential matters of concern

### **10.** Leukemia/Lymphoma

Depending on the type of leukemia or lymphoma, status of the disease, and type of treatment, some patients may have bleeding problems or delayed healing, or may be prone to infection. Gingival enlargement can be a sign of leukemia. Some adverse effects can result from the use of chemotherapeutic agents and may require dental management modifications.

### **11.** Epilepsy, Seizures, and Convulsions

A history of epilepsy should be identified, and the degree of seizure control should be determined. Specific triggers of seizures (e.g., odors, bright lights) should be identified and avoided. Some medications used to control seizures may affect dental treatment because of drug actions or adverse side effects. For example, gingival over growth is a wellrecognized adverse effect of diphenylhydantoin (Dilantin). Patients may discontinue the use of anticonvulsant medication without their doctor's knowledge and thus may be susceptible to seizures during dental treatment. Therefore, verification of patients' adherence to their medication schedule is important.

### **12. Psychiatric Patients**

Patients with a history of a behavioral disorder or psychiatric illness as well as the nature of the problem need to be identified. This information may help explain patients' unusual such as unexplainable or unusual conditions. Additionally, some psychiatric drugs have the potential to interact adversely with vasoconstrictors in local anesthetics. They also may produce adverse oral effects such as hyposalivation or xerostomia.

### **Oral Surgery**

Some patients may be excessively anxious or apprehensive about dental treatment, requiring stress reduction measures.

### 13. Stomach or Intestinal Ulcers, Gastritis, and Colitis

Patients with gastric or intestinal disease should not be given drugs that are directly irritating to the gastrointestinal tract, such as aspirin or nonsteroidal antiinflammatory drugs. Patients with colitis or a history of colitis may not be able to take certain antibiotics. Many antibiotics can cause a particularly severe form of colitis (i.e., pseudomembranous colitis), and elderly persons are more susceptible to this condition. Some drugs used to treat gastric or duodenal ulcers may cause dry mouth.

### 14. Hepatitis, Liver Disease, Jaundice, and Cirrhosis

Patients who have a history of viral hepatitis are of concern in dentistry because they may be asymptomatic carriers of the disease and can transmit it unknowingly to dental personnel or other patients. Of the several types of viral hepatitis, only hepatitis B, C, and D have carrier stages. The laboratory tests are available to identify affected patients. Standard infection control measures are mandatory. Patients also may have chronic hepatitis (B or C) or cirrhosis, with associated impairment of liver functio may result bleeding and less efficient metabolism of certain drugs, including local anesthetics and analgesics.

### **15.** Allergies or Hives

Patients may be allergic to some drugs or materials used in dentistry. Common drug allergens include antibiotics and analgesics. Latex allergy also is common, and in patients so affected, alternative materials such as vinyl or powderless gloves and vinyl dam material can be used to prevent an adverse reaction. True allergy to amide local anesthetics is uncommon. Dentists should procure a history regarding allergy by specifically asking patients how they react to a particular substance. Symptoms and signs with allergy include itching, urticaria (hives), rash, swelling, wheezing, angioedema, runny nose, and tearing eyes. Isolated signs and symptoms such as nausea, vomiting, heart palpitations, and fainting generally are not of an allergic origin but rather are manifestations of drug intolerance, adverse side effects, or psychogenic reactions.

### 16. Asthma

The type of asthma should be identified, as should the drugs taken and any precipitating factors or triggers. Stress may be a precipitating factor and should be minimized when possible. It often is helpful to ask whether the patient has visited the emergency room for acute treatment of asthma, because this historical detail would indicate more severe disease. A patient who uses an albuterol inhaler for treatment of acute attacks should be instructed to bring it to the dental appointment.

### **Oral Surgery**

### 17. Diabetes

Patients with diabetes mellitus must be identified to determine the type of diabetes, how it is being treated, and how well controlled it is. Patients with type 1 diabetes require insulin, whereas type 2 diabetes usually is controlled through diet and/or oral hypoglycaemic agents; however, some patients with type 2 diabetes eventually also require insulin. Those with type 1 diabetes have a greater number of complications and are of greater concern regarding management than are those with type 2 diabetes. Symptoms and signs of diabetes include excessive thirst and hunger, frequent urination, weight loss, and frequent infections. Long-term complications include blindness, hypertension, and kidney failure, each of which also may affect dental management. Patients with diabetes typically do not handle infection very well and also may have exaggerated periodontal disease. Patients who take insulin are at risk for episodes of hypoglycaemia in the dental office if meals are skipped or if infection is present.

### **18. Thyroid Disease**

Patients with uncontrolled hyperthyroidism are potentially hypersensitive to stress and a1- adrenergic effects of sympathomimetic, so the use of vasoconstrictors generally is contraindicated. In rare cases, infection or surgery can initiate a thyroid crisis—a serious medical emergency. These patients also may be easily upset emotionally and intolerant of heat, and they may exhibit tremors. Exophthalmos may be present. Patients with known hypothyroidism usually are taking a thyroid supplement; this medication regimen generally warrants no concern so long as the thyroid hormone level does not become too high.

### **19. Kidney Failure**

Patients with chronic kidney disease or a kidney transplant must be The potential abnormal identified. for drug metabolism, immunosuppressive drug therapy, bleeding problems, hepatitis, infection, blood pressure, and heart failure must be considered hiah in management. Certain drugs that are nephrotoxic should be avoided. Patients on hemodialysis do not require antibiotic prophylaxis.

### **20. Sexually Transmitted Diseases**

A variety of sexually transmitted diseases such as syphilis, gonorrhea, human immunodeficiency virus (HIV) infection, as well as AIDS, can have manifestations in the oral cavity because of oral genital contact or secondary to hematogenous dissemination in the blood or immune suppression. The dentist may be the first to identify these conditions. In addition, some sexually transmitted diseases, including HIV infection, hepatitis B and C, and syphilis, can be transmitted to the dentist through direct contact with oral lesions or infectious blood.

### **Oral Surgery**

### 21. Tobacco and Alcohol Use

The use of tobacco products is a risk factor that is associated with cancer, cardiovascular disease, pulmonary disease, and periodontal disease. Excessive use of alcohol is a risk factor for malignancy and heart disease, and may lead to liver disease. The combination of excessive alcohol and tobacco use is a significant risk factor for oral cancer.

### 22. Drug Addiction and Substance Abuse

Patients who have a history of intravenous drug use are at increased risk for infectious diseases such as hepatitis B or C, AIDS, and infective endocarditis. Narcotic and sedative medications should be prescribed with caution, if at all, for these patients, because of the risk of triggering a relapse. This caveat also applies to patients who are recovering alcoholics. Vasoconstrictors should be avoided in patients who are cocaine or methamphetamine users because the combination may precipitate arrhythmias or severe hypertension.

### 23. Tumors and Cancer

Patients who have had cancer are at risk for recurrence, so they should be closely monitored. Also, cancer treatment regimens including chemotherapeutic agents or radiation therapy may result in infection, gingival bleeding, oral ulcerations, dry mouth, mucositis, and impaired healing after invasive dental treatment, all of which represent significant management considerations. Patients with a history of intravenous bisphosphonate therapy for metastatic bone disease are at risk for osteonecrosis of the jaw, and surgical treatment should be managed cautiously.

### 24. Radiation Therapy and Chemotherapy

Patients with previous radiation treatment to the head, neck, or jaw must be carefully evaluated, because radiation can permanently destroy the blood supply to the jaws, leading to osteoradionecrosis after extraction or trauma. Irradiation of the head and neck can destroy the salivary glands, resulting in decreased saliva, increased dental caries, and mucositis. Fibrosis of masticatory muscles resulting in limited mouth opening also may occur. Chemotherapy can produce many undesirable adverse effects, most commonly a severe mucositis; however, such changes resolve with cessation of the chemotherapeutic agents.

### 25. Steroids

Cortisone and prednisone are examples of corticosteroids that are used in the treatment of many diseases. These drugs are important because their use can result in adrenal insufficiency and potentially render the patient unable to mount an adequate response to the stress of an infection or invasive dental procedure such as extractions or periodontal surgery. Generally, however, most routine dental procedures do not require administration of supplemental steroids.

#### **Oral Surgery**

#### 26. Pregnancy

Women who are or may be pregnant may need special consideration in dental management. Caution typically is warranted in the taking of radiographs, administration of drugs, and timing of dental treatment. Prolonged time in the dental chair should be avoided, to prevent the complication of supine hypotension. If supine hypotension develops, rolling the patient onto <u>her left side affords return</u> of circulation to the heart. Good dental hygiene is important to maintain during pregnancy. Preventive plaque control measures should be provided in the first trimester. Extensive reconstruction or significant surgical procedures are best postponed until after delivery. In spite of the safety of dental radiography, ionizing radiation should be avoided. The most important for the pregnant patient are the protective lead apron and the thyroid collar. In addition, the use of digital radiography markedly reduces radiation exposure.

#### **References:**

1. Dental Management of the Medically Compromised Patient ISBN: 978-0-323-08028-6 Copyright © 2013 by Mosby, an imprint of Elsevier Inc. Library of Congress Cataloging-in-Publication Data 2. Dental management of the medically compromised patient / James W. Little ... [et al.]. – 8th ed.p. ; Cm.Includesbibliographical references and index.ISBN 978-0-323-08028-6 (pbk. : alk. paper)

I. Little, James W., 1934- [DNLM: 1. Dental Care.

3. Dental Care for Chronically Ill. 3. Oral Manifestations. WU 29] 617.6-dc23.

# Armamentarium of Local Anaesthesia By Dr. Elham H. Abdulkareem

# The instruments required for initial examination are shown in Figure.



# The armamentarium

The equipment's necessary for the administration of local anaesthetic include:

- 1. The syringe
- 2. The needle
- 3. The local anaesthetic cartridge
- 4. Additional armamentarium:
- a) Topical antiseptic
- b) Topical anaesthetic
- c) Applicator sticks
- d) Cotton gauze (2\*2 inches)
- e) Haemostate

# The syringe

The syringe is the vehicle whereby the contents of the anaesthetic cartridge are delivered through the needle to the patient.

There are 7 types of syringe for local anaesthetic administration in use in dentistry which are:

- A. Non disposable:
- 1. Breech-loading metallic cartridge-type aspirating.
- 2. Breech-loading metallic cartridge-type non-aspirating.
- 3. Breech-loading plastic cartridge-type aspirating.
- 4. Breech-loading metallic cartridge-type self-aspirating.
- 5. Pressure
- 6. Jet injector
- B. Disposable
- C. Safety syringes.

Nonaspirating, breech loading cartridge syringe



Various types of aspirating cartridge syringes showing: A diamond, a fishhook, and a harpoon; as seen from above downwards





# American dental association criteria for acceptance of local anaesthetic syringes include the following:

 $\checkmark$  They must be durable and able to withstand repeated sterilization without damage.

 $\checkmark$  They should be capable of accepting a wide variety of cartridges and needles.

 $\checkmark$  They should be permit repeated use.

 $\checkmark$  They should be inexpensive, self-contained, light weight, and simple to use with one hand.

 $\checkmark$  They should provide for effective aspiration and be constructed so that blood may be easily observed in the cartridge.

## Breech-loading metallic cartridge-type aspirating syringe is the most commonly used.

The term breech-loading implies that the cartridge is inserted into the syringe from the side. Barrel.

The syringe consists of:

- $\Rightarrow$  Metal barrel
- $\Rightarrow$  Plunger united (piston)



- A double-ended needle is attached by means of a screw hub to other ended of the barrel. Opening of the hinge mechanism permits the insertion of a cartridge.
- The aspirating syringe has a device such as a tip or harpoon that is attached to the piston an is used to penetrate the thick rubber or silicone stopper at the opposite end of the cartridge.
- The manoeuvre of aspiration consists of withdrawal of the rubber plunger to create a negative pressure within the cartridge and is employed in order to ensure that a blood vessel has not been entered by the needle tip during its insertion into the soft tissues and prior to the injection of the solution. Positive aspiration results in a fine spiral of blood being clearly visible in the solution within the cartridge.

## The advantages are:

- Visible cartridge
- Aspiration with one hand
- Autoclavable
- Rust resistant
- Long lasting with proper maintenance
  The disadvantages are:
- Heavy weight
- May be too big size for small operators
- Possibility of infection with improper care.



A slight amount of aspiration may be achieved as follow:

Slight pressure on the piston produces a bulge in the rubber plunger. Which in turn exerts pressure on the solution in the cartridge, release of this produces suction. If the needle tip is in a blood vessel, a thin spiral of blood which rapidly diffuses will be seen in the solution. Pressure syringes introduced in the late 1970, pressure syringes brought about a renewed interest in the periodontal ligament injection and also called intraligamentary injection technique.

The 1st generation devices, using a pistol grip, are somewhat larger than the newer pen grip devices.

The advantages are:

- Measured dose
- Overcomes tissue resistance
- Cartridge protected

The disadvantages are:

• Cost

• Easy to inject too rapidly

Tilling the

Used for performing Periodontal Ligament (PDL) injections

- Compact and extremely durable
- Pen-style activator
- Injects only 0.06 ml of local anesthetic solution with each click of the activator level

Jet injector the primary uses to obtain topical anaesthesia prior to the insertion of a needle. In addition, it may be used to obtain mucosal anaesthesia of the palate.

The advantages are:

- Dose not require use of a needle
- Delivers very small volumes of local anaesthetic solution
- Used in topical anaesthesia

The disadvantages are:

- Inadequate for pulpal or regional block
- Some patients are disturbed by jet injection
- Cost

May damage periodontal tissues

# Care and handling

The recommendations concerning care of the syringes as following:

1. After each use thoroughly wash and rinse the syringe free of any local anaesthetic solution, saliva, or other foreign matter. Autoclave the syringe in the same manner as other surgical instruments.

2. After every five autoclaving's, dismantle the syringe and lightly lubricate all threaded joints and where the piston contacts the thumb ring and guide bearing.

3. Clean the harpoon with a brush after each use.

# **Clinical problems**

- Leakage during injection: When reloading a syringe with a second local anaesthetic cartridge and a needle already in place. One should be sure that the needle penetrates the center of the rubber diaphragm. An off-centre perforation will produce an ovoid puncture of the diaphragm that permits leakage of the anaesthetic solution around the outside of the metal needle and into the patient's mouth.
- Broken cartridge: A badly worn syringe may damage the cartridge, leading to breakage. This can result from a bent needle at its proximal end may not perforate the diaphragm on the cartridge. Positive pressure on the thumb ring increases intra-cartridge pressure, which can cause the cartridge to break.

## The needle

The needle permits the local anaesthetic solution to travel from the cartridge into the tissues surrounding the needle tip. Most needles used in dentistry are stainless steel and are disposable.

### Parts all needles have several components which include:

- $\clubsuit$  The bevel: is long, medium, and short.
- $\clubsuit$  The shank or shaft: the needle gauge and length.

Solution The hub: is the plastic or metal piece through which the needle attaches the syringe.

Solution The syringe adaptor: the interior surface of the needle

Solution The syringe penetrating end: is placed into the needle adaptor and perforates the rubber diaphragm of the cartridge.

the needles used in dental surgery, range from 24, 25, 27 and 30-gauges; and from 25 mm to 38-40 mm in length



Needles in dental practice are made up of stainless steel,

is most commonly used and is highly recommended.



Needles currently available are usually pre-sterilized and disposable

## When needles are selected for use in various injection technique, there are two factors must be considered which are:

- The gauge: Gauge refers to the diameter of the lumen of the needle; the smaller the number the greater the diameter of the lumen. A 30 gauge needle has a smaller internal diameter than a 25 gauge needle. The smaller diameter needles are less traumatic to the patient than needles with larges diameter. The most commonly used needles in dentistry are 25, 27, and 30 gauge
- Length:: Dental needles are available in two lengths: long approximately 40mm and short approximately 25mm. Needles should not be inserted into tissues to their hub unless it is absolutely necessary for the success of the injection. There are several reasons for this rule: needle breakage. **The weakest portion of the needle is at the hub so that the needle breakage occurs.** Long needles are preferred for all injection techniques requiring penetration of significant thicknesses of soft tissue.

# **Care and handling**

- 1. Needles must never be used on more than one patient.
- 2. Needles should be changed after three or four tissue penetration in same patient.
- 3. Needles should be covered with a protective sheath when not being used.
- 4. Needles must be properly disposed of after use.



### **Clinical problems**

- Pain on insertion: The used of a dull needle can lead to pain on initial penetration of the mucosa. This may be avoided by using sharp, new, disposable needles and the application of topical anaesthetic at the penetration site.
- Breakage: Bending weakens needles making them more likely to break on subsequent contact with hard tissues such as bone. Never attempt to force a needle against resistance. Smaller gauge needles are more likely to break than larger gauge needles.
- Pain on withdrawal: Pain on withdrawal is more likely that they occur when the needle tip forcefully contacts a hard surface as bone, therefore a needle should never be forced against resistance.
- Injury to the patient or administrator: A major cause is carelessness and inattention by the administrator, although sudden unexpected movement by the patient is also a frequent cause. The needle should remain capped until is to be used and should be made safe immediately after withdrawal from the mouth.

**The cartridge (carpule)** The dental cartridge is a glass cylinder containing the local anaesthetic drug. The glass cylinder itself can hold either 1.8ml, 2ml, or 2.2ml. The dental cartridge is by common usage, referred to as **a carpule** by dentist.

The prefilled dental cartridge consists of four parts are:

• Cylindrical glass tube

• **Stopper:** is located at the end of the cartridge that receives the harpoon of the aspirating syringe. The stopper was mixed with paraffin wax to produce an airtight seal against the glass walls of the cartridge. Glycerine was added in channels around the stopper as lubricant. Today the stopper treated by silicone, eliminating both the paraffin and the glycerine.

▲ Aluminum cap: is located at the opposite end of the cartridge from the rubber plunger. It fits around the neck of the glass cartridge holding the diaphragm in position.

• **Diaphragm:** is a semipermeable membrane, usually latex rubber through which the cartridge end of the needle penetrates. The perforation of the needle is centrically located and round forming a tight ceal around the needle.



### **Cartridge contents**

The solution contained within the dental cartridge are:

- Local anaesthetic drug: the drugs contained within the cartridge are listed by their percent concentration. The number of milligrams of the agent can be calculated by multiplying the percent concentration(e.g. 2% = 20mg/ml) so a 1.8 ml cartridge of a 2% solution contains 36 mg.
- A vasopressor drug is included in some anaesthetic cartridges to increase the safety and the duration of action of the local anaesthetic. Cartridges containing vasopressor also contain a chemical that serves as an antioxidant most frequently used sodium bisulfite.
- Sodium chloride: is added to the contents of the cartridge to make the solution isotonic with the tissue of the body.
- Distilled water: is used as the diluent to provide the volume of solution in the cartridge

n special

## Care and handling

- Cartridge should be stored at room temperature .
- Cartridge should be not used after their expiry date.
- Cartridge should be listed by their percent concentration.

## **Clinical problems**

- Bubble in the cartridge: A small bubble of approximately 1-2 mm diameter will frequently be found in the dental cartridge. It is composed of nitrogen gas which was bubbled into the local anaesthetic solution during manufacture to prevent oxygen from being trapped in the cartridge and potentially destroying the vasopressor. A large bubble which may be present with a plunger that is extruded beyond the rim of the cartridge, is the result of the freezing of the anaesthetic solution, such cartridges should not be used, since sterility of the solution cannot be assured.
- Extruded stopper: The stopper can be extruded when a cartridge is frozen and the liquid inside expands. In this case the solution can no longer be considered sterile and should not be used for injection. Frozen cartridges can be identified by the presence of a large air babble more than 2mm by the extruded stopper. An extruded stopper without bubble is indicative of prolonged storage in a chemical disinfecting solution and diffusion of the solution into the cartridge.

# Burning on injection: A burning sensation on injection of anesthetic solution may be the result of one of the following:

- Normal response to pH of the drug: The pH of the dental cartridge containing vasopressor is lower (3.3-4) than that without vasopressor (5.5-6), so the plain anaesthetics have a somewhat more rapid onset of clinical action and more comfortable.
- In addition the inclusion of sodium bisulfite as antioxidant into the anaesthetic solution to prevent the biodegradation of the vasopressor by oxygen which might be present in the cartridge during manufacture or which can diffuse through the semipermeable diaphragm after tilting, reacts with oxygen before the oxygen can destroy the vasopressor.
- 2. Cartridge containing sterilizing solution
- 3. overheated cartridge
- 4. cartridge containing a vasopressor

- Leakage during injection
- Broken cartridge

### Preparation of the armamentarium

Proper care and handling of the local anaesthetic armamentarium can prevent or at least minimize the development of complications associated with the syringe, needle, and cartridge.

### Step of the preparation:

1. Remove the sterilized syringe from its container.

2. Retract the piston fully prior to attempting to load the cartridge.

3. Insert the cartridge while the piston is fully retracted into the syringe, insert the rubber stopper end of the cartridge.

4. Engage the hook push the piston forward until the hook is firmly engaged in the plunger.

5. Attach the needle to the syringe after removal of the plastic cap from the syringe end of the needle.

6. Carefully, remove the plastic protective cap from the opposite end of the needle and expel a few drops of solution to test for proper flow.

7. The syringe is now ready for use.



# **Additional armamentarium**

- Topical antiseptic: Such as 5% Betadine, Povidone-iodine
- Topical anesthetic: Such as 2% Lidocaine jelly, or 5% Benzocaine, ointment (Mucopain).
- Applicator sticks.
- Cotton/gauze.
- Hemostat: To remove needle from the soft tissue in the event of needle breakage.
- Small stainless steel bowls.
- Cartridge dispenser and warmer





# Local anesthesia in Oral Surgery by Dr. Elham H. Abdulkareem Lecture 1 2020-2021

# Table 1.2: Timewise development of various local anesthetic agents

Year	Esters	Amides	Discoverer
1905	Procaine		Alfred Einhorn
1943		Lidocaine	Nils Lofgren
1948			Applied in clinical practice
1952	Propoxycaine		Clinton and Laskowsky
1953		Prilocaine	Prepared by Lofgren and Tegner
1956-57		Mepivacaine	Prepared by AF Ekenstam
1960			Introduced in dentistry
1957		Bupivacaine	AF Ekenstam
1969		Articaine	H Rusching et al
1971		Etidocaine	Takman

There are two methods for obtaining anesthesia in dentistry, local and general.

 Local anesthesia: It is a method, whereby a certain operative area is made insensitive to pain, without loss of consciousness.

The sensory or the efferent nerves are blocked at the periphery or at any point between operative field and the center in the brain.

 General anesthesia: It is a method, whereby a certain operative area is rendered insensitive to pain, with loss of consciousness, by blocking brain function. "Anesthesia" means loss of all forms of sensation including pain, touch, temperature and pressure perception and may be accompanied by impairment of motor function.

Another term "analgesia" is used to denote loss of sensation unaccompanied by loss of other forms of sensation.

As a general rule, a larger dose of the agent is required to obtain anesthesia than to obtain analgesia. When only a part of the body is involved then the terms "local analgesia" and "local anesthesia" are used.

# **FACTORS AFFECTING NERVE CONDUCTION**

# **Diffusion of Solution**

Local anesthetic agents are alkaloid bases, which are combined with acids, usually hydrochloric acid, to form water-soluble salts. All local anesthetic salts are formed by a combination of a weak base and a strong acid

 The purpose of using salts is to render the salts stable and soluble in water. Water solubility is necessary for their diffusion through the interstitial fluids to the nerve fiber

# **Injection of Local Anesthetic Agents Close to the Nerve Membrane**

For maximum action, the anesthetic agent, should be injected sufficiently close to the target nerve, so that adequate concentration is available for diffusion into the nerve.

# **pH of the Tissue**

The presence of a low pH, as in areas of infection (pH of pus is 5.5 to 5.6); may interfere with achieving adequate anesthesia by preventing deprotonization and liberation of the free base.

# **Lipid Solubility and Protein Binding**

 Higher the lipid solubility and percentage of protein binding, more rapid and long lasting are the effects.

The addition of a butyl group to procaine results in the formation of tetracaine, an agent having 16 times the anesthetic activity and 4 times the duration of the compound.
#### Type of Nerve

- The myelinated nerves usually require a greater concentration of local anesthetic solution. These nerves also require more time for blocking them.
- The non-myelinated nerve-fibers require lesser concentration and lesser time.

#### Size of Nerve

The larger the diameter of nerve fiber, the greater is the concentration required to prevent impulse conduction.

Since pain is the only modality of sensation in the tooth, all sensations are adequately eliminated when pain fibers are anesthetised.

Analgesia is produced as a result of blockage of afferent transmission,

whereas **relaxation of skeletal muscles** and inhibition of autonomic innervated structures cause blockade from afferent transmission.

#### **Presence of a Vasoconstrictor**

Local anesthetic solutions containing a vasoconstrictor are less absorbed than those without a vasoconstrictor. The solution which are rapidly absorbed into cardiovascular system, there is less likelihood of developing an adequate block anesthesia.

#### Injection into a Vascular Area

A local anesthetic solution injected into a highly vascular area is rapidly absorbed in cardiovascular system. The rapid absorption reduces the effective concentration in the vicinity of the nerve and causes a rapid termination of analgesia.

#### **Presence of Infection**

Infection reduces alkalinity of tissues which retards the deprotonization of local anesthetic agents. This prevents liberation of free alkaloid base, which is necessary for the development of effective analgesia.

The effectiveness of local anesthetic agents depends on the following:

- 1. The chemical nature of the drug used.
- 2. The concentration of the drug used.
- 3. The volume of solution injected.
- 4. The rate of diffusion of both the anesthetic salt and the free base.

5. Addition of vasoconstrictor. It influences the time during which the free base remains in contact with the nerve.

## Indications

In general, local anesthesia is used to render the teeth and supporting structures insensitive to painful procedures. Specifically, it is used for the following:

## **Oral Surgery**

1. To make needle insertion painless

- 2. Extraction of teeth and fractured roots the
- 3. Odontectomy
- 4. Treatment of alveolalgia
- 5. Alveolectomy
- 6. Apicoectomy
- 7. Incision and drainage of localised abscesses

8. Removal of cysts, residual infection areas, hypertrophic tissues and neoplastic growths, Ranula and salivary calculi

9. In the treatment of tic douloureux by producing prolonged anesthesia with a combination of a local anesthetic agent and alcohol injection, for blocking the involved nerve.

10. A therapeutic test to localise the source of vague pain about the face.

Tic douloureux or trigeminal neuralgia is a severe, stabbing pain to one side of the face.

#### **Conservative Dentistry**

The following operative and restorative procedures:

- i. Cavity preparation
- ii. Crown and bridge abutment preparation
- iii. Pulpotomy or pulpectomy.

## Periodontology

i. Surgical treatment of periodontal diseases

- ii. Deep scaling and prophylaxis treatment
- iii. Mucogingival surgical procedures.

### **Prosthodontics**

Giving denture patients relief from sore spots which are painful even though dentures have been relieved.

**Orthodontia** Separation of teeth.

## Radiology

To prevent gagging and retching caused by the contact of film with palatal tissues and posterior part of oral cavity. These tissues or the areas are anesthetised before placing the film. In these cases usually surface anesthesia is used.

## CONTRAINDICATIONS

These can be divided into two groups:

(a) Absolute contraindications, and (b) Relative contraindications

#### **Absolute Contraindications**

#### History of Allergy to Local Anesthetic Agents

Local anesthetic agents belonging to the same chemical group should not be used. However, local anesthetic agents in the different chemical group can be used.

In case, a patient gives history of allergy to an amide local anesthetic agent, an ester local anesthetic agent can be used.

History of allergy to any of the constituents of the local anesthetic solution.

- Bisulfites, in the form of Sodium-bisulfite and sodiummetabisulfite are used as anti-oxidants or as preservatives to the vasoconstrictor in the local anesthetic solutions.
- Other general preservatives present in the local anesthetic solution **include thymol**, **methylparaben**, **and chlorbutol**.
- The solution containing the constituent should be avoided.
- An alternate solution should be used, if possible.

#### Relative Contraindications

1. Fear and apprehension: Where the patient is uncooperative or refuses for regional analgesia.

2. Presence of acute inflammation or suppurative infection at the site of insertion of the needle: There are increased chances of dissemination of infection with the passage of the needle from the abscess area to the deeper tissues.

3. Infants or small children: These patients lack reasoning and understanding.

4. Mentally retarded patients: These patients are unable to cooperate.

5. Restricted mouth opening: When the patient cannot open his mouth sufficiently, in situations, such as (i) trismus, or (ii) partial or complete ankylosis of temporomandibular joint.

6. Patients with significant medical disease: (a) cardiovascular disease, (b) hepatic dysfunction, (c) renal dysfunction, and (d) clinical hyperthyroidism.

a. Patients with significant cardiovascular disease: All local anesthetic

solutions containing high concentrations of vasoconstrictor, such as epinephrine, as in gingival retraction cords, should be avoided.

Local anesthetic agents containing higher dilution of epinephrine, such as 1:100 000 or 1:200 000, or 3% mepivacaine or 4% prilocaine can be used (for nerve blocks).

b. Patients with significant hepatic dysfunction: All local anesthetic agents belonging to amide group undergo biotransformation in liver (fixed function oxidases). These agents are best avoided, if not, should be used judiciously.

c. Patients with significant renal dysfunction: All amides and esters should be avoided, however, these can be used judiciously.

d. Patients with clinical hyperthyroidism: High concentrations of vasoconstrictor, as in epinephrine gingival retraction cords, should be avoided. Local anesthetic agents containing higher dilutions of epinephrine such as 1:100 000 or 1:200 000, or 3% mepivacaine or 4% prilocaine (nerve blocks) can be used.

7. Major surgical procedures

8. Presence of certain anomalies or developmental defects: These situations make regional analgesia difficult or impossible.

9. Presence of congenital Methemoglobinemia: It can be idiopathic or congenital.

a. Articaine and prilocaine when used in large doses, can produce Methemoglobinemia (Bellamy et al, 1992). These two drugs should be avoided in patients with congenital Methemoglobinemia, and other clinical syndromes with reduced O<sub>2</sub> carrying capacity of blood; because, there is increased risk of producing clinically significant Methemoglobinemia. These agents may be administered, if absolutely necessary; however, their dose should be minimised. The maximum recommended dose of prilocaine is 8 mg/kg body weight.

Methemoglobinemia is less likely to occur at lower doses. Alternatively, local anesthetic agents belonging to other groups both amides or esters can be used.

b. Benzocaine, a topical local anesthetic agent, when used in very large doses can also induce methemoglobinemia (Guertler and Pearce, 1994; and Rodriguez et al, 1994).

10. Presence of atypical plasma cholinesterase:

- Cholinesterase, is an enzyme, present in plasma, and is required for biotransformation of all esters by causing hydrolysis.
- These agents undergo biotransformation (hydrolysis) in the blood by the enzyme plasma cholinesterase, which is produced in the liver.
- In such situations, it is preferable to use an amide type of local anesthetic agent. Amides do not present any risk of high blood levels, because these agents undergo biotransformation in the liver.

# **ADVANTAGES**

- 1. Patient remains awake and cooperative.
- 2. Little distortion of normal physiology; therefore can be used in poor risk patients.
- 3. Low incidence of morbidity.
- 4. Patient can leave hospital unescorted.
- 5. Additional trained personnel not required.
- 6. Technique not difficult to master.
- 7. Percentage of failure is small.
- 8. No additional expense to the patient.
- 9. Patient need not miss the previous meal. In fact, should have one. Patients should not come on empty

# DISADVANTAGES

- No true disadvantage to the use of regional analgesia; when the patient is mentally prepared and when there are no contraindications.
- In every instance, when satisfactory anesthesia can be achieved and the patient is cooperative, regional analgesia is the method of choice.

#### References

1. Archer WH. A Manual of Dental Anesthesia: An illustrated guide for students and Practitioners. 2nd ed. WB Saunders Company 1958.

2. Bellamy MC, Hopkins PM, Hallsall PJ, Ellis FR. A study into the incidence of methaemoglobinemia after "3-in-1" block with prilocaine. Anaesthesia 1992;47(12):1084-85.

3. Guertler AT, Pearce WA. A prospective evaluation of benzocaine-associated methaemoglobinemia in human beings. Ann Emerg Med 1994;24(4):626-30.

4. Malamed SF. Handbook of Local Anesthesia. 4th ed. New Delhi: Harcourt India Private Limited (Mosby) Indian Reprint 2001.

5. Prilocaine-induced methaemoglobinemia-Wisconsin, 1993, MMWR Morb Mortal Wkly Rep 1994;43(35):655-57.

6. Richard Bennett C. Monheim's Local Anesthesia and Pain Control in Dental Practice. 7th ed. New Delhi: CBS Publishers and Distributors 1990.

7. Rodriguez LF, Smolik LM, Zbehlik AJ. Benzocaine-induced methaemoglobinemia: A report of a severe reaction and review of the literature. Ann Pharmacother 1994;28(5):643-49.

# Local Anesthetic Agents By Dr. Elham H. Abdulkareem 2020-2021

#### **PROPERTIES OF AN IDEAL LOCAL ANESTHETIC AGENT**

- The properties that are desirable in a local anesthetic solution are as follows:
- 1. It should be non-irritating and produce no local reaction to the tissues to which it is applied.
- 2. It should not cause any permanent change in the nerve structure.
- 3. It should cause minimal systemic toxicity.
- 4. It must be effective when injected into the tissues and should have sufficient penetrating properties to be effective as a topical anesthetic, when applied topically to the mucous membrane.
- 5. It should have a short time of onset, if possible.
- 6. The duration of action must be long enough to allow completion of procedure.
- Bennett (1974) has added some properties which are as follows:
- 1. It should have enough potency to give complete anesthesia without the use of harmful concentrated solutions.
- 2. It should be relatively free from producing allergic reactions.
- 3. It should be stable in solution and readily undergo biotransformation in the body.
- 4. It should either be sterile or be capable of being sterilized by heat without deterioration.

## **CLASSIFICATIONS**

The local anesthetic agents can be classified in various ways.

## **On the Basis of Occurrence in Nature**

- 1. Naturally occurring, e.g. cocaine
- 2. Synthetic compounds
- Freely soluble, e.g. procaine
- Poorly soluble, e.g. benzocaine

3. Miscellaneous drugs with local anesthetic action, e.g. clove oil, phenol, chlorpromazine, certain antihistaminics such as diphenhydramine.

## **On the Basis of Chemical Structure** *Esters*

These can be further classified as:

i. Esters of benzoic acid, e.g. cocaine, benzocaine (ethylaminobenzoate), and butacaine.

ii. Esters of para-aminobenzoic acid, e.g. procaine, chloroprocaine, and propoxycaine.

#### Amides

For example, articaine, bupivacaine, lidocaine, mepivacaine, and prilocaine.

#### **On the Basis of Duration of Action**

a. Short-acting: Articaine, lidocaine, mepivacaine, prilocaine, etc.

b. *Long-acting:* Bupivacaine, etidocaine, bucricaine, etc.

# Lignocaine (lidocaine, xylocaine, octocaine, dentocaine)

It is the most commonly used local anesthetic agent in dentistry.

## Chemistry

- It is an amide.
- It is stable, as it can be stored for a long time at room temperature.
- It withstands boiling and autoclaving.
- It is compatible with all types of vasoconstrictors.

## **Biotransformation**

Lidocaine undergoes biotransformation in liver.

a. Metabolism: It is metabolised in liver by microsomal fixed-function oxidases.

b. *Excretion:* Lidocaine and its breakdown products are excreted to some extent in urine, by kidney/

*Potency:* 2 times as potent as procaine. Today, it is taken as the standard for comparison of various other local anesthetic agents.

Toxicity: 2 times as toxic as procaine.

Time of onset of action: Rapid (2-3 minutes).

Duration of action: It depends upon:

i. Type of injection: Nerve block has longer duration than infiltration.

ii. Amount of vasoconstrictor used in the solution.

Effective dental concentration: 2%

Anesthetic half-life: 1.6 hours

Topical anesthetic action: It has topical anesthetic effect. It forms an excellent surface anesthetic. It is used in the following forms topically:

• 2% in the form of jelly

- 5% in the form of ointment
- 10% and 15% in the form of spray (aerosol).

#### Maximum Recommended Dose (MRD)

It is used in the concentration of 2%. It simply means 2 g of solute is contained in 100 ml of solution, which means 2000 mg of the solute are present in 100 ml of solution. Hence, it means 20 mg of the solute is contained in 1 ml of solution. In other words 1 ml of 2% lignocaine solution contains 20 mg of the local anesthetic agent (lignocaine).

the Maximum Recommended dose for lidocaine with or without a vasoconstrictor is 4.4 mg/kg body weight. The Maximum Recommended Doses are mentioned here as shown in Table 7.1, for individuals with their body weights ranging from 10 to 70 kg.

Table 7.1: The maximum recommended doses for various body wieghts					
Body weight (kg)	Dose (Body weight x 4.4) = mg	MRD (ml)	MRD (Cartridges) (approx)		
10	44	2.2	1.1		
20	88	4.4	2.2		
30	132	6.6	3.3		
40	176	8.8	4.4		
50	220	11	5.5		
60	264	13.2	6.6		
70	300	15.4	7.7		

### Availability in Dentistry

i. Dental cartridges: 2% lidocaine with epinephrine 1:80, 000 in India. ii. Vials:

a. 2% lidocaine, without epinephrine (Lidocaine Plain)

b. 2% lidocaine with epinephrine 1:80,000, 1:100,000, and 1:200,000.

#### **Toxicity**

The deleterious effects of toxic doses on various systems are as follows:

#### **Central Nervous System**

The manifestations of stimulation vary from mild restlessness to severe convulsions. The depression is manifested as drowsiness to loss of consciousness.

#### **Respiratory System**

a. In small doses it causes mild bronchodilatation.

b. In large doses it causes respiratory arrest (apnea) (the most common cause of death related to overdose of local anesthetic agent). In majority of cases, respiratory arrest precedes cardiac arrest.

#### Cardiovascular System

The effects on cardiovascular system vary with the dose used.

a. *Moderately large doses:* It produces overall inhibition on the contractility of heart muscle, in the form of:

i. A decrease in the electrical excitability of myocardium.

ii. A decrease in the force of contraction (negative inotropic effect).

iii. A decrease in the rate of electrical impulse conduction (negative chronotropic effect).

Table 7.6: Showing the comparison of various other local anesthetic agents				
Properties	Mepivacaine	Bupivacaine	Prilocaine	
Trade names	Scandonest, Polocaine, Carbocaine	Marcaine, Sensorcaine	Citanest, Citanest forte	
Chemical structure	Amide, with N- methyl group at hydrophilic end	Amide, with a butyl group replacing ethyl group at hydrophilic end	Amide, a derivative of toluidine	
Onset	1½ - 2 min	6-10 min	2-4 min	
Duration of action	2-3 hr	6-8 hr	2-3 hr	
Potency	Similar to lignocaine	4 times as potent as lignocaine	Similar to lignocaine	
Toxicity	Similar to lignocaine	Similar to lignocaine	Lesser than lignocaine	
Metabolism	By hepatic fixed- function oxidases	By hepatic amidases	By hepatic amidases	
Excretion	Kidney	Kidney	Kidney	
pН				
i. Plain solution	4.5	4.5-6.0	4.5	
ii. Solution with	3.0-3.5	3.0-4.5	3.0-4.0	
Effective dental	2% with	0.25% and 0.5%	3% and 4%	
concentration	vasoconstrictor and 3% without vasoconstrictor			
Anesthetic half-life	1.9 hours	2.7 hours	1.6 hours	
Maximum recom-	6.6 mg/kg BW,	1.3 mg/kg BW, not	6 mg/kg BW; not to	
mended dose	not to exceed 400 mg	to exceed 90 mg	exceed 400 mg	
Contraindications	2		<ul> <li>Patients with congenital or idiopathic methe- moglobulinemia</li> <li>Patients taking acetaminophen or phenacetin</li> </ul>	
Others uses		i. Spinal anesthesia,		
		ii. Epidural anesthesi	a	

# SELECTION OF LOCAL ANESTHETIC AGENTS

- There are several anesthetic agents available for use in dentistry. The agents are selected on the basis of the needs of a particular patient.
- 1. Duration of action
- 2. Need for control of postoperative pain
- 3. Physical and mental status of the patient
- 4. Concomitant medications.

#### **Duration of Action**

It is an important aspect to be considered. An approximate duration for completing of the surgical procedure for which anesthesia is required, should be taken into account. The working time should be adequate. On the basis of duration of action, various local anesthetic agents may be arbitrarily grouped as follows:

## 1. Ultra-short acting agents: Where the duration of action is less than 30 minutes:

- i. Procaine without a vasoconstrictor
- ii. 2-chloroprocaine (1.2% or 3%) without a vasoconstrictor
- iii. 2% lidocaine without a vasoconstrictor
- iv. 4% prilocaine without a vasoconstrictor for infiltration.

#### 2. Short-acting agents: Where the duration of action is 45 to 75 minutes.

- i. 2% lidocaine with 1:100 000 epinephrine
- ii. 2% mepivacaine with 1:200 000 levonordefrine

**3.** *Medium-acting agents:* Where the duration of action is 90 to 150 minutes: 4% prilocaine with 1:200,000 epinephrine

**4. Long-acting agents: Where the duration of action is 180 minutes or longer.** 0.5% bupivacaine with 1:200,000 epinephrine

#### **Need for Control of Postoperative Pain**

Most of the oral surgical procedures result in varying amount of postoperative pain. The local anesthetic agent serves as an additional medication that sometimes eliminates the need for postoperative analgesics. The local anesthetic agents that have shown to produce longer duration of analgesia even after the other sensations have returned are bupivacaine and etidocaine.

#### **Physical, Medical and Mental Status of the Patient**

- Any co-existing medical conditions, such as hypertension or diabetes mellitus should be considered.
- A patient with a history of allergy to a specific local anesthetic agent, or of any of the other components used along with the local anesthetic agents should be considered.
- Patients with a history of malignant hyperpyrexia, the use of amide derivatives is contraindicated.
- The mental status of the patients has to be evaluated. Small children and mentally challenged adults are sometimes fascinated with the numbness and the tingling sensation produced by the local anesthetic agents. These patients land up into traumatising their lip, tongue or cheeks intentionally or inadvertently. Hence for these reasons, the use of long acting local anesthetic agents is contraindicated in such patients.

## **Concomitant Medications**

The use of vasoconstrictors is contraindicated in patients who are taking monoamino-oxidase (MAO) inhibitors, tricyclic (TCA) antidepressants, etc.