

المادة محاضرات تحليل دالي ٢

الفصل الثاني

المرحلة الرابعة

القسم الرياضيات

الكلية التربية للعلوم الصرفة

الجامعة الانبار

المحاضرة الاولى

Example:

Every open and closed balls in normed space are convex. (31)

Sol. let X be normed space; $x, y \in B_r(x_0)$ and $0 \leq \lambda \leq 1 \Rightarrow \|x - x_0\| < r$ and $\|y - x_0\| < r$
we need to prove $\lambda x + (1-\lambda)y \in B_r(x_0)$.

$$\lambda x + (1-\lambda)y - x_0 = \lambda(x - x_0) + (1-\lambda)(y - x_0)$$

$$\begin{aligned} \Rightarrow \|\lambda x + (1-\lambda)y - x_0\| &= \|\lambda(x - x_0) + (1-\lambda)(y - x_0)\| \\ &\leq |\lambda| \|x - x_0\| + |1-\lambda| \|y - x_0\| \\ &< |\lambda| r + |1-\lambda| r = r \end{aligned}$$

because $|\lambda| = \lambda$ and $|1-\lambda| = 1-\lambda$ ($\lambda, 1-\lambda > 0$)

$\Rightarrow \lambda x + (1-\lambda)y \in B_r(x_0) \Rightarrow B_r(x_0)$ is convex.

Similarly, we can prove $\overline{B_r(x_0)}$ is convex.

Theorem:

Let $\{x_n\}$ and $\{y_n\}$ be two sequences in normed space X such that $x_n \rightarrow x$ and $y_n \rightarrow y$. Then

1. $x_n + y_n \rightarrow x + y$; 2. $\lambda x_n \rightarrow \lambda x$, $\forall \lambda \in F$.
3. $\|x_n\| \rightarrow \|x\|$ 4. $\|x_n - y_n\| \rightarrow \|x - y\|$
5. If $\{\lambda_n\}$ is a sequence in F such that $\lambda_n \rightarrow \lambda$, then $\lambda_n x_n \rightarrow \lambda x$.

Proof: (1) $\|(x_n + y_n) - (x + y)\| = \|(x_n - x) + (y_n - y)\|$

$$\leq \|x_n - x\| + \|y_n - y\|$$

(32)

Since $\|x_n - x\| \rightarrow 0$ and $\|y_n - y\| \rightarrow 0$ as $n \rightarrow \infty$

So, $\|(x_n + y_n) - (x + y)\| \rightarrow 0$, as $n \rightarrow \infty$, i.e.

$$x_n + y_n \rightarrow x + y$$

(3) Since $|\|x_n\| - \|x\|| \leq \|x_n - x\|$ and $\|x_n - x\| \rightarrow 0$ as $n \rightarrow \infty$, i.e.

$$\|x_n\| \rightarrow \|x\|.$$

$$(4) \quad |\|x_n - y_n\| - \|x - y\|| \leq \|(x_n - y_n) - (x - y)\|$$

$$\leq \|x_n - x\| + \|y_n - y\|$$

we have $\|x_n - x\| \rightarrow 0$ and $\|y_n - y\| \rightarrow 0$, as $n \rightarrow \infty$

Hence $|\|x_n - y_n\| - \|x - y\|| \rightarrow 0$ as $n \rightarrow \infty$.

$$(5) \quad \|\lambda_n x_n - \lambda x\| = \|\lambda_n x_n - \lambda_n x + \lambda_n x - \lambda x\|$$

$$= \|\lambda_n (x_n - x) + (\lambda_n - \lambda)x\|$$

$$\leq |\lambda_n| \|x_n - x\| + |\lambda_n - \lambda| \|x\|$$

Since $\|x_n - x\| \rightarrow 0$ and $|\lambda_n - \lambda| \rightarrow 0$, as $n \rightarrow \infty$

$\Rightarrow \|\lambda_n x_n - \lambda x\| \rightarrow 0$, as $n \rightarrow \infty$.

Example: let $X = F$, we define the function

$\|\cdot\|: X \rightarrow \mathbb{R}$ by $\|x\| = |x|$, $\forall x \in X$. show

that X is Banach space.

Sol.: First, we need to show that X is a normed space

• since $|x| \geq 0$, for all $x \in X \Rightarrow \|x\| \geq 0$.

• let $x \in X \Rightarrow \|x\| = 0 \Leftrightarrow |x| = 0 \Leftrightarrow x = 0$.

• let $x \in X$ and $\alpha \in F \Rightarrow \|\alpha x\| = |\alpha x| = |\alpha| |x|$ (33)
 $= |\alpha| \|x\|$.

• let $x, y \in X$, $\|x+y\| = |x+y| \leq |x| + |y|$

$\Rightarrow \|x+y\| \leq \|x\| + \|y\|$.

$\Rightarrow X$ is normed space.

Since \mathbb{R} or \mathbb{C} is complete space, F is complete space, hence X is Banach space.

Remark: let F^n denoted the set of all ordered n -tuples of elements in F of fixed $n \in \mathbb{N}$, i.e.

$F^n = \{x = (x_1, \dots, x_n); x_i \in F, i=1, 2, \dots, n\}$. Then

F^n is a vector space under the following addition and multiplication by scalar

1. $x+y = (x_1, \dots, x_n) + (y_1, \dots, y_n) = (x_1+y_1, \dots, y_n+x_n)$
 for all $x, y \in F^n$.

2. $\lambda x = \lambda (x_1, \dots, x_n) = (\lambda x_1, \dots, \lambda x_n), \forall x \in F^n, \lambda \in F$

Example: let $X = F^n$, we define the function

$\|\cdot\|: X \rightarrow \mathbb{R}$ by $\|x\| = \left(\sum_{i=1}^n |x_i|^2\right)^{\frac{1}{2}} \forall x \in F^n$.

Show that X is Banach space.

Sol.: First, we need to show that X is normed space

1. since $|x_i| \geq 0 \forall |i| \leq n \Rightarrow \|x\| \geq 0$

2. let $x \in X$, $\|x\| = 0 \Leftrightarrow \left(\sum_{i=1}^n |x_i|^2\right)^{\frac{1}{2}} = 0 \Leftrightarrow x_i^2 = 0$

For $i=1, 2, \dots, n$

So, $\|x\|=0 \Leftrightarrow x_i=0, \forall i=1, 2, \dots, n \Leftrightarrow x=0$

(34)

3. let $x \in X$ and $\lambda \in F \Rightarrow \|\lambda x\| = \left(\sum_{i=1}^n |\lambda x_i|^2 \right)^{1/2}$
 $= |\lambda| \left(\sum_{i=1}^n |x_i|^2 \right)^{1/2}$

4. let $x, y \in X \Rightarrow \|x+y\| = \left(\sum_{i=1}^n |x_i + y_i|^2 \right)^{1/2}$
 $= |\lambda| \|x\|$

By using Minkowski inequality $\leq \left(\sum_{i=1}^n |x_i|^2 \right)^{1/2} + \left(\sum_{i=1}^n |y_i|^2 \right)^{1/2}$
 $= \|x\| + \|y\|$

$\Rightarrow X$ is a normed space.

Second, to show that X is complete.

let $\{x_m\}$ be a Cauchy sequence in X , $\exists k \in \mathbb{Z}^+$ \Rightarrow

$$\|x_m - x_l\| < \epsilon, \forall m, l > k$$

$$\Rightarrow \|x_m - x_l\|^2 < \epsilon^2 \quad \forall m, l > k \quad \text{--- (1)}$$

Since $x_m - x_l = (x_1^m - x_1^l, x_2^m - x_2^l, \dots, x_n^m - x_n^l)$

because $x_m \in F^n \Rightarrow x_m = (x_1^m, x_2^m, \dots, x_n^m)$

$$\text{So, } \|x_m - x_l\|^2 = \sum_{i=1}^n |x_i^m - x_i^l|^2 \quad \text{--- (2)}$$

From (1) or (2) $\sum_{i=1}^n |x_i^m - x_i^l|^2 < \epsilon^2 \quad \forall m, l > k$

$$\Rightarrow |x_i^m - x_i^l| < \epsilon \quad \forall m, l > k$$

$$\Rightarrow |x_i^m - x_i^l| < \epsilon \quad \forall m, l > k$$

So that for each i , the sequence $\{x_m\}$ is Cauchy sequence in F .

المحاضرة الثانية

Since F is complete, then for each i , the (35)
sequence $\{x_m\}$ converges to a point, say $x_i \in F$

$$\Rightarrow x_m \rightarrow x_i \quad \forall 1 \leq i \leq n$$

$$\text{put } x = (x_1, \dots, x_n) \Rightarrow x \in F^n$$

we must prove $x_m \rightarrow x$

$$\text{let } \epsilon > 0, \text{ for all } m > k, \text{ we have } \|x_m - x\|^2 = \sum_{i=1}^n |x_m^i - x_i|^2 < \epsilon^2$$

$$\Rightarrow \|x_m - x\| < \epsilon \quad \forall m, k > k$$

$$\Rightarrow \{x_m\} \text{ converges to } x \in F^n = X$$

$$\Rightarrow X \text{ is Complete} \Rightarrow X \text{ is Banach space.}$$

Example: let $X = \mathbb{R}^n$, we define the function

$$\|\cdot\|: X \rightarrow \mathbb{R} \text{ by } \|x\| = \sum_{i=1}^n |x_i|.$$

for all $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. show that X

is Banach space.

Sol. First, we need to show that X is normed space.

1. since $|x_i| \geq 0 \quad \forall 1 \leq i \leq n \Rightarrow \|x\| \geq 0$

2. let $x \in X, \|x\| = 0 \Leftrightarrow \sum_{i=1}^n |x_i| = 0$

$$\Leftrightarrow |x_i| = 0, \quad \forall 1 \leq i \leq n$$

$$\Leftrightarrow x_i = 0, \quad \forall 1 \leq i \leq n$$

$$\Leftrightarrow x = 0.$$

3. let $x \in X$ and $\lambda \in \mathbb{R}$

(36)

$$\lambda x = \lambda(x_1, \dots, x_n) = (\lambda x_1, \dots, \lambda x_n)$$

$$\Rightarrow \|\lambda x\| = \sum_{i=1}^n |\lambda x_i| = |\lambda| \sum_{i=1}^n |x_i| = |\lambda| \|x\|.$$

4. let $x, y \in X$

$$\begin{aligned} \|x+y\| &= \sum_{i=1}^n |x_i + y_i| \leq \sum_{i=1}^n (|x_i| + |y_i|) \\ &= \sum_{i=1}^n |x_i| + \sum_{i=1}^n |y_i| \end{aligned}$$

$$\Rightarrow \|x+y\| \leq \|x\| + \|y\|.$$

$\Rightarrow X$ is normed space

Second, since $X = \mathbb{R}^n$ is complete space

$\Rightarrow X$ is Banach space.

Example: let $X = \mathbb{R}^n$, we define the function

$$\|\cdot\|: X \rightarrow \mathbb{R} \text{ by } \|x\| = \max\{|x_1|, |x_2|, \dots, |x_n|\}$$

for all $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. Then X is Banach space.

Remark: let $C[a, b]$ be the set of all real-valued bounded continuous functions, defined on $[a, b]$, i.e.

$f \in C[a, b]$ iff $f: [a, b] \rightarrow \mathbb{R}$ is bounded and conts. function. Then $C[a, b]$ is vector space

under the following addition and scalar multiplication

1. $(f+g)(x) = f(x) + g(x), \forall f, g \in C[a, b].$

2. $(\alpha f)(x) = \alpha f(x), \forall f \in C[a, b], \alpha \in \mathbb{F}.$

Example: let $X = C[a, b]$, we define the function (37)

$\| \cdot \| : X \rightarrow \mathbb{R}$ by $\|f\| = \max \{ |f(x)| ; a \leq x \leq b \}$

for all $f \in X$. Prove that X is ~~normed~~ ^{normed} space.

Sol. First, we need to show that is a normed space.

1. Since $|f(x)| \geq 0, \forall x \in [a, b] \Rightarrow \|f\| \geq 0$.

$$\begin{aligned} 2. \|f\| = 0 &\Leftrightarrow \max \{ |f(x)| ; a \leq x \leq b \} = 0 \\ &\Leftrightarrow |f(x)| = 0, x \in [a, b] \\ &\Leftrightarrow f(x) = 0, x \in [a, b] \\ &\Leftrightarrow f = 0 \end{aligned}$$

3. let $f \in X$ and $\alpha \in \mathbb{R}$

$$\begin{aligned} \|\alpha f\| &= \max \{ |(\alpha f)(x)| ; a \leq x \leq b \} \\ &= \max \{ |\alpha f(x)| ; a \leq x \leq b \} \\ &= \max \{ |\alpha| |f(x)| ; a \leq x \leq b \} \\ &= |\alpha| \max \{ |f(x)| ; a \leq x \leq b \} \\ &= |\alpha| \|f\|. \end{aligned}$$

$$\begin{aligned} 4. \|f+g\| &= \max \{ |(f+g)(x)| ; a \leq x \leq b \} \\ &= \max \{ |f(x) + g(x)| ; a \leq x \leq b \} \\ &\leq \max \{ |f(x)| + |g(x)| ; a \leq x \leq b \} \\ &= \max \{ |f(x)| ; a \leq x \leq b \} + \max \{ |g(x)| ; a \leq x \leq b \} \\ &= \|f\| + \|g\|. \end{aligned}$$

whenever $f, g \in X$.

$\Rightarrow X$ is normed space

Example: let $X = C[0, 1]$, we define the (38)
function $\|\cdot\|: X \rightarrow \mathbb{R}$, by $\|f\| = \int_0^1 |f(x)| dx$ for
all $f \in X$, $x \in [0, 1]$. Show that X is normed space
but not Banach space.

Sol. First, to show X is a normed space.

1. Since $|f(x)| \geq 0$ for all $x \in [0, 1] \Rightarrow \|f\| \geq 0$.

2. (i) If $f = 0$, then $\int_0^1 |f(x)| dx = 0 = \|f\|$.

(ii) If $\|f\| = 0$, then $\int_0^1 |f(x)| dx = 0$

Since $|f(x)| \geq 0$ and f is cont., then $|f(x)| = 0 \Rightarrow f = 0$.

3. Let $f \in X$ and $\alpha \in \mathbb{R}$

$$\begin{aligned}\|\alpha f\| &= \int_0^1 |(\alpha f)(x)| dx = \int_0^1 |\alpha f(x)| dx \\ &= \int_0^1 |\alpha| |f(x)| dx = |\alpha| \int_0^1 |f(x)| dx \\ &= |\alpha| \|f\|.\end{aligned}$$

4. Let $f, g \in X$

$$\begin{aligned}\|f+g\| &= \int_0^1 |(f+g)(x)| dx = \int_0^1 |f(x)+g(x)| dx \\ &\leq \int_0^1 (|f(x)| + |g(x)|) dx \\ &= \int_0^1 |f(x)| dx + \int_0^1 |g(x)| dx \\ &= \|f\| + \|g\|.\end{aligned}$$

$\Rightarrow X$ is normed space.

We now show that X is not complete.

المحاضرة الثالثة

Consider the sequence $\{f_n\}$ in X defined as follows (39)

$$f_n(x) = \begin{cases} 1, & 0 \leq x \leq \frac{1}{2} \\ -nx + \frac{1}{2}n + 1, & \frac{1}{2} < x \leq \frac{1}{2} + \frac{1}{n} \\ 0, & \frac{1}{2} + \frac{1}{n} < x \leq 1 \end{cases}$$

Then $\{f_n\}$ is a Cauchy sequence in X , because, if $m > n \gg 3$, then

$$\begin{aligned} \|f_m - f_n\| &= \int_0^1 |(f_m - f_n)(x)| dx = \int_0^1 |f_m(x) - f_n(x)| dx \\ &= \int_0^{\frac{1}{2}} |1 - 1| dx + \int_{\frac{1}{2}}^{\frac{1}{2} + \frac{1}{n}} |f_m(x) - f_n(x)| dx \end{aligned}$$

$$\begin{aligned} \|f_m - f_n\| &\leq \int_{\frac{1}{2}}^{\frac{1}{2} + \frac{1}{n}} |f_m(x)| dx + \int_{\frac{1}{2}}^{\frac{1}{2} + \frac{1}{n}} |f_n(x)| dx \\ &= \int_{\frac{1}{2}}^{\frac{1}{2} + \frac{1}{n}} |-mx + \frac{1}{2}m + 1| dx + \int_{\frac{1}{2}}^{\frac{1}{2} + \frac{1}{n}} |-nx + \frac{1}{2}n + 1| dx \end{aligned}$$

Since $-mx + \frac{1}{2}m + 1 > 0$, when $\frac{1}{2} < x < \frac{1}{2} + \frac{1}{n}$

$$\|f_m - f_n\| \leq \frac{1}{2m} + \frac{1}{2n} \Rightarrow \|f_m - f_n\| \rightarrow 0 \text{ as}$$

$\Rightarrow \{f_n\}$ is Cauchy sequence. But this sequence is not convergent in X .

For, if there existed a $f \in X$ such that $f_n \rightarrow f$

$$\Rightarrow f(x) = \begin{cases} 1, & 0 \leq x < \frac{1}{2} \\ 0, & \frac{1}{2} < x \leq 1 \end{cases} \text{ This contradiction because } f \text{ is not cont.}$$

Theorem: let M be a subspace of Banach space X . Then M is Banach space if and only if it is closed in X .

Proof: suppose M is a Banach space and to prove that M is closed i.e. $M = \overline{M}$

we have always $M \subset \overline{M} \dots \textcircled{1}$

let $x \in \overline{M}$, there is a sequence $\{x_n\}$ in M such that

$$x_n \rightarrow x$$

$\Rightarrow \{x_n\}$ is Cauchy sequence in M

we have M is complete

$\Rightarrow x_n \rightarrow x \in M$, because the convergent point is unique.

$$\Rightarrow \overline{M} \subset M \dots \textcircled{2}$$

From (1) and (2), we get $M = \overline{M} \Rightarrow M$ is closed.

conversely, Assume that M is closed set in X .

let $\{x_n\}$ be a Cauchy sequence in M .

Since $M \subset X \Rightarrow \{x_n\}$ is a Cauchy sequence in X

Since X is complete space, there is $x \in X \ni x_n \rightarrow x$

So, $x_n \in M \Rightarrow x \in \overline{M}$, M is closed i.e. $M = \overline{M}$

$$\Rightarrow x \in M$$

$\Rightarrow \{x_n\}$ is Cauchy sequence converge in M

$\Rightarrow M$ is complete $\Rightarrow M$ is Banach space.

Theorem: Every finite dimensional normed space ⁽⁴¹⁾ is complete.

Proof: let X be a finite dimensional normed space with $\dim X = n > 0$ and let $\{e_1, e_2, \dots, e_n\}$ be a basis for X , take $\{x_m\}$ any Cauchy sequence in X , i.e.

$$\|x_m - x_k\| \rightarrow 0 \text{ as } m, k \rightarrow \infty \quad \dots (1)$$

$$\text{Since } x_m, x_k \in X \Rightarrow x_m = \sum_{i=1}^n \alpha_i^m e_i; \alpha_i^m \in F$$

$$\text{also } x_k = \sum_{i=1}^n \alpha_i^k e_i, \alpha_i^k \in F$$

$$\Rightarrow x_m - x_k = \sum_{i=1}^n (\alpha_i^m - \alpha_i^k) e_i$$

Since $\{e_1, \dots, e_n\}$ is linear independent, by Lemma of linear combination, there is $c > 0$ such that

$$\|x_m - x_k\| = \left\| \sum_{i=1}^n (\alpha_i^m - \alpha_i^k) e_i \right\| \geq c \sum_{i=1}^n |\alpha_i^m - \alpha_i^k| \quad \dots (2)$$

From (1) and (2), we have $\sum_{i=1}^n |\alpha_i^m - \alpha_i^k| \rightarrow 0$ as $m, k \rightarrow \infty$ for $i=1, 2, \dots, n$

$$\Rightarrow |\alpha_i^m - \alpha_i^k| \rightarrow 0 \text{ as } m, k \rightarrow \infty \text{ for } i=1, 2, \dots, n$$

For $i=1, 2, \dots, n \Rightarrow \{\alpha_i^m\}$ is Cauchy sequence in F .

Since F is either \mathbb{R} or \mathbb{C} and each for \mathbb{R} or \mathbb{C} are complete $\Rightarrow \exists \alpha_i \in F \ni \alpha_i^m \rightarrow \alpha_i$.

$$\text{Put } x = \sum_{i=1}^n \alpha_i e_i \Rightarrow x_m \rightarrow x, x \in X$$

$\Rightarrow X$ is complete.

Corollary: Every finite dimensional subspace M (42) of a normed space X is closed.

Proof: Since M is a finite dimensional subspace of a normed space $X \Rightarrow M$ is complete space $\Rightarrow M$ is closed.

Note: The infinite dimensional subspace of Banach space need not be closed.

Definition: Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be two norms on a vector space X . We say that $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent (or $\|\cdot\|_1$ is equivalent to $\|\cdot\|_2$), written $\|\cdot\|_1 \cong \|\cdot\|_2$ if there exist positive real numbers a and b such that $a\|x\|_1 \leq \|x\|_2 \leq b\|x\|_1$ for all $x \in X$.

Example: Let $\|x\|_1 = \sum_{i=1}^n |x_i|$ and $\|x\|_2 = \left(\sum_{i=1}^n x_i^2\right)^{1/2}$ for all $x \in \mathbb{R}^n$. Show that $\|\cdot\|_1 \cong \|\cdot\|_2$.

Sol. From Cauchy's inequality, we have $\sum_{i=1}^n |x_i y_i| \leq \left(\sum_{i=1}^n x_i^2\right)^{1/2} \left(\sum_{i=1}^n y_i^2\right)^{1/2}$ for all $x_i, y_i \in \mathbb{R}$, $1 \leq i \leq n$.

Put $y_i = 1$ for all $i = 1, 2, \dots, n$, we have

$$\sum_{i=1}^n |x_i| \leq \left(\sum_{i=1}^n x_i^2\right)^{1/2} \cdot \left(\sum_{i=1}^n 1\right)^{1/2}$$

So, $\|x\|_1 \leq \|x\|_2 \cdot \sqrt{n} \Rightarrow \frac{1}{\sqrt{n}} \|x\|_1 \leq \|x\|_2$

Set $a = \frac{1}{\sqrt{n}}$ and $b = 1$, we have $\|x\|_2 \leq \|x\|_1$.

Hence $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent i.e.

$$\|\cdot\|_1 \cong \|\cdot\|_2.$$

المحاضرة الرابعة

Lemma (Linear Combination)

(43)

Let $\{x_1, x_2, \dots, x_n\}$ be a linear independent set of vectors in normed space X . Then there is a number $c > 0$, such that $\|\sum_{i=1}^n \alpha_i x_i\| \geq c \sum_{i=1}^n |\alpha_i|$, for all $\alpha_i \in F, i = 1, 2, \dots, n$.

Theorem: On a finite dimensional vector space all norms are equivalent.

Proof: Let X be a finite dimensional vector space with $\dim X = n > 0$ and $\|\cdot\|_1, \|\cdot\|_2$ be two norms on X .

To prove $\|\cdot\|_1 \cong \|\cdot\|_2$.

Let $\{e_1, e_2, \dots, e_n\}$ be a basis for $X \Rightarrow \forall x \in X$ has a unique representation $x = \sum_{i=1}^n \alpha_i e_i, \alpha_i \in F$

$$\begin{aligned} \|x\|_1 &= \left\| \sum_{i=1}^n \alpha_i e_i \right\|_1 \leq \sum_{i=1}^n |\alpha_i| \|e_i\|_1 \\ &\leq \max \|e_i\|_1 \sum_{i=1}^n |\alpha_i| \end{aligned}$$

$$\text{set } \max \|e_i\|_1 = k \Rightarrow \|x\|_1 \leq k \sum_{i=1}^n |\alpha_i| \dots \text{ (1)}$$

since $\{e_1, e_2, \dots, e_n\}$ is basis for X and using L.C

$$\text{Lemma, } \exists c > 0, \exists \left\| \sum_{i=1}^n \alpha_i e_i \right\|_2 \geq c \sum_{i=1}^n |\alpha_i|$$

$$\Rightarrow \|x\|_2 \geq c \sum_{i=1}^n |\alpha_i| \dots \text{ (2)}$$

From (1) and (2), we get

$$\frac{k}{c} \|x\|_1 \leq \sum_{i=1}^n |\alpha_i| \leq \|x\|_2$$

$$\Rightarrow \frac{k}{c} \|x\|_1 \leq \|x\|_2 \dots \text{ (3)}$$

Similarly, we can

$$\|x\|_2 \leq \frac{c}{k} \|x\|_1 \quad \text{--- (4)}$$

From (3) and (4), we obtain

$$a = \frac{k}{c} \|x\|_1 \leq \|x\|_2 \leq b = \frac{c}{k} \|x\|_1$$

$$\Rightarrow \|\cdot\|_1 \cong \|\cdot\|_2$$

Continuity

Def. Let X and Y be two normed spaces. A function $f: X \rightarrow Y$ is called

1. Continuous at $x_0 \in X$, if for each $\epsilon > 0$, there is $\delta > 0$ such that $x \in X$, $\|x - x_0\| < \delta \Rightarrow \|f(x) - f(x_0)\| < \epsilon$.
or equivalently, a function f is continuous at $x_0 \in X$, if for every sequence $\{x_n\}$ in X converging to x_0 , the sequence $\{f(x_n)\}$ in Y converges to $f(x_0) \in Y$ i.e.

$$x_n \rightarrow x_0 \Rightarrow f(x_n) \rightarrow f(x_0)$$

2. Compact if $f(X)$ contained in compact subset of Y .

3. Completely continuous if it is both continuous and compact.

4. Finite dimensional if it is compact function and $f(X)$ contained in a finite dimensional subspace of Y .

Theorem:

(45)

Let X be a normed space. Then the function $f: X \rightarrow \mathbb{R}$, $f(x) = \|x\|$ is continuous, the norm $\|\cdot\|$ on X is cont. function.

Proof: let $x_0 \in X$ and $\{x_n\}$ seq. in $X \ni x_n \rightarrow x_0$ as $n \rightarrow \infty$.

$$\text{Now, } |f(x_n) - f(x_0)| = |\|x_n\| - \|x_0\|| \leq \|x_n - x_0\|$$

Since $x_n \rightarrow x_0 \Rightarrow \|x_n - x_0\| \rightarrow 0$ as $n \rightarrow \infty$
 $\Rightarrow |f(x_n) - f(x_0)| \rightarrow 0$ as $n \rightarrow \infty$

$$\Rightarrow f(x_n) \rightarrow f(x_0)$$

$\Rightarrow f$ is cont. at x_0 and x_0 is arbitrary point.
 $\Rightarrow f$ is cont. on X .

Theorem: let X be a normed space. Then functions $f: X \times X \rightarrow X$, $f(x, y) = x + y$ and $g: F \times X \rightarrow X$, $g(\alpha, x) = \alpha x$ are continuous, in other words, vector addition and scalar multiplication are jointly continuous.

Proof: let $x_0, y_0 \in X$ and $\{x_n\}, \{y_n\}$ in X such that $x_n \rightarrow x_0$ and $y_n \rightarrow y_0$ as $n \rightarrow \infty$.

$$\text{Now: } \|f(x_n, y_n) - f(x_0, y_0)\| = \|(x_n + y_n) - (x_0 + y_0)\|$$

$$= \|(x_n - x_0) + (y_n - y_0)\|$$

(46)

$$\leq \|x_n - x_0\| + \|y_n - y_0\|$$

$$\leq \rightarrow 0 + \rightarrow 0 = 0, \text{ as } n \rightarrow \infty$$

$$\Rightarrow f(x_n, y_n) \rightarrow f(x_0, y_0) \text{ as } n \rightarrow \infty$$

$\Rightarrow f$ is continuous at (x_0, y_0) and (x_0, y_0) is any point in $X \times X$, hence f is continuous.

Also, let $x_0 \in X$, $\alpha \in F$ and $\{x_n\}$ in X , $\{\alpha_n\}$ in F such that $x_n \rightarrow x_0$ and $\alpha_n \rightarrow \alpha$ as $n \rightarrow \infty$

$$\text{Now, } \|g(\alpha_n, x_n) - g(\alpha, x_0)\| = \|\alpha_n x_n - \alpha x_0\|$$

$$= \|\alpha_n x_n - \alpha_n x_0 + \alpha_n x_0 - \alpha x_0\|$$

$$= \|\alpha_n (x_n - x_0) + (\alpha_n - \alpha) x_0\|$$

$$\leq |\alpha_n| \|x_n - x_0\| + |\alpha_n - \alpha| \|x_0\|$$

Since $\|x_n - x_0\| \rightarrow 0$ and $|\alpha_n - \alpha| \rightarrow 0$ as $n \rightarrow \infty$, we have $\|g(\alpha_n, x_n) - g(\alpha, x_0)\| \rightarrow 0$ as $n \rightarrow \infty$

$$g(\alpha_n, x_n) \rightarrow g(\alpha, x_0), \text{ as } n \rightarrow \infty.$$

g is continuous at (α, x_0) and (α, x_0) is any point in

$F \times X$, hence g is continuous.

Corollary:

Every normed space X is topological linear space.

المحاضرة الخامسة

Continuous Linear Functions:

(47)

Recall that a function $f: X \rightarrow Y$ from a linear space X into linear space Y is called a linear if:

$$f(\alpha x + \beta y) = \alpha f(x) + \beta f(y), \text{ for all } x, y \in X \text{ and } \alpha, \beta \in F.$$

Remarks:

i. Linear function of linear space X into field F is called linear functional on X .

ii. Let $L(X, Y)$ denote the set of all linear functions from a linear space X into a linear space Y . Then $L(X, Y)$ is a vector space under the following addition and scalar multiplication

1. $(f + g)(x) = f(x) + g(x), \forall f, g \in L(X, Y).$

2. $(\alpha f)(x) = \alpha f(x), \forall f \in L(X, Y) \text{ and } \alpha \in F.$

If $Y = F$, we write $L(X)$ instead of $L(X, F)$. The space of all linear functionals defined on a linear space X is called the algebraic dual space and denoted by X' , i.e. $X' = L(X, F)$.

3. We say that X, Y are linear isomorphic (we write $X \cong Y$), then there is a bijective linear function

$f: X \rightarrow Y$ such function is called linear isomorphism.

Theorem:

(48)

Let X be a linear space over a field F .

1. If $x \in X$ and a function $T_x: X' \rightarrow F$ defined by $T_x(f) = f(x)$, for all $f \in X'$, then T_x is linear functional i.e. $T_x \in X''$ and it is called Evaluation functional induced by x .

2. If the function $\psi: X \rightarrow X''$ defined by $\psi(x) = T_x$ for all $x \in X$, then ψ injection linear function and ψ is called canonical function.

Proof: (1) let $f, g \in X'$, $\alpha, \beta \in F$

$$\begin{aligned} T_x(\alpha f + \beta g) &= (\alpha f + \beta g)(x) = (\alpha f)(x) + (\beta g)(x) \\ &= \alpha f(x) + \beta g(x) = \alpha T_x(f) + \beta T_x(g). \end{aligned}$$

$$\Rightarrow T_x \in X''.$$

(2) let $x, y \in X$, $\alpha, \beta \in F$

$$\Rightarrow \psi(\alpha x + \beta y) = T_{\alpha x + \beta y}$$

$$\begin{aligned} \text{for all } f \in X', \quad T_{\alpha x + \beta y}(f) &= f(\alpha x + \beta y) \\ &= \alpha f(x) + \beta f(y) \\ &= \alpha T_x(f) + \beta T_y(f) \\ &= (\alpha T_x + \beta T_y)(f) \end{aligned}$$

$$\Rightarrow \psi(\alpha x + \beta y) = \alpha T_x + \beta T_y = \alpha \psi(x) + \beta \psi(y).$$

$\Rightarrow \psi$ is linear function.

Now, to prove that ψ is injection, let $x, y \in X$ such that $\psi(x) = \psi(y)$

$$\Rightarrow T_x = T_y \Rightarrow T_x(f) = T_y(f) \text{ for } f \in X' \quad (49)$$

$$\Rightarrow f(x) = f(y) \text{ for all } f \in X'$$

$$\Rightarrow f(x-y) = 0 \text{ for all } f \in X'$$

$$\Rightarrow x-y = 0 \Rightarrow x=y \Rightarrow \psi \text{ is injective.}$$

Definition: Let X be a linear space over a field F .

We say that X is an Algebraically Reflexive if

ψ is an onto, where ψ is defined above theorem.

Theorem: Every finite dimensional space is algebraically reflexive.

Proof: Let X be a finite dimensional space over a field F . $\Rightarrow \dim X' = \dim X$, so that X' finite dimensional. $\Rightarrow \dim X'' = \dim X$, so that X'' finite dimensional.

Since $\psi: X \rightarrow X''$ is injective and X', X'' are finite dimensional and $\dim X'' = \dim X$, then ψ is onto.

Remark:

Recall that a function f from a topological space X into topological space Y , i.e. $f: X \rightarrow Y$ is called continuous at a point $x \in X$ if every neighborhood U of $f(x)$ in Y there is a neighborhood V of x in $X \ni$

$f(V) \subset U$. If f is continuous at every point, it is called continuous. A function $f: X \rightarrow Y$ is continuous iff each open (resp. closed) set U in Y the set $f^{-1}(U)$ is open (resp. closed) set in X .

Def. let (X, d) and (Y, d^*) be metric spaces. (50)

A function $f: X \rightarrow Y$ is called an Isometry if

- (1) f is bijective.
- (2) $d^*(f(x), f(y)) = d(x, y)$ for all $x, y \in X$.

Def. let X and Y be a normed spaces. An isometric isomorphism of X into Y is a one-one linear function f of X into Y such that $\|f(x)\| = \|x\|$ for every $x \in X$. Also we say that X is isometrically isomorphic or (congruent) to Y if there exists an isomorphism of X onto Y .

Remark: let f be an isometric isomorphism of X into Y where X and Y are normed spaces. let $x, y \in X$. Then $\|f(x) - f(y)\| = \|f(x - y)\| = \|x - y\|$. Thus f preserves distances and so it is an isometry.

Def. let X and Y be normed spaces. A topological isomorphism of X into Y is a 1-1 linear function f of X into Y such that f and f^{-1} are continuous on their respective domains. Also we say that X is topologically isomorphic to Y if there exists a topological isomorphism of X into Y . In other words, X and Y are topologically isomorphic provided there exists a homeomorphism of X onto which is also a linear function.

Remark: Topological isomorphism space need not be isometrically isomorphic. In fact there do exist examples of pairs spaces which are topologically isomorphic but not congruent.

المحاضرة السادسة

Theorem:

let X and Y be a normed spaces. Then X and Y are topologically isomorphic iff there exists a linear function of X onto Y and positive constant α, β such that $\alpha \|x\| \leq \|f(x)\| \leq \beta \|x\|$.

Proof: suppose X and Y are topologically isomorphic, then there exist a linear function f of X onto Y such that f and f^{-1} are continuous.

since f is cont^s iff there exists a positive constant α such that $\|f(x)\| \leq \alpha \|x\|$ for all $x \in X$.

Again f^{-1} is cont^s iff there exists a positive constant β such that $\beta \|x\| \leq \|f(x)\|$, for all $x \in X$.

It follows that X and Y are topologically isomorphic iff there exists a linear function of X onto Y and positive constants α and β such that $\beta \|x\| \leq \|f(x)\| \leq \alpha \|x\|$.

Theorem: let X and Y be topological linear spaces and let $f: X \rightarrow Y$ be a linear function. If f is cont^s at 0, then it is continuous.

Proof: let $x \in X$ and U be neighborhood of $f(x)$ in Y .

Then $U = f(x) + W$, where W is neighborhood of 0 in Y .

Since f is cont^s at 0 in X , then there exist a neighbor.

V of 0 in X such that $f(V) \subset W \Rightarrow x+V$ is neighbor of x in X .

To show that $f(x+V) \subset U$.

let $z \in f(x+V) \Rightarrow \exists y \in x+V \ni f(y) = z$ (52)
 Since $y \in x+V \Rightarrow y-x \in V \Rightarrow f(y-x) \in f(V)$
 $\Rightarrow f(y) - f(x) \in f(V) \Rightarrow z - f(x) \in f(V)$
 $\Rightarrow z \in f(x) + f(V) \Rightarrow z \in U$
 $\Rightarrow f(x+V) \subset U$
 $\Rightarrow f$ is cont'. at x , x is arbitrary point
 $\Rightarrow f$ is cont'.

Theorem: let X and Y be a normed spaces and
 let $f: X \rightarrow Y$ be a linear function. Then f is cont'.
 either at every point of X or no point of X .

Proof: let x_1 and x_2 be any two points of X and
 suppose f is cont'. at x_1 . Then to each $\epsilon > 0$, there
 exists $\delta > 0 \ni \|x - x_1\| < \delta \Rightarrow \|f(x) - f(x_1)\| < \epsilon$.

Now, $\|x - x_2\| < \delta \Rightarrow \|(x + x_1 - x_2) - x_1\| < \delta$
 $\Rightarrow \|f(x + x_1 - x_2) - f(x_1)\| < \epsilon$
 $\Rightarrow \|f(x) + f(x_1) - f(x_2) - f(x_1)\| < \epsilon$
 $\Rightarrow \|f(x) - f(x_2)\| < \epsilon$

$\Rightarrow f$ is cont' at x_2 , then f is cont'.

Theorem: let X and Y be a Banach spaces. If
 $f: X \rightarrow Y$ is cont', linear and onto function, then
 f is open.

Proof: let G be open set in X . We want to
 show that $f(G)$ is open in Y .

Let $y \in f(G)$, then $y = f(x)$ for some $x \in G$ (53)
 Since G is open set in X , there is $r > 0 \ni B_r(x) \subset G$
 $\Rightarrow f(B_r(x)) \subset f(G)$.

Since $B_r(x) = x + B_r(0) \Rightarrow x + B_r(0) \subset G$

By Lemma, there is an open sphere $B'_r(0)$ in Y center at origin such that $B'_r(0) \subset f(B_r(0))$

$$\begin{aligned} \Rightarrow y + B'_r(0) &\subseteq y + f(B_r(0)) = f(x) + f(B_r(0)) \\ &= f(x + B_r(0)) = f(B_r(x)) \subset f(G) \end{aligned}$$

since $y + B'_r(0) = B'_r(y) \Rightarrow B'_r(y) \subset f(G)$

$\Rightarrow f(G)$ is open, thus f is an open.

Def.: Let X and Y be any non-empty sets and let

$f: X \rightarrow Y$ be a function. The set

$$\{(x, y) \in X \times Y, y = f(x)\} = \{(x, f(x)) ; x \in X, f(x) \in Y\}$$

is called the graph of f . We shall denote the graph of f by f_G . i.e.

$$f_G = \{(x, y) \in X \times Y, y = f(x)\} = \{(x, f(x)) ; x \in X, f(x) \in Y\}.$$

In the case X and Y are normed spaces. Then $X \times Y$ is normed spaces. We will now generalize the above notion of graph.

Closed linear Functions

(54)

Def. let X and Y be a normed spaces and let D be a subspace of X . The linear function $f: D \rightarrow Y$ is called closed if every sequence $\{x_n\}$ in D such that $x_n \rightarrow x \in X$ and $f(x_n) \rightarrow y$, then $x \in D$ and $y = f(x)$.

Theorem: let X and Y be a normed spaces and let D be a subspace of X . The linear function $f: D \rightarrow Y$ is closed iff its graph f_G is closed subspace.

Proof: suppose that $f: D \rightarrow Y$ is closed and to prove that f_G is closed subspace.

let (x, y) be any limit point of f_G , i.e. $(x, y) \in \overline{f_G}$.

Then there is sequence of points in f_G , $(x_n, f(x_n))$ where $x_n \in D$ such that $(x_n, f(x_n)) \rightarrow (x, y)$

$$\Rightarrow (x_n, f(x_n)) - (x, y) \rightarrow 0 \Rightarrow \|x_n - x\| \rightarrow 0 \text{ and } \|f(x_n) - y\| \rightarrow 0$$

$$\Rightarrow x_n \rightarrow x \text{ and } f(x_n) \rightarrow y$$

Since $f: D \rightarrow Y$ is closed, then $x \in D$ and $f(x) = y$

$$\Rightarrow (x, y) \in f_G \Rightarrow f_G \text{ is closed.}$$

Conversely, let the graph f_G is closed. To prove that the linear function $f: D \rightarrow Y$ is closed.

let $\{x_n\}$ be a sequence in D such that $x_n \rightarrow x \in X$

$$\text{and } f(x_n) \rightarrow y \Rightarrow (x_n, f(x_n)) \rightarrow (x, y)$$

$$\Rightarrow (x, y) \in \overline{f_G} \text{ since } \overline{f_G} \text{ is closed } \Rightarrow \overline{f_G} = f_G$$

المحاضرة السابعة

$\Rightarrow (x, y) \in f_G \Rightarrow x \in D$ and $y = f(x)$.

(55)

\Rightarrow linear function $f: D \rightarrow Y$ is closed.

Theorem: let X and Y be a Banach spaces. If

$f: X \rightarrow Y$ is a linear function, then f is conts.

iff its graph is closed.

Proof: suppose that f is conts. To prove f_G is closed

let (x, y) be any limit point of f_G i.e. $(x, y) \in \bar{f}_G$

$\Rightarrow \exists (x_n, f(x_n)) \in f_G \ni (x_n, f(x_n)) \rightarrow (x, y)$

$\Rightarrow (x_n, f(x_n)) - (x, y) \rightarrow 0 \Rightarrow \|(x_n, f(x_n)) - (x, y)\| \rightarrow 0$

$\Rightarrow \|x_n - x, f(x_n) - y\| \rightarrow 0$

$\Rightarrow \|x_n - x\| \rightarrow 0 + \|f(x_n) - y\| \rightarrow 0$

$\Rightarrow x_n \rightarrow x$ and $f(x_n) \rightarrow y$

Since f is conts. and $x_n \rightarrow x \Rightarrow f(x_n) \rightarrow f(x)$

$\Rightarrow f(x) = y \Rightarrow (x, y) = (x, f(x)) \in f_G$.

$\Rightarrow f_G$ is closed.

Conversely let f_G be closed. To show that

f is conts. ?

"Boundedness"

(56)

Def. Let A be a subset of a topological linear space X over F . We say that A is a bounded if for any neighborhood V of 0 in X , there is a real number $\lambda > 0$ such that $A \subset \lambda V$, and we say that X is locally bounded if there is a bounded neighborhood V of 0 in X .

Note: Let A be a subset of a normed space X . A function $f: A \rightarrow X$ is compact if $f(B)$ is a compact subset of X whenever B is bounded subset of A .

Theorem: Let X be a topological linear space over a field F and $A, B \subseteq X$. Then

1. If A is finite, then A is bounded.
2. If B is bounded and $A \subseteq B$, then A is bounded.
3. If A and B are bounded sets, then $A \cap B$, $A \cup B$, $A + B$ are bounded sets.
4. If A is bounded, αA is bounded for all $\alpha \in F$.
5. If A is bounded, \bar{A} is bounded.

Proof: (1) Since A is finite set, then $A = \{a_1, \dots, a_n\}$. Let V be a neighbor. of 0 in X . Then there exists a balanced neighbor. W of 0 in $X \ni W \subset V$.

Since every neighbor. is absorbing set, then W is absorbing set, so for all $x \in X \exists \lambda > 0 \ni \lambda x \in W$.

Since $A \subset X \Rightarrow a_i \in X \forall i=1,2,\dots,n$
 $\Rightarrow \exists \lambda_i > 0 \ni \lambda_i a_i \in W \forall i=1,2,\dots,n$

(57)

Take $\lambda = \max\{\lambda_1, \dots, \lambda_n\}$

Since W balanced set $\Rightarrow \bigcup_{i=1}^n \lambda_i W = \lambda W$

$\Rightarrow A \subseteq \bigcup_{i=1}^n \lambda_i W \Rightarrow A \subset \lambda W \Rightarrow A \subset \lambda V$

$\Rightarrow A$ is bounded.

2. Let V be a neighbor. of 0 in X .

Since B is bounded set, then $\exists \lambda > 0 \ni B \subset \lambda V$

Since $A \subseteq B \Rightarrow A \subset \lambda V \Rightarrow A$ is bounded.

3. (i) since $A \cap B \subset A$ and A is bounded
 $\Rightarrow A \cap B$ is bounded.

(ii) Let V be a neighbor. of 0 in X , there is balanced neighbor. W of 0 in X such that $W \subset V$.

Since A and B are bounded, then there exists

$\lambda_1, \lambda_2 > 0 \ni A \subset \lambda_1 W$ and $B \subset \lambda_2 W$.

Take $\lambda = \max\{\lambda_1, \lambda_2\}$, since $W \subset V$

$\Rightarrow \lambda W \subset \lambda V \Rightarrow A \cup B \subset \lambda V \Rightarrow A \cup B$ bounded

(iii) Let V be a neighbor. of 0 in X , there is a symmetric neigh. W of 0 in $X \ni W + W \subset V \Rightarrow$ there is a balanced neighbor. U of 0 in X such that $U \subset W$.

Since A, B are bounded, then there exist, $\lambda_1, \lambda_2 > 0 \ni$
 $A \subset \lambda_1 U$ and $B \subset \lambda_2 U$.

Take $\lambda = \max\{\lambda_1, \lambda_2\}$

Since $U \subset W \Rightarrow \lambda(U+U) \subset \lambda(W+W) \subset \lambda V$

$\Rightarrow A+B \subset \lambda V \Rightarrow A+B$ is bounded.

المحاضرة الثامنة

4. If $\alpha = 0$, then $\alpha A = \{0\} \Rightarrow \alpha A$ is bounded. (58)

If $\alpha \neq 0$, let V be a neighbor. of 0 in X , there is a balanced neighbor. W of 0 in $X \ni W \subset V$.

Since A is bounded, there is $\lambda > 0 \ni A \subset \lambda W$.

Take $r = \lambda |\alpha| \Rightarrow r > 0$

since W is balanced and $d \leq |\alpha| \Rightarrow \alpha W \subset |\alpha| W$
 $\Rightarrow \lambda \alpha W \subset \lambda |\alpha| W$.

Since $A \subset \lambda W \Rightarrow \alpha A \subset \lambda \alpha W \subset \lambda |\alpha| W = r W$

Since $W \subset V \Rightarrow r W \subset r V \Rightarrow \alpha A \subset r V$

$\Rightarrow \alpha A$ is bounded.

5. Let V be a neighbor. of 0 in X , there is a neighbor. W of 0 in $X \ni \bar{W} \subset V$.

since A is bounded, there is $\lambda > 0$ such that

$$A \subset \lambda W \Rightarrow \bar{A} \subset \overline{\lambda W} = \lambda \bar{W}$$

we have $\bar{W} \subset V \Rightarrow \lambda \bar{W} \subset \lambda V \Rightarrow \bar{A} \subset \lambda V$.

$\Rightarrow \bar{A}$ is bounded.

Def. let A be a subset of a topological ^{linear} space X over F . We say that A is a Totally bounded if for any neighbor. V of 0 in X , there exists a finite subset B of X such that $A \subset B + V$.

Theorem: If A is a totally bounded of a topological linear space over F , then for any neighbor. V of 0 in X , there exists a finite subset A_0 of $A \ni A \subset V + A_0$.

Theorem: let X be topological linear space (59) over a field F and $A, B \subseteq X$. Then

1. If A is finite, then A is totally bounded.
2. If A is totally bounded, then A is bounded.
3. If B is a totally bounded and $A \subseteq B$, then A is totally bounded.
4. If A, B are totally bounded sets, then $A \cap B, A \cup B, A + B$ are totally bounded sets.

Proof: (1) since $A \subseteq A + V$, for every neighbor. V of 0 in X . $\Rightarrow A$ is totally bounded.

(2) let V be a neighbor. of 0 in X , there is balanced neighbor. W of 0 in X such that $W \subseteq V$.

Since W is balanced neighbor. of 0 in X and A is totally bounded set, there exists a finite subset B of X such that $A \subseteq B + W$.

Since B is finite $\Rightarrow B$ is bounded, $\Rightarrow \exists \alpha > 0 \Rightarrow B \subseteq \alpha W$.

Also W balanced $\Rightarrow \alpha W + W \subseteq (\alpha + 1)W$.

Take $\lambda = \alpha + 1 \Rightarrow A \subseteq \lambda W \subseteq \lambda V$
 $\Rightarrow A$ is bounded set.

3. let V be neighbor. of 0 in X .

Since B is a totally bounded, \exists finite subset D of X
 $\Rightarrow B \subseteq D + V$.

we have $A \subseteq B \Rightarrow A \subseteq D + V \Rightarrow A$ is totally bounded.

4. (i) we have $A \cap B \subseteq A$ and A is totally bounded
 $\Rightarrow A \cap B$ is totally bounded by using (3).

(ii) let V be a neighbor. of 0 in X . (60)

we have A and B are totally bounded, \exists finite subsets $D_1, D_2 \ni A \subset D_1 + V$ and $B \subset D_2 + V$.

Take $D = D_1 \cup D_2 \Rightarrow D$ is finite subsets and

$A \cup B \subset D + V \Rightarrow A \cup B$ is totally bounded.

(iii) let V be a neighbor. of 0 in X , \exists symmetric neighb.

W of 0 in $X \ni W + W \subset V$

we have A and B are totally bounded, then there are finite subset D_1, D_2 such that

$A \subset D_1 + W$ and $B \subset D_2 + W$.

Take $D = D_1 \cup D_2 \Rightarrow D$ is finite subset,

$A + B \subset D + W + W \subset D + V \Rightarrow A + B$ is totally bounded.

Def. let X be topological linear space over F .

1. A sequence $\{x_n\}$ in X is said to be Converge to the point $x \in X$ if for every neighbor. V of 0 in X , there exist, $K \in \mathbb{Z}^+ \ni x_n \in x + V \quad \forall n \geq K$ and we write $x_n \rightarrow x$ or $\lim_{n \rightarrow \infty} x_n = x$

2. A sequence $\{x_n\}$ in X is said to Cauchy sequence if \forall neighbor. V of 0 in X , $\exists K \in \mathbb{Z}^+ \ni x_n - x_m \in V$ for all $n, m \geq K$.