

الكورس ٢ - زمر

Quotient Group

المحاضرة ١

Here is a fundamental construction of a new group from a given group.

Theorem (1): Let G/K denotes the family of all the cosets of a subgroup K of G . If K is a normal subgroup, then:

$$aK bK = abK$$

for all $a, b \in G$, and G/K is a group under this operation

Definition (2): The group G/K is called the **quotient group**; when G is finite, its order G/K is the index $[G:K]$ (presumably, this is the reason quotient groups are so called).

We can now prove the converse of Proposition 2.91(ii).

Proposition (3): Every normal subgroup K of a group G is the kernel of some homomorphism.

Proof:

Define the natural map $\pi: G \rightarrow G/K$ by $\pi(a) = aK$. With this notation, the formula $aK bK = abK$ can be rewritten as $\pi(a)\pi(b) = \pi(ab)$; thus, π is a (surjective) homomorphism. Since K is the identity element in G/K ,

$$\ker \pi = \{a \in G : \pi(a) = K\} = \{a \in G : aK = K\} = K$$

المحاضرة ٢

First Isomorphism Theorem

The following theorem shows that every homomorphism gives rise to an isomorphism, and that quotient groups are merely constructions of homomorphic images.

Theorem (1): (First Isomorphism Theorem)

If $f: G \rightarrow H$ is a homomorphism, then:

$$G / \ker f \cong \text{im } f$$

Where $\text{im } f = f(H)$. In more detail, if we put $\ker f = K$, then the function $\phi: G/K \rightarrow f(H)$ is given by: $\phi: aK \mapsto f(a)$ for each $a \in G$, is an isomorphism.

Proof:

It is clear that $\ker f$ is a normal subgroup of G , and we can easily show that ϕ is well-defined. Let us now see that ϕ is a homomorphism. Since f is a homomorphism and $\phi(aK) = f(a)$,
 $\phi(aK bK) = \phi(abK) = f(ab) = f(a)f(b) = \phi(aK)\phi(bK)$.

Also ϕ is surjective and injective. Therefore, $\phi: G/K \rightarrow \text{im } f$ is an isomorphism.

Remark (2):

1. Here is a minor application of the first isomorphism theorem. For any group G , the identity function $f: G \rightarrow G$ is a surjective homomorphism with $\ker f = 1$.

المحاضرة ٣

Proposition (3):

1. If H and K are subgroups of a group G , and if one of them is a normal subgroup, then HK is a subgroup of G . Moreover, $HK = KH$.
2. If both H and K are normal subgroups, then HK is a normal subgroup.

Proof:

1. Assume first that K is normal in G . We claim that $HK = KH$. If $hk \in HK$, then:
 $hk = hkh^{-1}h = k1h \in KH$

where $k_1 = hkh^{-1}$, then $k_1 \in K$, because K is normal subgroup ₂

Hence, $HK = KH$. For the reverse inclusion, write $kh = hh^{-1}kh = hk^2 \in HK$, where $k^2 = h^{-1}kh$. (Note that the same argument shows that $HK = KH$ if H is normal subgroup of G .) ³

المحاضرة ٤

¹

Third Isomorphism Theorem

Lecture 11

In the following lecture we study the third important theorem of fundamental isomorphism theorem.

Theorem (1): (Third Isomorphism Theorem)

If H and K are normal subgroups of a group G with $K \leq H$, then H/K is normal in G/K and $(G/K)/(H/K) \cong G/H$. ²

Proof:

Define $f: G/K \rightarrow G/H$ by $f(aK) = aH$. Note that f is a (well-defined) function, for if $aK = bK$, then $a^{-1}b \in K$. But $K \subseteq H$, thus $a^{-1}b \in H$, and so $aH = bH$, and we are done. It is easy to see that f is an epimorphism.

Now $\ker f = H/K$. Also clearly H/K is a normal subgroup of G/K . Since f is monomorphism, so by the first isomorphism theorem we have: $(G/K)/(H/K) \cong G/H$

The third isomorphism theorem is easy to remember: the K 's in the fraction $(G/K)/(H/K)$ can be canceled. One can better appreciate the first isomorphism theorem after having proved the third one. The quotient group $(G/K)/(H/K)$ consists of cosets (of H/K) whose representatives are themselves cosets (of G/K). 3

Here is another construction of a new group from two given groups.

Definition (2): If H and K are groups, then their **direct product**, denoted by $H \times K$, is the set of all ordered pairs (h, k) equipped with the following operation:

$$(h, k)(h_1, k_1) = (hh_1, kk_1)$$

It is routine to check that $H \times K$ is a group [the identity element is (e, e_1) and $(h, k)^{-1} = (h^{-1}, k^{-1})$].

Remark (3): let G and h be groups. Then $H \times K$ is abelian if and only if both H and K are abelian.

We end the tenth lecture by the following example.

Example: $Z \times 2Z$ is the direct product between $(Z, +)$ and $(2Z, +)$ groups.

The identity element is $(0, 0)$, and the inverse element of (a, b) is $(-a, -b)$.

We now show that HK is a subgroup. Since $e \in H$ and $e \in K$, we have $e = e \cdot e \in HK$. If $hk \in HK$, then $(hk)^{-1} = k^{-1}h^{-1} \in KH = HK$. If $hk, h_1k_1 \in HK$, then $h_1^{-1}kh_1 = ke \in K$ and $Hkh_1k_1 = hh_1(h_1^{-1}kh_1)k_1 = (hh_1)(kek_1) \in HK$. Therefore, HK is a subgroup of G .

2. If $g \in G$, then:

$$ghkg^{-1} = (ghg^{-1})(gkg^{-1}) \in HK$$

Therefore, HK is normal in G . $ghkg^{-1} =$

$$(ghg^{-1})(gkg^{-1}) \in HK. \text{ Therefore, } HK \text{ is normal in } G.$$

Definition (1): If H and K are subgroups of a finite group G , then then the **Product Formula** is:

$$|HK||H \cap K| = |H||K|$$

المحاضرة ٥

One can shorten the list of items needed to verify that a subset is, in fact, a subgroup.

Proposition (4): A subset H of a group G is a subgroup if and only if H is nonempty and, whenever $x, y \in H$, then $xy^{-1} \in H$.

Proof: If H is a subgroup, then it is nonempty, for $1 \in H$. If $x, y \in H$, then $y^{-1} \in H$, by part (iii) of the definition, and so $xy^{-1} \in H$, by part (ii).

Conversely, assume that H is a subset satisfying the new condition. Since

H is nonempty, it contains some element, say, h .

Taking $x = h = y$, we see that $e = hh^{-1} \in H$, and so

part (i) holds. If $y \in H$, then set $x = e$ (which we can now do because e

$\in H$), giving $y^{-1} = ey^{-1} \in H$, and so part (iii) holds. Finally, we know that $(y^{-1})^{-1} = y$, by. Hence, if $x, y \in H$, then $y^{-1} \in H$ and so $xy = x(y^{-1})^{-1} \in H$. Therefore, H is a subgroup of G . Since every subgroup contains e , one may replace the hypothesis “ H is nonempty” in Proposition by “ $e \in H$ ”.

Note that if the operation in G is addition, then the condition in the proposition is that H is a nonempty subset of G such that $x, y \in H$ implies $x - y \in H$.

Proposition (5): Let G be a finite group, and $a \in G$. Then the order of a , is the number of elements in $\langle a \rangle$.

Definition (6): If G is a finite group, then the number of elements in G , denoted by $|G|$, is called the **order of G** .

Definition (7): If X is a subset of a group G , such that X generates G , then G is called **finitely generated**, and G generated by X .

In particular; If $G = \langle \{a\} \rangle$, then G is generated by the subset $X = \{a\}$.

Definition (8):

A group G is called **cyclic** if $G = \langle a \rangle$; that is G can be generated by only one element say a , and this element is called a generator of G .

المحاضرة ٦

Lecture 13

If H and K are subgroups of a group G with H is normal in G , then HK is a subgroup of G and $H \cap K$ is normal in K .

Moreover:

$$K / (H \cap K) \cong HK / H$$

Proof:

We begin by showing first that HK/H makes sense, and then describing its elements. Since H is normal subgroup of G , then HK is a subgroup of G . Normality of H in HK follows :

from a more general fact: if $H \trianglelefteq S \trianglelefteq G$ and if H is normal in G , then H is normal in S .

We can easily show that each coset $xH \trianglelefteq HK/H$ has the form kH for some $k \in K$. It follows that the function $f: K \rightarrow HK/H$, given by $f(k) = kH$, is surjective. Moreover, f is a homomorphism, for it is the restriction of the natural map $\pi: G \rightarrow G/H$. Since $\ker \pi = H$, it follows that $\ker f = H \cap K$ and so $H \cap K$ is a normal subgroup of K . The first isomorphism theorem gives:

$$K/(H \cap K) \cong HK/H$$

Remark (3):

The second isomorphism theorem gives the product formula in the special case when one of the subgroups is normal: if $K/(H \cap K) \cong HK/H$, then: $|K/(H \cap K)| = |HK/H|$, and so $|HK| = |H| |K/(H \cap K)| = |H| |K|$.

المحاضرة ٧

Definition (6): If $f: G \rightarrow H$ is a homomorphism, define

$$\mathbf{kernel\ } f = \{x \in G : f(x) = e\}$$

$$\mathbf{and\ image\ } f = \{h \in H : h = f(x) \text{ for some } x \in G\}.$$

We usually abbreviate kernel f to $\ker f$ and image f to $\text{im } f$

So that if $f: G \rightarrow H$ is a homomorphism and B is a subgroup of H then $f^{-1}(B)$ is a subgroup of G containing $\ker f$.

Note: Kernel comes from the German word meaning “grain” or “seed” (corn comes from the same word).

Its usage here indicates an important ingredient of a homomorphism, we give it without proof.

Proposition: Let $f: G \rightarrow H$ be a homomorphism.
(i) $\ker f$ is a subgroup of G and $\operatorname{im} f$ is a subgroup of H .

(ii) If $x \in \ker f$ and if $a \in G$, then $axa^{-1} \in \ker f$.

(iii) f is an injection if and only if $\ker f = \{e\}$.

Normal Subgroups

Definition (1): A subgroup K of a group G is called **normal**, if for each $k \in K$ and $g \in G$ imply $gkg^{-1} \in K$. that is $gKg^{-1} \subseteq K$ for every $g \in G$.

Definition (2):

Define the **center of a group G** , denoted by $Z(G)$, to be

$Z(G) = \{z \in G: zg = gz \text{ for all } g \in G\}$;

that is, $Z(G)$ consists of all elements commuting with every element in G . (Note that the equation zg

123

gz can be rewritten as $z = gzg^{-1}$, so that no other elements in G are conjugate to z .

Remark (3):

Let us show that $Z(G)$ is a subgroup of G . We can easily show that $Z(G)$ is subgroup of G . It is clear that $Z(G) \neq \emptyset$ since $1 \in Z(G)$, for 1 commutes with everything. Now, If $y, z \in Z(G)$, then $yg = gy$ and $zg = gz$ for all $g \in G$. Therefore, $(yz)g = y(zg) = y(gz) = (yg)z = g(yz)$, so that yz commutes with everything, hence $yz \in Z(G)$. Finally, if $z \in Z(G)$, then $zg = gz$ for all $g \in G$; in particular, $zg^{-1} = g^{-1}z$. Therefore,

$$gz^{-1} = (zg^{-1})^{-1} = (g^{-1}z)^{-1} = z^{-1}g$$

(we are using $(ab)^{-1} = b^{-1}a^{-1}$ and $(a^{-1})^{-1} = a$).

So that $Z(G)$ is subgroup of G .

Clearly the center $Z(G)$ is a normal subgroup; since if $z \in Z(G)$ and $g \in G$, then $gzg^{-1} = zgg^{-1} = z \in Z(G)$

A group G is abelian if and only if $Z(G) = G$. At the other extreme are groups G for which $Z(G) = \{1\}$; such groups are called centerless. For example, it is easy to see that $Z(S_3) = \{1\}$; indeed, all large symmetric groups are centerless.

$\{1\}$. By the first isomorphism theorem, we have $G/\{1\} \cong G$

2. Given any homomorphism $f:G \rightarrow H$, one should immediately ask for its kernel and its image; the first isomorphism theorem will then provide an isomorphism

$G/\ker f \cong \text{im } f$. Since there is no significant difference between

isomorphic groups, the first isomorphism theorem also says that there is no significant difference between quotient groups and homomorphic images.