



كلية : التربية للعلوم الصرفة

القسم او الفرع : الرياضيات

المرحلة: الثالثة

أستاذ المادة : م.د. ناديه علي ناظم

اسم المادة باللغة العربية : التحليل الرياضي

اسم المادة باللغة الإنجليزية : Mathematical Analysis

اسم الحاضرة السادسة باللغة العربية: المتتابعات في الفضاء المترى

اسم المحاضرة السادسة باللغة الإنجليزية: sequences in metric space:

Mathematical Analysis

Dr. Nadia Ali

Teaching at the University of Anbar
College of Education for Pure Sciences
Department of Mathematics

2022 – 2023

Chapter Three

Sequences in Metric Space

Definition 3.1: Let S be any set a function f whose domain is the set N and the range is S is

Called a sequence in S .

i.e. $f: N \rightarrow S$, where $\forall n \in N, \exists x_n \in S$ s.t $f(n) = x_n$

1. $\left\langle \frac{1}{5n} \right\rangle = \frac{1}{5}, \frac{1}{10}, \frac{1}{15}, \dots$
2. $\left\langle \frac{1}{n+1} \right\rangle = \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$
3. $\left\langle 4 \right\rangle = 4, 4, 4, \dots$
4. $\left\langle n - 3 \right\rangle = -2, -1, 0, 1, \dots$

Definition 3.2: Let (X, d) be a metric space and $\langle X_n \rangle$ be seq. in X , then $\langle X_n \rangle$ is said to be converges to appoint in X , if $\forall \epsilon > 0, \exists k \in N$ s.t $d(X_n, x) < \epsilon, \forall n > k$. We write $X_n \rightarrow x$ or $\lim_{n \rightarrow \infty} X_n = x$, x is called

A Limit point of $\langle X_n \rangle$.

If $\forall n > K$, does not Converge, them $\langle X_n \rangle$ is called divergent Sequence.
Not that: K depend on ϵ only.

التغير الهندسي للتعریف التقارب

$$(X_n \rightarrow x)$$

يعني الكرة التي مركزها x ونصف قطرها ϵ تمتلك عدد غير منتهي من حدود او نقاط المتتابعة
لأنه X_n

$\forall \epsilon > 0, \exists k \in N$ s.t $d(X_n, x) < \epsilon, \forall n > k \Rightarrow X_n \in B(x, \epsilon)$.

Ex: Let $\langle X_n \rangle = \langle 1 \rangle$ constant seq. show that $\lim_{n \rightarrow \infty} X_n = 1$

$\langle 1 \rangle$ converges to 1 since $\forall \epsilon > 0, \exists k \in N$

s.t $d(X_n, x) = |1 - 1| = 0 < \epsilon, \forall n > k$

Ex: Let $\langle X_n \rangle$ be a seq. defined by $X_n = \begin{cases} n & \text{if } n \leq 50 \\ 3 & \text{if } n \geq 50 \end{cases}$. show that

$$\lim_{n \rightarrow \infty} X_n = 3$$

Solution:

$$\langle X_n \rangle \geq 1, 2, 3, \dots, 50, 3, 3, 3, \dots$$

$$\forall \epsilon > 0, \exists k = 50 \text{ s.t } d(X_k, x) = |3 - 3| = 0 < \epsilon$$

Example 3.1: Show that $\lim_{n \rightarrow \infty} X_n = 2$, where $\langle X_n \rangle = \langle \frac{2n-3}{n+1} \rangle$

Solution:

$$\forall \epsilon > 0, \text{ to find } K \in \mathbb{N} \text{ s.t } d(X_n, x) < \epsilon, \forall n > K ?$$

$$\begin{aligned} d(X_n, x) &= \left| \frac{2n-3}{n+1} - 2 \right| = \left| \frac{2n-3-2(n+1)}{n+1} \right| \\ &= \left| \frac{2n-3-2n-2}{n+1} \right| = \left| \frac{-5}{n+1} \right| = \frac{5}{n+1} \end{aligned}$$

$\forall \epsilon > 0$, by Arch. Property $\rightarrow \exists K \in \mathbb{N} \exists$

$$\forall k > 5 \rightarrow \frac{5}{\epsilon} < k.$$

$$\forall n > K \rightarrow n+1 > k+1 \text{ and } k+1 > k, k > \frac{5}{\epsilon}$$

$$\Rightarrow n+1 > k+1 > k > \frac{5}{\epsilon}$$

$$\frac{1}{n+1} < \frac{\epsilon}{5}, \forall n > k$$

Exercise 3.1:

1. Let $\langle X_n \rangle = \langle \frac{2}{\sqrt{n}} \rangle$, show that $\lim_{n \rightarrow \infty} X_n = 0$

2. Let $\langle X_n \rangle = \langle \frac{5n-4}{2-3n} \rangle$, show that $\lim_{n \rightarrow \infty} X_n = -\frac{5}{3}$

3. Let $\langle X_n \rangle = \langle \frac{2-7n}{1-5n} \rangle$, show that $\lim_{n \rightarrow \infty} X_n = \frac{7}{5}$

Show that the following sequence are divergent

1. $\langle X_n \rangle = \langle \sqrt{n} \rangle$
2. $\langle X_n \rangle = \langle (-1)^n \rangle$
3. $\langle X_n \rangle = 3^n$
4. $\langle X_n \rangle = \langle \frac{n^2}{2n-1} \rangle$

Theorem 3.1: If $\langle X_n \rangle$ is convergent sequence in (X, d) , then $\langle X_n \rangle$ has a unique limit point.

Proof:

Suppose $\langle X_n \rangle$ has two limit points x and y with $x \neq y$ and $d(x, y) = \epsilon$

Since $X_n \rightarrow y \Rightarrow \forall \epsilon > 0, \exists k_2 \in N \text{ s.t. } d(x, y) < \frac{\epsilon}{2}$

Let $k = \max\{k_1, k_2\}$

Since $d(x, y) \leq d(x, x_n) + d(x_n, y) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$

$\Rightarrow d(x, y) < \epsilon, \forall \epsilon > 0$

This true only when $d(x, y) = 0 \Rightarrow x = y \rightarrow C!$

$\therefore \langle X_n \rangle$ has a unique limit point.

Definition 3.3: A seq. $\langle X_n \rangle$ is called bounded if the set $\{X_n : n \in N\}$ is bounded

i.e. $\langle x_n \rangle$ is bounded if $\exists M > 0 \text{ s.t. } d(x_n, x_m) \leq M, \forall n, \forall m$