Derivatives and Integrals

Differentiation:

We previous studies, we defined the slope of the curve at a point as the limit of secant slopes, and this limit called derivative measure rate of which a function changes, and it is one of the most important ideas in calculus.

Derivatives are used to calculate velocity and acceleration, to estimate the rate of spread of a distance, to set of production to find the best dimensions of figures (cylindrical, circle ...etc), and for many applications.

In this chapter, we develop techniques to calculate derivatives easily and learn how to use derivatives to approximate complicated functions.

The derivative of the function f(x) respect to the variable x is the function $\hat{f}(x)$ whose value at x is

$$\dot{f}(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

We say that f is differentiable (has derivative) at x if \hat{f} exists at every point in the domain of f.

The slope of tangent line:

The slope of the curve y = f(x) at the point $p(x_{\circ}, f(x_{\circ}))$ is the number

$$m = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

The tangent line to the curve at p is the line through p with this slope.

In the previous studies the slope $m = \frac{\Delta y}{\Delta x}$

$$\therefore m = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{f(x_\circ + h) - f(x_\circ)}{(x_\circ + h) - x_\circ} = \frac{f(x_\circ + h) - f(x_\circ)}{h}$$

We say that f is differentiable (has derivative) at x if f exists at every point in the domain of f, then we call f differentiable.

But if *f* is not differentiable at every point in the domain of *f*, for example sqort or rotational functions, then we write z = x + h, then h = z - x and *h* approaches 0 if *z* approaches *x*.

Therefore, an equivalent definition of the derivative is as follows

$$\dot{f}(x) = \lim_{z \to x} \frac{f(z) - f(x)}{z - x}$$

Example1/ Derivative of square root function:

a- find the derivative of $y = \sqrt{x}$ for x > 0

b- Find the tangent line equation to the curve $y = \sqrt{x}$ at x=4 Solution:

a-

$$\hat{f}(x) = \lim_{z \to x} \frac{f(z) - f(x)}{z - x} = \lim_{z \to x} \frac{\sqrt{z} - \sqrt{x}}{z - x} = \lim_{z \to x} \frac{(\sqrt{z} - \sqrt{x})}{(\sqrt{z} - \sqrt{x})(\sqrt{z} + \sqrt{x})} \\
= \lim_{z \to x} \frac{1}{\sqrt{z} + \sqrt{x}} = \frac{1}{\sqrt{x} + \sqrt{x}} = \frac{1}{2\sqrt{x}}$$

b-

The slope of the curve at x = 4 is $f(x) = \frac{1}{2\sqrt{x}}$, then $m = f(4) = \frac{1}{2\sqrt{4}} = \frac{1}{4}$ The tangent line through the point = 4, and $y = f(x) = \sqrt{x}$ Then $= f(4) = \sqrt{4} = 2$, then the point p(x, y) = p(4, 2)The equation of tangent line through the point p(4, 2) and the slope $(m = \frac{1}{4})$ equal $y = m(x - x_1) + y_1 \xrightarrow{yields} y = \frac{1}{4}(x - 4) + 2 = \frac{1}{4}x + 1$ In another method:

$$f(x) = \sqrt{x} = x^{1/2}$$
$$m = f(x) = \frac{1}{2}x^{\frac{1}{2}-1} = \frac{1}{2}x^{-1} = \frac{1}{2\sqrt{x}}$$

: the slope at
$$x = 4$$
 is $m = \hat{f}(4) = \frac{1}{2\sqrt{4}} = \frac{1}{4}$

$$y_1 = f(x_1) = f(4) = \sqrt{4} = 2$$

Then point $p(x_1, y_1) = p(4, 2)$

The equation of tangent line through the point p(4,2) and the slope $(m = \frac{1}{4})$ equal $y = m(x - x_1) + y_1 \xrightarrow{yields} y = \frac{1}{4}(x - 4) + 2 = \frac{1}{4}x + 1$

Differentiable on an interval; one sided derivatives:

A function y = f(x) is differentiable on an open interval (finite or infinite) if it has a derivative at each point of the interval.

 $\lim_{h \to 0^+} \frac{f(a+h) - f(a)}{h}$ Right-hand derivative at a

 $\lim_{h \to 0^{-}} \frac{f(b+h) - f(b)}{h}$ Left-hand derivative at b

Example 2/Function y = |x| is not differentiable at the origin, show that the function is differentiable on $(-\infty, 0)$ and $(0, \infty)$ but has no derivative at x = 0.

Solution:

To the right of the origin $\frac{d}{dx}(|x|) = \frac{d}{dx}(x) = 1$

To the left of the origin $\frac{d}{dx}(|x|) = \frac{d}{dx}(-x) = -1$

 \therefore derivative on the left \neq derivative on the right

Example3/ Show that the function $f(x) = \sqrt{x}$ is not differentiable at x = 0

Solution: $\frac{d}{dx}\sqrt{x} = \frac{1}{2}x^{\frac{1}{2}-1} = \frac{1}{2\sqrt{x}}$

We apply the definition to examine if the derivative exists at x = 0

$$m = \lim_{h \to 0} \frac{f(x_{\circ} + h) - f(x_{\circ})}{h} = \lim_{h \to 0} \frac{\sqrt{(x+h)} - \sqrt{x}}{h} = \lim_{h \to 0} \frac{\sqrt{h}}{h}$$
$$= \lim_{h \to 0} \frac{\sqrt{h}}{\sqrt{h}\sqrt{h}} = \lim_{h \to 0} \frac{1}{\sqrt{h}} = \infty$$

 \therefore the function is not have derivative at x = 0

Example4/ Find derivative functions and values using the definition, calculate the derivatives of the functions, then find the values of the derivatives as specified.

1-
$$f(x) = 4 - x^2$$
, $f'(-3)$, $f'(0)$, $f'(1)$

Solution:

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \to 0} \frac{4 - (x+h)^2 - (4 - x^2)}{h}$$
$$= \lim_{h \to 0} \frac{4 - x^2 - 2hx - h^2 - 4 + x^2}{h} = \lim_{h \to 0} \frac{-h(2x+h)}{h}$$
$$= -\lim_{h \to 0} 2x + h = -2x$$
$$2 - f(x) = (x-1)^2 + 1, \quad f'(-1), f'(0), f'(2)$$

Solution:

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

=
$$\lim_{h \to 0} \frac{\left[\left((x+h) - 1\right)^2 + 1\right] - \left[(x-1)^2 + 1\right]}{h}$$

=
$$\lim_{h \to 0} \frac{\left[(x+h)^2 - 2(x+h) + 1 + 1\right] - \left[x^2 - 2x + 1 + 1\right]}{h}$$

=
$$\lim_{h \to 0} \frac{\left[x^2 + 2xh + h^2 - 2x - 2h + 2 - x^2 + 2x - 2\right]}{h}$$

=
$$\lim_{h \to 0} \frac{2xh + h^2 - 2h}{h} = \lim_{h \to 0} \frac{h(2x+h-2)}{h} = 2x - 2$$

$$\therefore f'(-1) = -4, f'(0) = -2, f'(2) = 2$$

Example5/ Find the derivative of the following functions:

1-
$$y = 2x^3 \rightarrow \frac{dy}{dx} = \frac{d}{dx}(2x^3) = 6x^2$$

2-
$$r = \frac{s^3}{2} + 1 \rightarrow \frac{dr}{ds} = \frac{d}{ds} \left(\frac{s^3}{2} + 1 \right) = \frac{d}{ds} \frac{s^3}{2} + \frac{d1}{ds} = \frac{3s^3}{2} + 0$$

3- $v = t - \frac{1}{t} \rightarrow \frac{dv}{dt} = \frac{d}{dt} \left(t - \frac{1}{t} \right) = \frac{d}{dt} \left(t \right) - \frac{d}{dt} \left(\frac{1}{t} \right) = 1 - \left(-t^{-2} \right) = 1 + \frac{1}{t^2}$
4- $p = \frac{1}{\sqrt{q+1}} \rightarrow \frac{dp}{dq} = \frac{d}{dq} \left(\frac{1}{\sqrt{q+1}} \right) = \frac{d}{dq} \left(q + 1 \right)^{-1/2} = \frac{-1}{2} \left(q + 1 \right)^{\frac{-1}{2} - 1} = \frac{-1}{2} \left(q + 1 \right)^{\frac{-3}{2}} = \frac{-1}{2(q+1)^{\frac{3}{2}}}$

Example6/ Differentiable the following functions and find the slope of the tangent line at the given value of the independent variable, and find the equation of tangent line.

1-
$$f(x) = x + \frac{9}{x}$$
, at $x = -3$

Solution:

The equation of tangent line $= m(x - x_1) + y_1$, then we need slope (m) and the point $p(x_1, y_1)$

$$y_1 = f(-3) = -3 + \frac{9}{-3} = -6, \qquad \therefore p(-3, -6)$$
$$f'(x) = m = \frac{d}{dx} \left(x + \frac{9}{x} \right) = \frac{dx}{dx} + \frac{d}{dx} 9x^{-1} = 1 + 9(-1)x^{-2} = 1 - \frac{9}{x^2}$$

∴ the slope at x = -3 → $m = 1 - \frac{9}{(-3)^2} = 1 - \frac{9}{9} = 1 - 1 = 0$ ∴ equation of tangent line $y = 0(x - x_1) + (-6) = -6$ ∴ tangent line is horizontal line, where y = -6

2-
$$k(x) = \frac{1}{2+x^2}$$
 at $x = 2$
 $\therefore y_1 = k(2) = \frac{1}{2+x^2} = \frac{1}{6}$ $\therefore p(2, \frac{1}{6})$

The slope $(m) = k'(x) = \frac{d}{dx}(2+x^2)^{-1} = -1(2+x^2)^{-2} \cdot 2x$ $= \frac{-2x}{(2+x^2)^2}$ \therefore the slope at $x = 2 \rightarrow m = \frac{-2(2)}{(2+2^2)^2} = \frac{-4}{36} = -\frac{1}{9}$ \therefore equation of tangent line at $p\left(2,\frac{1}{6}\right)$ and $m = \frac{-1}{9}$ is $y = m(x-x_1) + y_1 \rightarrow y = \frac{-1}{9}(x-2) + \frac{1}{6} = \frac{-1}{9}x + \frac{2}{9} + \frac{1}{6}$ $= \frac{-1}{9}x + \frac{21}{54}$

3- $s = t^3 - t^2$ at t = -1 $y_1 = s(-1) = (-1)^3 - (-1)^2 = -2$ $\therefore p(-1, -2)$ $m = s' = \frac{ds}{dt} = \frac{d}{dt}(t^3 - t^2) = \frac{d}{dt}t^3 - \frac{d}{dt}t^2 = 3t^2 - 2t$ the slope at t = -1 $\rightarrow m = 3(-1)^2 - 2(-1) = 3 + 2 = 5$ The equation of tangent line at the point p(-1, -2) and m= 5 equal y = 5(x - (-1)) + (-2) = 5x + 3

Example7/ Find equation of the tangent line at the indicated point for the following function:

Solution:

$$y = 8(x-2)^{\frac{-1}{2}} \to y' = \frac{dy}{dx} = \frac{d}{dx}8(x-2)^{\frac{-1}{2}} = 8\frac{d}{dx}(x-2)^{\frac{-1}{2}}$$
$$y' = m = 8\left(\frac{-1}{2}\right)(x-2)^{\frac{-3}{2}} = \frac{-4}{(x-2)^{\frac{3}{2}}}$$

The slope at x = 6 equal $m = \frac{-4}{(6-2)^{\frac{3}{2}}} = \frac{-4}{(4)^{\frac{3}{2}}} = \frac{-4}{\sqrt[2]{4^3}} = \frac{-4}{\sqrt[2]{64}} = \frac{-4}{8} = -\frac{1}{2}$

∴ equation of line $y = m(x - x_1) + y_1 \rightarrow y = -\frac{1}{2}(x - 6) + 4$ ∴ $y = -\frac{1}{2}x + 7$

2- $w = g(z) = 1 + \sqrt{4-z}$, p(z,w) = (3,2)

Solution:

$$m = w' = \frac{dg(z)}{dz} = \frac{d}{dz} 1 + \frac{d}{dz} (4-z)^{\frac{1}{2}} = 0 + \frac{1}{2} (4-z)^{\frac{-1}{2}} = \frac{1}{2(4-z)^{\frac{1}{2}}}$$

$$\therefore slope \ at \ x = 3 \ equal \quad m = \frac{1}{2(4-3)^{\frac{1}{2}}} = \frac{1}{2}$$

 \therefore equation of tangent line equal to $y = m(x - x_1) + y_1$

$$y = \frac{1}{2}(x-3) + 2 = \frac{1}{2}x - \frac{3}{2} + 2 = \frac{1}{2}x + \frac{1}{2}$$

Example8/ Find the values of the following derivatives:

1-
$$\frac{ds}{dt}$$
 at $t = -1$ if $s = 1 - 3t^2$

Solution:

$$\frac{ds}{dt} = \frac{d}{dt}(1 - 3t^2) = \frac{d}{dt}1 - 3\frac{d}{dt}t^2 = 0 - 3 \times 2t = -6t$$
$$\therefore \quad \frac{ds}{dt} \quad at \ t = -1 \quad equal \quad \frac{ds}{dt} = 6$$
$$2 - \frac{dy}{dx} \quad at \ x = \sqrt{3} \quad if \quad y = 1 - \frac{1}{x}$$

Solution:

$$\frac{dy}{dx} = \frac{d}{dx} \left(1 - \frac{1}{x} \right) = \frac{d}{dx} 1 - \frac{d}{dx} x^{-1} = 0 - (-1)x^{-2} = \frac{1}{x^2}$$
$$\frac{dy}{dx} \text{ at } x = \sqrt{3} \text{ equal } \frac{dy}{dx} = \frac{1}{3}$$
$$3 - \frac{dr}{d\theta} \text{ at } \theta = 0 \quad \text{if} \quad r = \frac{2}{\sqrt{4-\theta}}$$

Solution:

$$\frac{dr}{d\theta} = \frac{d}{d\theta} 2(4-\theta)^{\frac{-1}{2}} = 2\frac{d}{d\theta}(4-\theta)^{\frac{-1}{2}} = 2\left(-\frac{1}{2}\right)(4-\theta)^{\frac{-3}{2}}$$
$$= \frac{-1(-1)}{(4-\theta)^{\frac{3}{2}}}$$
$$\therefore \frac{dr}{d\theta} \text{ at } \theta = 0 \text{ equal } \frac{dr}{d\theta} = \frac{1}{4^{\frac{3}{2}}} = \frac{1}{\sqrt[2]{4^3}} = \frac{1}{\sqrt[2]{64}} = \frac{1}{8}$$

Differentiation Rules:

We can differentiate functions without having to apply the definition of the derivative each time.

Powers, multiples, sums and differences, derivative product rule, derivative quotient rule, power rule for negative integers, second and higher order derivative.

1- The first rule of every constant function is zero.

If *f* has the constant value f(x) = c then

$$\frac{df}{dx} = \frac{d}{dx}(c) = 0 \text{ for example } f(x) = 8, \therefore \frac{df}{dx} = \frac{d}{dx} = 0$$

2- Power rule for positive integers, if *n* is a positive integer, then

$$\frac{d}{dx}x^{n} = nx^{n-1} \text{ for example } f(x) = x^{2} :: \frac{df}{dx} = \frac{d}{dx}x^{2} = 2x$$

3- Constant multiple rule

If u is a differentiable function of x, and c is a constant, then

$$\frac{d}{dx}(cu) = c\frac{du}{dx} \text{ for example } f(x) = 3x^2 \quad \therefore \frac{df}{dx} = \frac{d}{dx}(3x^2) = 3\frac{dx^2}{dx}$$
$$= 3 \times 2x = 6x$$

4- **Derivative sum rule**, if *u* and *v* are differentiable functions of *x*, then their sum u + v is differentiable at every point where *u* and *v* are to be differentiable at each point.

$$\frac{d}{dx}(u+v) = \frac{du}{dx} + \frac{dv}{dx} \text{ for example } y = x^4 + 12x \quad \therefore \frac{dy}{dx}$$
$$= \frac{d}{dx}x^4 + \frac{d}{dx}(12x) = 4x^3 + 12$$

And for difference rule

$$\frac{d}{dx}(u-v) = \frac{du}{dx} - \frac{dv}{dx} \text{ for example } y = x^4 - 12x \quad \therefore \frac{dy}{dx} = \frac{d}{dx}x^4 - \frac{d}{dx}(12x) = 4x^3 - 12$$

5- **Derivative product rule**, if u and v are differentiable at x, then

$$\frac{d}{dx}(uv) = u\frac{dv}{dx} + v\frac{du}{dx} \text{ for example } y = \frac{1}{x}\left(x^2 + \frac{1}{x}\right) \therefore \frac{dy}{dx}$$
$$= \frac{1}{x}\frac{d}{dx}\left(x^2 + \frac{1}{x}\right) + \left(x^2 + \frac{1}{x}\right)\frac{d}{dx}\left(\frac{1}{x}\right)$$
$$= \frac{1}{x}\left[\frac{d}{dx}x^2 + \frac{d}{dx}(x^{-1})\right] + \left(x^2 + \frac{1}{x}\right)\frac{d}{dx}(x^{-1})$$
$$= \frac{1}{x}\left(2x - \frac{1}{x^2}\right) + \left(x^2 + \frac{1}{x}\right)(-x^{-2}) = \left(2 - \frac{1}{x^3}\right) - \left(1 + \frac{1}{x^3}\right)$$
$$= 2 - \frac{1}{x^3} - 1 - \frac{1}{x^3} = 1 - \frac{2}{x^3}$$

6- **Derivative quotient rule**, if *u* and *v* are differentiable at *x*, and if $v(x) \neq 0$, then the quotient $\frac{u}{v}$ is differentiable at *x*, then

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v\frac{du}{dx} - u\frac{dv}{dx}}{v^2} \quad \text{for example } \frac{d}{dx}\left[\frac{f(x)}{g(x)}\right]$$
$$= \frac{g(x)f'(x) - f(x)g'(x)}{g^2(x)}$$

Example9/ $y = \frac{t^2 - 1}{t^2 + 1}$

$$\frac{dy}{dx} = \frac{(t^2+1) \cdot 2t - (t^2-1) \cdot 2t}{(t^2+1)^2} = \frac{2t^3 + 2t - 2t^3 + 2t}{(t^2+1)^2} = \frac{4t}{(t^2+1)^2}$$

7- **Power rule for negative integers**, if *n* is a negative integer and $x \neq 0$ then $\frac{d}{dx}(x^n) = nx^{n-1}$

Example $10/\frac{d}{dx}\left(\frac{1}{x}\right) = \frac{d}{dx}(x^{-1}) = (-1)(x^{-2}) = -\frac{1}{x^2}$

8- Second and higher order derivatives

If y = f(x) is a differentiable function, then its derivative f'(x) is also a function. If f' is also differentiable, then we can differentiable f' to get a new function of x denoted by f''. f'' called second derivative, f''' third derivative,....,etc.

$$f^{//}(x) = \frac{d^2 y}{dx^2} = \frac{d}{dx} \left(\frac{dy}{dx}\right) = \frac{dy^{\prime}}{dx} = y^{\prime/}$$

Example 11/ $y = x^6$ $\therefore y' = 6x^5$, $y'' = \frac{dy'}{dx} = \frac{d}{dx}(6x^5) = 30x^4$, $y'' = \frac{d}{dx}y'' = \frac{d}{dx}30x^4 = 120x^3$

In general
$$y^n = \frac{d}{dx}y^{(n-1)} = \frac{d^ny}{dx^n}$$

Example 12/ Finding higher derivatives for the function $y = x^3 - 3x^2 + 2$

$$y' = 3x^2 - 6x$$
, $y'' = 6x - 6$, $y''' = 6$, $y''' = y^{(4)} = 0$

Example13/ Find the first and second derivative for the following functions:

1-
$$y = -x^{2} + 3 \rightarrow y' = -2x$$
, $y'' = -2$
2- $s = 5t^{3} - 3t^{5} \rightarrow s' = 15t^{2} - 15t^{4}$, $s'' = 30t - 60t^{3}$
3- $y = x^{5} + x^{3} + x \rightarrow y' = 5x^{4} + 3x^{2} + 1$, $y'' = 20x^{3} + 6x$

Example14/ How a circles area changes with its diameter.

Solution: the area of a circle is related to its diameter by the equation $A = \frac{\pi}{4}D^2$, where $A = \pi r^2 = \pi (\frac{D}{2})^2 = \frac{\pi}{4}D^2$

How fast does the area change with respect to the diameter when the diameter is 10 meter?

Solution:
$$\frac{dA}{dD} = \frac{\pi}{4} 2D = \frac{\pi D}{2}$$

When D = 10 m, the area is changing at rate $\frac{\pi}{2}(10) = 5\pi m$

Motion along a line: displacement, velocity, speed and acceleration:

Suppose that object is moving along a coordinate line (say an s-axis), so that we know its position S on that line as a function of time:

$$S = f(t)$$

The displacement of the object over the time interval from t to $t + \Delta t$

$$\Delta S = f(t + \Delta t) - f(t)$$

And average velocity of the object over that time interval is

$$v_{av} = \frac{displacement}{travel time} = \frac{\Delta S}{\Delta t} = \frac{f(t + \Delta t) - f(t)}{\Delta t}$$

To find the body's velocity at the exact instant t, we take the limit of the average velocity over the interval from t to $t + \Delta t$ as Δt shrinks to Zero. This limit is the derivative of f with the respect to t.

Velocity (instantaneous velocity) is the derivative of position with respect to time. If body's position at time t is S = f(t), then the bodys velocity at time t is

$$V(t) = \frac{dS}{dt} = \lim_{\Delta t \to 0} \frac{f(t + \Delta t) - f(t)}{\Delta t}$$

Speed is the absolute value of velocity

$$Speed = |V(t)| = \left|\frac{dS}{dt}\right|$$

Acceleration is the derivative of velocity with respect to time

$$a(t) = \frac{dV}{dt} = \frac{d^2}{dt^2}$$

Example 15/ Free fall of a heavy ball released from rest at time = 0 sec , where the equation of free fall is $S = 4.9t^2$.

a- How many meters does the ball fall in first 2 sec?

b- What are the velocity, speed and acceleration?

Solution:

a- during the first 2 sec, the ball falls $S(2) = 4.9 (2)^2 = 19.6 m$ b- at any time t, $V(t) = S'(t) = \frac{d}{dt}(4.9t^2) = 9.8t$ at $t = 2 \ second \rightarrow V(2) = 9.8 \times 2 = 19.6 \ m/sec$ Speed = $|V(2)| = |19.6| = 19.6 \ m/sec$

acceleration $a(t) = V'(t) = S''(t) = 9.8 \ m/sec^2$

Derivatives of Trigonometric functions:

$$\frac{d}{dx}(sinx) = cosx , \quad \frac{d}{dx}(cosx) = -sinx$$
$$\frac{d}{dx}(tanx) = sec^{2}x , \quad \frac{d}{dx}(secx) = secx tanx$$
$$\frac{d}{dx}(cotx) = -csc^{2}x , \quad \frac{d}{dx}(cscx) = -cscx cotx$$
$$tanx = \frac{sinx}{cosx} , \quad cotx = \frac{cosx}{sinx} , \quad secx = \frac{1}{cosx} \text{ and } cscx = \frac{1}{sinx}$$

Example 16/ Prove that $\frac{d}{dx}(tanx) = sec^2 x$

Solution:
$$\frac{d}{dx}(tanx) = \frac{d}{dx}\left(\frac{sinx}{cosx}\right) = \frac{cosx \cdot cosx - sinx (-sinx)}{cos^2 x} = \frac{cos^2 x + sin^2 x}{cos^2 x} = \frac{1}{cos^2 x} = sec^2 x$$

Example 17/ Derivative the following trigonometric functions:

Solutions:

1-
$$y = x^2 - sinx \rightarrow \frac{dy}{dx} = 2x - cosx$$

2- $y = x^2 sinx \rightarrow y' = \frac{dy}{dx} = x^2 cosx + sinx(2x)$
3- $y = \frac{sinx}{x} \rightarrow y' = \frac{dy}{dx} = \frac{xcosx - sinx(1)}{x^2} = \frac{xcosx - sinx}{x^2}$

Example 18/ Find second derivative $y^{\prime\prime}$ for the function = secx ?

Solution: y = secx, $y' = \frac{dy}{dx} = secx tanx$

$$y'' = \frac{dy'}{dx} = \frac{d}{dx}(\sec x \tan x) = \sec x \frac{d}{dx} \tan x + \tan x \frac{d}{dx}(\sec x)$$
$$= \sec x(\sec^2 x) + \tan x(\sec x \tan x) = \sec^3 x + \sec x \tan^2 x$$

Derivative of a composition function:

The derivative of the composite function f(g(x))at x is the derivative of f at g(x) times the derivative of g at x called (Chain Rule), if f(u) is differentiable at the point u = g(x)and g(x) is differentiable at x. Then the composite function $(f \circ g)(x) = f(g(x))$ is differentiable at , and

$$(f \circ g)/(x) = f/(g(x)) \cdot g/(x)$$
 or $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

Example 19/ The function $y = \frac{3}{2}x = \frac{1}{2}(3x)$ is the composite of the functions $= \frac{1}{2}u$ and u = 3x. How are the derivatives of these functions related?

Solution:

We have $\frac{dy}{dx} = \frac{3}{2}$, $\frac{dy}{du} = \frac{1}{2}$ and $\frac{du}{dx} = 3$ Since $\frac{3}{2} = \frac{1}{2}$. 3, We see that $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

Example 20/ The function $y = 9x^4 + 6x^2 + 1 = (3x^2 + 1)^2$ is the composite of $y = u^2$ and $u = 3x^2 + 1$. Calculate derivatives?

Solution:

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \dots \dots \dots (1)$$
$$\frac{dy}{du} = 2u \dots \dots (2) , \qquad \frac{du}{dx} = 6x \dots \dots (3)$$

By substituting equations 2 and 3 in equation 1, then

$$\frac{dy}{dx} = 2u.6x = 2(3x^2 + 1).6x = 12x(3x^2 + 1) = 36x^3 + 12x$$

Example 21/ An object moves along the x - axis so that its position at any time $t \ge 0$ is given by $x(t) = \cos(t^2 + 1)$. Find the velocity of the object as a function of .

Solution:

The velocity $\frac{dx}{dt}$, x is a composite function Method 1/Let $t^2 + 1 = u$ $\therefore x = \cos u$, $\frac{dx}{dt} = \frac{dx}{du}\frac{du}{dt}$ $\frac{dx}{du} = -\sin u$, $\frac{du}{dt} = 2t$ $\therefore \frac{dx}{dt} = -\sin u \cdot 2t = -\sin(t^2 + 1) \cdot 2t = -2t\sin(t^2 + 1)$ Method 2/ $(f \circ g)(x) = f(g(x))$, $(f \circ g)/(x) = f/(g(x)) \cdot g/(x)$ $Velocity = \frac{dx}{dt} = -\sin(t^2 + 1) \cdot 2t$

Example 22/ Derivative $sin(x^2 + x)$ with respect to x.

Solution: $\frac{d}{dx}(x^2 + x) = \cos(x^2 + x).(2x + 1)$

Example 23/ Find the derivative of $g(t) = \tan(5 - \sin 2t)$.

Solution:
$$g'(t) = \frac{dg}{dt} = \frac{d}{dt} \tan(5 - \sin 2t) = \sec^2(5 - \sin 2t) \cdot \frac{d}{dt}(5 - \sin 2t) = \sec^2(5 - \sin 2t) \cdot \left(0 - \cos 2t \cdot \frac{d}{dt}(2t)\right) = \sec^2(5 - \sin 2t) \cdot \left(-\cos 2t \cdot 2t - \sin 2t\right) \cdot \left(-\cos 2t \cdot 2t\right) \cdot \left(-\cos 2t \cdot 2t - \sin 2t\right) \cdot \left(-$$

The chain rule with powers of a function:

If f is a differentiable function of u and if u is a differentiable function of x, then substituting y = f(u) into the chain rule formula

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \longrightarrow \frac{d}{dx} f(u) = f'(u) \frac{du}{dx}$$

If *n* is a positive or negative integer and $f(u) = u^n$

$$\therefore f'(u) = nu^{n-1} \quad \rightarrow \quad \frac{d}{dx}u^n = nu^{n-1}\frac{du}{dx}u^n$$

Example 24/ $\frac{d}{dx}(5x^3 - x^4)^7 = 7(5x^3 - x^4)^6 \cdot (15x^2 - 4x^3)$ Example 25/ $\frac{d}{dx}(\frac{1}{3x-2}) = \frac{d}{dx}(3x-2)^{-1} = -1(3x-2)^{-2} \cdot (3) = \frac{-3}{(3x-2)^2}$ Example 26/ $\frac{d}{dx}\sin(\frac{\pi x}{180}) = \frac{\pi}{180}\cos\frac{\pi x}{180}$ Example 27/ If x = 2t + 3 and $y = t^2 - 1$, find the value of $\frac{dy}{dx}$ at x = 6.

Solution: $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2t}{2} = t \dots \dots (1)$

from origin equation $x = 2t + 3 \rightarrow t = \frac{x - 3}{2} \dots \dots (2)$

By substituting equation 2 in equation 1, then $\frac{dy}{dx} = \frac{x-3}{2} = \frac{6-3}{2} = \frac{3}{2}$

Antiderivative and Integrals:

A function F is an antiderivative of f on an interval I if

$$F' = f(x)$$

Example 28/

$$f(x) = 2x \quad \rightarrow (integral)F(x) = x^2 , \therefore F'(x) = 2x = f(x)$$

 $g(x) = cosx \rightarrow (integral) G(x) = sinx , \therefore G'(x) = cosx = g(x)$

Antiderivative formulas:

No.	Function	General antiderivative	Notes
1	x^n	$\frac{x^{n+1}}{n+1} + c$	$n \neq -1$
2	sinkx	$-\frac{\cos kx}{k}+c$	$k \ a \ constant$, $k eq 0$
3	coskx	$\frac{sinkx}{k} + c$	$k \ a \ constant$, $k eq 0$
4	sec ² x	tanx + c	
5	csc^2x	-cotx + c	
6	secx tanx	secx + c	
7	cscx cotx	-cscx + c	

Example 29/ Find the general antiderivative of $f(x) = \frac{3}{\sqrt{x}} + \sin 2x$

Solution:
$$f(x) = 3(x)^{\frac{-1}{2}} + \sin 2x$$
, $F(x) = 3\frac{x^{\frac{-1}{2}+1}}{\frac{-1}{2}+1} + {\binom{2}{2}}\sin 2x = 3\frac{x^{\frac{1}{2}}}{\frac{1}{2}} + {\binom{1}{2}(2\sin 2x)} = 6\sqrt{x} + \frac{1}{2}(-\cos 2x) = 6\sqrt{x} - \frac{1}{2}(\cos 2x) + c$

Note/ The function $F(x) = x^2$ is not function has derivative 2x, also the function $x^2 + 1$ has derivative 2x, therefore, we can write 2x + 1, where *c* represent arbitrary constant, therefore the two functions different by constant *c*.

Example 30/f(x) = sinx that satisfies F(0) = 3

Solution: $F(x) = -\cos x + c \rightarrow F(0) = -\cos 0 + c \rightarrow 3 = -1 + c \rightarrow \therefore c = 4$, $\therefore F(x) = -\cos x + 4$

Integrals:

A special symbol is used to denote the collection of all antiderivatives of a function .

Definition: The set of all antiderivatives of f is the **indefinite integral** of f with respect to x ,

$$\int f(x)dx$$

Example 31/Indefinite integration done by term and rewriting the constant of integration $\int (x^2 - 2x + 5) dx$.

Solution: If we recognize that $\left(\frac{x^3}{3}\right) - x^2 + 5x$ is antiderivative of $x^2 - 2x + 5$. We can evaluate the integral as

 $\int (x^2 - 2x + 5)dx = \frac{x^3}{3} - x^2 + 5x + c$

where c represent arbitrary constant

$$Or \int (x^2 - 2x + 5)dx$$

= $\int x^2 dx - \int 2x \, dx + \int 5 \, dx$
= $\int x^2 dx - 2 \int x \, dx + 5 \int dx = (\frac{x^3}{3} + c_1) - 2(\frac{x^2}{2} + c_2)$
+ $5(x + c_3)$
 $\therefore \int (x^2 - 2x + 5)dx = \frac{x^3}{3} + c_1 - x^2 - 2c_2 + 5x + 5c_3$
 $C = c_1 - 2c_2 + 5c_3$
 $\therefore \int (x^2 - 2x + 5)dx = \frac{x^3}{3} - x^2 + 5x + C$

Example 32/ Find the following indefinite integrals, and check your answers by differentiation:

1-
$$\int (x+1)dx$$
, 2- $\int (5-6x)dx$, 3- $\int \left(3t^2 + \frac{t}{2}\right)dt$, 4- $\int (2x^3 - 5x+7)dx$, 5- $\int x^{\frac{-1}{3}}dx$

Solution:

$$1-\int (x+1)dx = \int xdx + \int dx = \frac{x^2}{2} + x + c$$

$$2-\int (5-6x)dx = \int 5dx - 6\int xdx = 5x - 6\frac{x^2}{2} = 5x - 3x^2 + c$$

$$3-\int \left(3t^2 + \frac{t}{2}\right)dt = \int 3t^2dt + \frac{1}{2}\int tdt = 3\frac{t^3}{3} + \frac{1}{2}\frac{t^2}{2} = t^3 + \frac{1}{4}t^2 + c$$

$$4-\int (2x^3 - 5x + 7)dx = 2\int x^3dx - 5\int xdx + 7\int dx = 2\frac{x^4}{4} - 5\frac{x^2}{2} + 7x = \frac{1}{2}x^4 - \frac{5}{2}x^2 + 7x + c$$

$$5-\int x\frac{-1}{3}dx = \frac{x\frac{-1}{3}+1}{\frac{-1}{3}+1} = \frac{2}{3}x^{\frac{2}{3}} + c$$

H.W 1/ Find the indefinite integrals:

$$\begin{array}{l} 1-\int -2costdt , 2-\int 7\sin\left(\frac{\theta}{3}\right)d\theta , 3-\int 3cos5\theta d\theta , 4-\\ \int \frac{2}{5}sec\theta tan\theta d\theta , 5-\int \frac{csc\theta cot\theta}{2}d\theta \end{array}$$

H.W 2/ Verify the following formulas by differentiation:

$$1- \int (7x-2)^3 dx = \frac{(7x-2)^4}{28} + c \quad , \ 2- \int (3x+5)^{-2} dx = -\frac{(3x+5)^{-1}}{3} + c$$
$$3- \int \sec^2(5x-1) dx = \frac{1}{5} \tan(5x-1) + c \quad , \ 4- \int \csc^2\left(\frac{x-1}{3}\right) dx = -3\cot\left(\frac{x-1}{3}\right) + c \quad , \ 5- \int \frac{1}{(x+1)^2} dx = \frac{1}{x+1} + c$$

H.W 3/ Solve the initial value problems in the following formulas:

$$1 - \frac{dy}{dx} = 2x - 7 , \quad y(2) = 0$$

$$2 - \frac{dy}{dx} = 10 - x , \quad y(0) = -1$$

$$3 - \frac{dy}{dx} = \frac{1}{x^{2}} + x , \quad x > 0 ; \quad y(2) = 1$$

$$4 - \frac{ds}{dt} = 1 + \cos t , \quad s(0) = 4$$

$$5 - \frac{dv}{dt} = \frac{1}{2} \operatorname{sect tant} , \quad v(0) = 1$$

Definite integral:

Let f(x) be a function defined on a closed interval [a, b], we say that a number I is the definite integral of f over [a, b].

The symbol for the number *I* in the definition of the definite integral is

$$\int_{a}^{b} f(x) dx$$
 , or $\int_{a}^{b} f(t) dt$ or $\int_{a}^{b} f(u) du$

Properties of definite integrals:

1-
$$\int_{a}^{b} f(x)dx = -\int_{b}^{a} f(x)dx$$

2- $\int_{a}^{b} f(x)dx = 0$ when $a = b$

3-
$$\int_{a}^{b} kf(x)dx = k \int_{a}^{b} f(x)dx$$
 constant multiple k.
4-

$$\int_{a}^{b} (f(x) \pm g(x)) dx = \int_{a}^{b} f(x) dx \pm \int_{a}^{b} g(x) dx \text{ sum and differences}$$

5-
$$\int_{a}^{b} f(x) dx + \int_{b}^{c} f(x) dx = \int_{a}^{c} f(x) dx \text{ Additivity}$$

Example 33/ Find the following definite integrals if:

$$\int_{-1}^{1} f(x)dx = 5 , \qquad \int_{1}^{4} f(x)dx = -2 , \qquad \int_{-1}^{1} h(x)dx = 7$$

$$1 - \int_{4}^{1} f(x)dx = -\int_{1}^{4} f(x)dx = -(-2) = 2$$

$$2 - \int_{-1}^{1} [2f(x) + 3h(x)]dx = 2 \int_{-1}^{1} f(x)dx + 3 \int_{-1}^{1} h(x)dx = 2(5) + 3(7) = 31$$

$$3 - \int_{-1}^{4} f(x)dx = \int_{-1}^{1} f(x)dx + \int_{1}^{4} f(x)dx = 5 + (-2) = 3$$

H.W 4/Using properties of integrals and known values to find other integrals, suppose that f and g are integrable and that equal:

$$\int_{1}^{2} f(x)dx = -4 , \quad \int_{1}^{5} f(x)dx = 6 , \quad \int_{1}^{5} g(x)dx = 8$$

Find the following integrals

$$\int_{2}^{2} f(x)dx = -4 , \quad \int_{5}^{1} g(x)dx , \quad \int_{1}^{2} 3f(x)dx , \quad \int_{2}^{5} f(x)dx$$

Solution:

$$\int_{2}^{2} f(x)dx = \int_{2}^{1} f(x)dx + \int_{1}^{2} f(x)dx = -\int_{1}^{2} f(x)dx + \int_{1}^{2} f(x)dx = -(-4) + (-4) = 0$$

H.W 5/ Prove that $\int_{-\pi}^{\pi} \cos x \, dx = 0$ $\int_{0}^{\frac{\pi}{4}} \sec^2 x \, dx = 1$

H.W 6/ Evaluate the following indefinite integrals by using the given substitutions to reduce the integrals to standard form.

$$1 - \int \sin 3x \, dx$$
, $u = 3x$; $2 - \int x \sin(2x^2) dx$, $u = 2x^2$

Solution:

$$1-\int \sin 3x \, dx \quad ; u = 3x \to x = \frac{1}{3}u \to dx = \frac{1}{3}du$$

$$\therefore \int \sin 3x \, dx = \frac{1}{3}\int \sin u \, du = \frac{1}{3}(-\cos u) = -\frac{1}{3}\cos(3x)$$

$$Or \quad \int \sin 3x \, dx = \frac{3}{3}\int \sin 3x \, dx = \frac{1}{3}(-\cos u) = -\frac{1}{3}\cos(3x)$$

Example 34/ Evaluate the following definite integrals by using the given substitutions to reduce the integrals to standard form.

$$\int_{x=-1}^{x=1} 3x^2 \sqrt{x^3 + 1} \, dx \quad , \quad Let \ u = x^3 + 1$$

Solution:

method1:
$$u = x^3 + 1 \rightarrow du = 3x^2 dx \rightarrow dx = \frac{1}{3x^2} du$$

when $x = -1$ $\therefore u = (-1)^3 + 1 = 0$
and when $x = 1$ $\therefore u = (1)^3 + 1 = 2$
 $\therefore \int_{x=-1}^{x=1} 3x^2 \sqrt{x^3 + 1} dx = \int_{u=0}^{u=2} 3x^2 \sqrt{u} \frac{1}{3x^2} du = \int_{u=0}^{u=2} u^{\frac{1}{2}} du = \frac{u^{\frac{1}{2}+1}}{\frac{1}{2}+1} \Big]_0^2 = \frac{2}{3} u^{\frac{3}{2}} \Big]_0^2 = \frac{2}{3} \Big[2^{\frac{3}{2}} - 0^{\frac{3}{2}} \Big] = \frac{2}{3} \Big[2\sqrt{2} \Big] = \frac{4\sqrt{2}}{3}$

Method2: transform the integral as an indefinite integral change back to x, and use the original x - limits.

$$\int_{x=-1}^{x=1} 3x^2 \sqrt{x^3 + 1} \, dx = \int \sqrt{u} \, du = \frac{2}{3} u^{\frac{3}{2}} + c = \frac{2}{3} (x^3 + 1)^{\frac{3}{2}} + c$$
$$\therefore \int_{x=-1}^{x=1} 3x^2 \sqrt{x^3 + 1} \, dx = \frac{2}{3} (x^3 + 1)^{\frac{3}{2}} \Big]_{-1}^{-1} = \frac{4\sqrt{2}}{3}$$

H.W 7/ Evaluate the following definite integrals by using the given substitutions to reduce the integrals to standard form.

$$\int_{rac{\pi}{4}}^{rac{\pi}{2}} cot heta csc^2 heta \; d heta$$
 , Let $u=cot heta$

Note:

a- If f is even, then $\int_{-a}^{a} f(x) dx = 2 \int_{0}^{a} f(x) dx$ b- If f is odd, then $\int_{-a}^{a} f(x) dx = 0$ Example 35/ $\int_{-2}^{2} x dx = 0$

Example 36/ $\int_{-2}^{2} x^{2} dx = 2 \int_{0}^{2} x^{2} dx = 2 \frac{x^{3}}{3} \Big]_{0}^{2} = \frac{2}{3} [2^{3} - 0^{3}] = \frac{16}{3}$

Area between curves:

If *f* and *g* are continuous with $f(x) \ge g(x)$ throughout [a, b], then the area of the region between the curves y = f(x) and y = g(x) from *a* to *b* is the integral of (f - g) from *a* to *b*.

$$A = \int_{a}^{b} [f(x) - g(x)] \, dx$$

Example 37/ Find the area of the region enclosed by the parabola $y = 2 - x^2$ and the line y = -x.

Solution: The limits of integration are found by solving $y = 2 - x^2$ and the line y = -x for x.

$$2 - x^{2} = -x \rightarrow x^{2} - x - 2 = 0 \rightarrow (x - 2)(x + 1) = 0$$

either $x - 2 = 0 \rightarrow \therefore x = 2$ or $x + 1 = 0 \rightarrow \therefore x = -1$

The region runs from x = -1 to x = 2, and the limits of integration are a = -1 and b = 2.



The area between curves is

$$A = \int_{a}^{b} [f(x) - g(x)] \, dx = \int_{-1}^{2} [(2 - x^2) - (-x)] \, dx$$
$$A = \int_{-1}^{2} (2 - x^2 + x) \, dx = \left[2x - \frac{x^3}{3} + \frac{x^2}{2}\right]_{-1}^{2} = \left[4 - \frac{8}{3} + \frac{4}{2}\right] - \left[-2 + \frac{1}{3} + \frac{1}{2}\right] = \frac{27}{6} = 4.5 \ (unit)^2$$

Example 38/ Find the area of the region enclosed by the parabola $y = x^2$ and the line y = x + 2.

Solution: The limits of integration are found by solving $y = x^2$ and the line y = x + 2 for x.

$$x^{2} = x + 2 \rightarrow x^{2} - x - 2 = 0 \rightarrow (x - 2)(x + 1) = 0$$

either $x - 2 = 0 \rightarrow \therefore x = 2$ or $x + 1 = 0 \rightarrow \therefore x = -1$

The region runs from x = -1 to x = 2, and the limits of integration are a = -1 and b = 2.



The area between curves is

$$A = \int_{a}^{b} [f(x) - g(x)] dx = \int_{-1}^{2} [(x+2) - x^{2}] dx$$
$$A = \int_{-1}^{2} (x+2 - x^{2}) dx = \left[\frac{x^{2}}{2} + 2x - \frac{x^{3}}{3}\right]_{-1}^{2}$$
$$= \left[2 + 4 - \frac{8}{3}\right] - \left[\frac{1}{2} - 2 + \frac{1}{3}\right] = \frac{27}{6} = 4.5 \ (unit)^{2}$$

Area under a curve by integration:

Case1- Curves which are entirely above the x - axis:

In this case, we find the area by simply finding the integral

$$A = \int_{a}^{b} f(x) \, dx$$

Where the area under the curve y = f(x) from x = a to x = b

Case2- Curves which are entirely below the x - axis:

In this case, the integral gives a negative number, we need to take the absolute value of this to find our area.

$$A = \left| \int_{a}^{b} f(x) \, dx \right|$$

Case3- Part of the curve is below the x - axis, part of it is above the x - axis:

In this case, we have to sum the individual parts, taking the absolute value for the section where the curve is below the x - axis from (x = a to x = c).

$$A = \left| \int_{a}^{c} f(x) \, dx \right| + \int_{c}^{d} f(x) \, dx$$

If we don't do it like this, the "negative" area (the part below the x - axis) will be subtract from the "positive" part, and our total area will not be correct.

Case4- Certain Curves are much easier to sum vertically:

In some cases, it is easier to find the area if we take vertical sums. Sometimes the only possible way is to sum. We need to re-express this as x = f(y), and we need to sum from bottom to top. So,

$$A = \int_{c}^{d} f(y) \, dy$$

Area between two curves using integration:

Area bounded by the curves y_1 and y_2 and the lines x = a and x = b. We see that if we subtract the area under lower curve $y_1 = f_1(x)$, from the area under the upper curve $y_2 = f_2(x)$, then we will find the required area. This can be achieved in one step

$$A = \int_{a}^{b} (y_2 - y_1) dx$$

Likewise, we can sum vertically by re-expressing both functions so that they are functions of y, and we find

$$A = \int_c^d (x_2 - x_1) dy$$

Where, c and d as the limits on the integral.

Example39/ Find the area underneath the curve $y = x^2 + 2$ from x = 1 to x = 2.



Solution: $A = \int_{1}^{2} (x^{2} + 2) dx = \left[\frac{x^{3}}{3} + 2x\right]_{1}^{2} = \left[\frac{8}{3} + 4\right] - \left[\frac{1}{3} + 2\right] = \left[\frac{20}{3}\right] - \left[\frac{7}{3}\right] = \frac{13}{3} unit^{2}$

Example40/ Find the area bounded by $y = x^2 - 4$, the axis and the lines x = -1 and x = 2.

Solution:



$$A = \left| \int_{-1}^{2} (x^2 - 4) dx \right| = \left[\frac{x^3}{3} - 4x \right]_{-1}^{2} = \left[\frac{8}{3} - 8 \right] - \left[-\frac{1}{3} + 4 \right]$$
$$= \left[\frac{-16}{3} \right] - \left[\frac{-1 + 12}{3} \right] = \frac{-16}{3} - \frac{11}{3} = \frac{-27}{3} = |-9|$$
$$= 9unit^2$$

Example41/ What the area bounded by the curve $y = x^3$ and the lines x = -2 and x = 1.

Solution:



$$A = \left| \int_{-2}^{0} x^3 \, dx \right| + \int_{0}^{1} x^3 \, dx = \left| \left[\frac{x^4}{4} \right]_{-2}^{0} \right| + \left[\frac{x^4}{4} \right]_{0}^{1} = \left| 0 - \frac{16}{4} \right| + \left[\frac{1}{4} \right]_{-2}^{1} \right|$$
$$= 4 + \frac{1}{4} = 4.25$$

Example42/Find the area of the region bounded by the curve $y = \sqrt{x-1}$, the y – axis and the lines y = 1 and y = 5.

Solution:



In this case, we express *x* as a function of *y* :

$$y = \sqrt{x - 1}$$
, $\to y^2 = x - 1$, $\to x = y^2 + 1$

So, the area is given by:

$$A = \int_{1}^{5} (y^{2} + 1) dy = \left[\frac{y^{3}}{3} + y\right]_{1}^{5} = \left[\frac{5^{3}}{3} + 5\right] - \left[\frac{1^{3}}{3} + 1\right]$$
$$= \left[\frac{125}{3} + 5\right] - \left[\frac{1}{3} + 1\right] = \frac{140}{3} - \frac{4}{3} = \frac{136}{3} = 45\frac{1}{3}$$

Example43/ Find the area bounded by $y = x^3$, x = 0 and y = 3.

Solution: we must convert

x = 0 to y function, where substitute it in function $y = x^3$, to get on y = 0, also we must convert the function $y = x^3$ to $x = \sqrt[3]{y} = y^{\frac{1}{3}}$

The area equal $A = \int_{c}^{d} f(y) dy = \int_{0}^{3} y^{\frac{1}{3}} dy = \left[\frac{y^{\frac{1}{3}+1}}{\frac{1}{3}+1}\right]_{0}^{3} = \left[\frac{y^{\frac{4}{3}}}{\frac{4}{3}}\right]_{0}^{3} = \frac{3}{4}\left[(3)^{\frac{4}{3}} - (0)^{\frac{4}{3}}\right] = \frac{3}{4}\left[(3)^{\frac{4}{3}}\right] = 3.245 \text{ unit}^{2}$

Or another method : We substitute y = 3 in $y = x^3$, to get $x = \sqrt[3]{3}$ And the integral limitations equal x = 0 and $x = \sqrt[3]{3}$.

Then , we can use horizontally method to find area between the above function y = 3 and the below function $y = x^3$.



The area equal $A = \int_0^{\sqrt[3]{3}} (3-x^3) dx = \left[3x - \frac{x^4}{4}\right]_0^{\sqrt[3]{3}} = \left[3\sqrt[3]{3} - \frac{(\sqrt[3]{3})^4}{4}\right] - [0] = [4.326 - 1.08] = 3.245 \ unit^2$

Example44/ Find the area between the curves $y = x^2 + 5x$ and $y = 3 - x^2$ between x = -2 and x = 0.

Solution:

$$A = \int_{-2}^{0} [(3 - x^2) - (x^2 + 5x)] dx = \int_{-2}^{0} [-2x^2 - 5x + 3] dx$$
$$= \left[\frac{-2x^3}{3} - \frac{5x^2}{2} + 3x\right]_{-2}^{0} = [0] - \left[\frac{16}{3} - 10 - 6\right]$$
$$= 0 - \left[\frac{16}{3} - 16\right] = -\left[\frac{-32}{3}\right] = \frac{32}{3} = 10\frac{2}{3} \text{ unit}^2$$

Note/ some of the shaded area is above the x - axis and some of it is below, therefore don't worry about taking absolute value, where the formula takes care of that automatically.

Note/ we can take any from the functions if it below or above x - axis, and if we get on the negative value of area, then we can take the absolute value of it.

Note/ if in question gave you only two functions for the curves, then from equating them, you get on the integral limitations x_1 and x_2 or y_1 and y_2 .

Example45/ Find the area between the curves $y = x^2 + 5x$ and $y = 3 - x^2$.

Solution: by equating the two function to get on the integral limitations

$$x^{2} + 5x = 3 - x^{2} \rightarrow x^{2} + x^{2} + 5x - 3 = 0 \rightarrow 2x^{2} + 5x - 3 = 0$$

$$\therefore (2x-1)(x+3) = 0 , either 2x - 1 = 0, \therefore x = 0.5, or x + 3 = 0, \\ \therefore x = -3$$

$$A = \int_{-3}^{0.5} [(x^2 + 5x) - (3 - x^2)] dx = \int_{-3}^{0.5} (2x^2 + 5x - 3) dx$$
$$= \left[\frac{2x^3}{3} + \frac{5x^2}{2} - 3x\right]_{-3}^{0.5}$$
$$= \left[\frac{2(0.5)^3}{3} + \frac{5(0.5)^2}{2} - 3(0.5)\right] at$$
$$= [0.08 + 0.625 - 1.5] - [18 + 22.5 + 9]$$
$$= [-0.795] - [13.5] = |-14.295| = 14.295 unit^2$$

Volume of Solid:

Many solid objects, especially those made on a lathe have a circular cross-section and curved sides. In this section, we see how to find the volume of such objects using integration.

Example46/ Consider the area bounded by the straight line y = 3x, x - axis, and x = 1.

Solution: when the shaded is rotated 360° about the x - axis, a volume is generated, and the resulting solid is a Cone.

To find this volume, we could take slices

The typical disk with dimensions, radius = y, and height = dx, where the volume of Cylinder is given by $V = \pi r^2 h$, and because radius r = y, and height dx = h, then

$$V = \pi y^2 dx$$

Adding the volumes of the disks (with infinitely small dx, we obtain the following formula:

$$V = \pi \int_{a}^{b} y^{2} dx$$
, which means
 $V = \pi \int_{a}^{b} [f(x)]^{2} dx$

Where, y = f(x) is the equation of the curve whose area is being rotated. *a and b* are limits of the area being rotated. *dx* shows that the area is being rotated about the x - axis.

By applying volume formula to the earlier example, we have:

$$V = \pi \int_{a}^{b} y^{2} dx = V = \pi \int_{0}^{1} (3x)^{2} dx = 9\pi \int_{0}^{1} x^{2} dx = 9\pi \left[\frac{x^{3}}{3}\right]_{0}^{1}$$
$$= 3\pi (unit)^{3}$$

And we can find the volume of the cone using the following:

$$V = \frac{1}{3}\pi r^2 h = \frac{\pi}{3}(3)^2 \cdot 1 = 3\pi \text{ (unit)}^3 \text{ (Check ok.)}$$

Example47/ Find the volume if the area bounded by the curve $y = x^3 + 1$, *the* x - axis, and the limits of x = 0 and x = 3 is rotated around *the* x - axis.

$$V = \pi \int_{a}^{b} y^{2} dx = \pi \int_{0}^{3} (x^{3} + 1)^{2} dx = \pi \int_{0}^{3} (x^{6} + 2x^{3} + 1)^{2} dx$$
$$= \pi \int_{0}^{3} [x^{6} dx + 2x^{3} dx + dx] = \pi \left[\frac{x^{7}}{7} + 2\frac{x^{4}}{4} + x \right]_{0}^{3}$$
$$= \pi [312.4 + 40.25] = 355.9 \pi (unit)^{3}$$

And the area can be finding by the following:

$$A = \int_0^3 (x^3 + 1)dx = \left[\frac{x^4}{4} + x\right]_0^3 = \left[\frac{81}{4} + 3\right] - [0] = 43.25 \ (unit)^2$$

Volume by rotating the area enclosed between two curves:

If we have two curves y_2 and y_1 that enclose some area and we rotate that area around the x - axis, then the volume of the solid formed is given by

$$V = \pi \int_{a}^{b} [(y_2)^2 - (y_1)^2] \, dx$$

Where the limits for the region indicated by the vertical lines at x = a and x = b. y_1 and y_2 , represent lower and upper functions, respectively.

Example48/ A cup is made by rotating the area between functions $y = 2x^2$ and y = x + 1 with $x \ge 0$ around the x - axis. Find the volume of the material needed to make the cup. Units are *cm*.

Solution: to find integration limits, we equate the two functions, then we get

$$x + 1 = 2x^2 \rightarrow 2x^2 - x - 1 = 0 \rightarrow (2x + 1)(x - 1) = 0$$

either $(2x + 1) = 0 \rightarrow \therefore x = \frac{-1}{2}$ (neglible because we have $x \ge 0$)

or
$$(x - 1) = 0 \rightarrow \therefore x = 1$$
, therefore the limitations are $x = 0$ and $x = 1$

$$V = \pi \int_0^1 [(x+1)^2 - (2x^2)^2] \, dx = \pi \int_0^1 [x^2 + 2x + 1 - 4x^4] \, dx$$
$$= \pi \left[\frac{x^3}{3} + 2\frac{x^2}{2} + x - 4\frac{x^5}{5} \right]_0^1 = \pi \left[\frac{1}{3} + 1 + 1 - \frac{4}{5} \right]$$
$$= \pi \left[\frac{5 + 15 + 15 - 12}{15} \right] = \frac{23}{15}\pi \ cm^3$$

Rotation around the y-axis:

When the shaded area is rotated 360° about the y - axis, the volume that is generated can be found by:

$$V = \pi \int_c^d x^2 dy$$
, which means $V = \pi \int_c^d [f(y)]^2 dy$

Where: x = f(y) is the equation of the curve expressed in terms of y. c and d are the upper and lower y limits of the area being rotated. dy shows that the area is being rotated about the y - axis.

Example49/ Find the volume of the solid of revolution generated by rotating the curve $y = x^3$ between y = 0 and y = 4 about the y - axis.

Solution: If $y = x^3$, then $x = \sqrt[3]{y} = y^{\frac{1}{3}}$

$$V = \pi \int_0^4 (y^{\frac{1}{3}})^2 \, dy = \pi \int_0^4 y^{\frac{2}{3}} \, dy = \pi \left[\frac{y^{\frac{2}{3}+1}}{\frac{2}{3}+1} \right]_0^4 = \pi \left[\frac{y^{\frac{5}{3}}}{\frac{5}{3}} \right]_0^4 = \pi \frac{3}{5} \left[y^{\frac{5}{3}} \right]_0^4$$
$$= \pi \frac{3}{5} \left[4^{\frac{5}{3}} - 0^{\frac{5}{3}} \right] = \pi \frac{3}{5} \left[10.079 - 0 \right] = 19 \ (unit)^3$$

Example50/ Find the volume of the solid of revolution generated by rotating the curve y = x between y = 0 and x = 2 about the y - axis.

Solution: we can find the volume by two methods, in this case take about y - axis, therefore *if* y = x *become* x = y

the limitations between
$$y = 0$$
, and we substitute $x = 2$ in equation $y = x$, then get on $y = 2$

$$V = \pi \int_0^2 y^2 \, dy = \pi \left[\frac{y^3}{3} \right]_0^2 = \frac{\pi}{3} [2^3 - 0^3] = \frac{\pi}{3} [8 - 0] = \frac{8\pi}{3}$$
$$= 8.378 \ (unit)^3$$

Arc Length of a Curve by Using Integration:

If the horizontal distance is dx (a small change in x) and the vertical height of the triangle is dy (a small change in y), then the length of the curved *arc* dr is approximated as:

$$dr = \sqrt{(dx)^2 + (dy)^2}$$

And this equation represents general form of the length of the curve.

The arc length of the curve y = f(x) from x = a to x = b is given by:

$$length = r = \int_{a}^{b} \sqrt{1 + (\frac{dy}{dx})^2} \, dx$$

Note/ we are assuming the function y = f(x) is continuous in the region x = a to x = b (otherwise, the formula won't work).

Arc Length by Using Radian Measurement:

In this section, we see some of the common applications of radian measure, including arc length, area of sector of a circle, and angular velocity.

The length s, of an arc of a circle radius r subtended by θ (in radians) is given by:

$$s = r\theta$$

If r is in meters, s will also be in meters.

Example51/ Find the length of the *arc* of a circle with radius 4 *cm* and central angle 5.1 *radians*.

Solution:

$$s = r\theta = 4 \times 5.1 = 20.4 \ cm$$

Area of Sector:

The area of a sector with central angle θ (in radians) is given by:

$$Area = \frac{\theta r^2}{2}$$

If r is measured in cm, the area will be in cm^2 .

Example 52/ Find the area of the sector with radius 7*cm* and central angle 2.5 *radians*.

Solution: $Area = \frac{\theta r^2}{2} = \frac{2.5 \times 7^2}{2} = 61.25 \ cm^2$

Angular velocity: The time rate of change of angle θ by a rotating body is the angular velocity, written ω . It is measured in *radius/second*.

If v is a linear velocity in $(\frac{m}{s})$ and r is the radius of the circle in (m), then

$$v = r\omega$$

Example53/ A bicycle with tyres 90 *cm* in diameter is travelling at 25 *Km/h*. What is the angular velocity of the tyre in *radians per second*.

Solution: we convert the units to meters

$$25\frac{Km}{h} = \frac{25000}{3600} = 6.94 \text{ m/s}$$
$$r = \frac{90}{2} = 45 \text{ cm} = 0.45 \text{ m}$$
So, $\omega = \frac{v}{r} = \frac{6.94}{0.45} = 15.43 \text{ rad/sec}$

Finite sums:

Sigma notation enables us to write a sum with many terms in the compact form

$$\sum_{k=1}^{n} a_k = a_1 + a_2 + a_3 + \dots + a_{n-1} + a_n$$

The index k ends at k=n

Summation symbol $\rightarrow \sum_{k=1}^{n} a_k$ is a formula for the k_{th} term

Index k start at k=1

Example 36/

$$\sum_{k=1}^{11} k^2 = 1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2 + 7^2 + 8^2 + 9^2 + 10^2 + 11^2$$

$$\sum_{k=1}^{5} 2^{k} = 2^{1} + 2^{2} + 2^{3} + 2^{4} + 2^{5}$$

H.W 8/ Write the sums of the following notations:

$$1 - \sum_{k=1}^{2} \frac{6k}{k+1} , \quad 2 - \sum_{k=1}^{4} \cos k\pi , \quad 3 - \sum_{k=1}^{5} \sin k\pi , \quad 4 - \sum_{k=1}^{6} 2^{k-1}$$

$$5 - \sum_{k=0}^{5} 2^{k} , \quad 6 - \sum_{k=0}^{4} (2k+1) , \quad 7 - \sum_{k=1}^{2} \frac{k}{k+1}$$

Solution :

7-
$$\sum_{k=1}^{2} \frac{k}{k+1} = \frac{1}{1+1} + \frac{2}{2+1} = \frac{1}{2} + \frac{2}{3} = \frac{7}{6}$$

Algebra rules for finite sums:

- 1- Sum rules: $\sum_{k=1}^{n} (a_k + b_k) = \sum_{k=1}^{n} a_k + \sum_{k=1}^{n} b_k$
- 2- Difference rule: $\sum_{k=1}^{n} (a_k b_k) = \sum_{k=1}^{n} a_k \sum_{k=1}^{n} b_k$
- 3- Constant multiple rule: $\sum_{k=1}^{n} Ca_k = C \cdot \sum_{k=1}^{n} a_k$
- 4- Constant value rule: $\sum_{k=1}^{n} C = n.C$, $Ex / \sum_{k=1}^{3} 5 = 3.5 = 15$