


---

### Partitions of set:

Let  $S$  be any nonempty set. A partition ( $\Pi$ ) of  $S$  is a subdivision of  $S$  into nonoverlapping, nonempty subsets. A partition of  $S$  is a collection  $\{A_i\}$  of non-empty subsets of  $S$  such that:

- 1)  $A_i \neq \emptyset$ , where  $i=1,2,3,\dots$
- 2) The sets of  $\{A_i\}$  are mutually disjoint  
or  $A_i \cap A_j = \emptyset$  where  $i \neq j$ .
- 3)  $\cup A_i = S$ , where  $A_1 \cup A_2 \cup \dots \cup A_i = S$

The partition of a set into five cells,  $A_1, A_2, A_3, A_4, A_5$ , can be represented by Venn diagram



Example 1: let  $A = \{1,2,3,n\}$

$$A_1 = \{1\}, A_2 = \{3,n\}, A_3 = \{2\}$$

$\Pi = \{A_1, A_2, A_3\}$  is a partition on  $A$  because it satisfies the three above conditions.

**Example 2 :** Consider the following collections of subsets of  $S = \{1,2,3,4,5,6,7,8,9\}$

- (i)  $\{\{1,3,5\}, \{2,6\}, \{4,8,9\}\}$
- (ii)  $\{\{1,3,5\}, \{2,4,6,8\}, \{5,7,9\}\}$
- (iii)  $\{\{1,3,5\}, \{2,4,6,8\}, \{7,9\}\}$

Then

- (i) is not a partition of  $S$  since 7 in  $S$  does not belong to any of the subsets.
- (ii) is not a partition of  $S$  since  $\{1,3,5\}$  and  $\{5,7,9\}$  are not disjoint.
- (iii) is a partition of  $S$ .

---

### FINITE SETS, COUNTING PRINCIPLE:

A set is said to be finite if it contains exactly  $m$  distinct elements where  $m$  denotes some nonnegative integer. Otherwise, a set is said to be infinite. For example, the empty set  $\emptyset$  and the set of letters of English alphabet are finite sets, whereas the set of even positive integers,  $\{2,4,6,\dots\}$ , is infinite.

If a set  $A$  is finite, we let  $n(A)$  or  $\#(A)$  denote the number of elements of  $A$ .

Example: If  $A = \{1,2,a,w\}$  then

$$n(A) = \#(A) = |A| = 4$$

Lemma: If  $A$  and  $B$  are finite sets and disjoint Then  $A \cup B$  is finite set and:

$$n(A \cup B) = n(A) + n(B)$$

**Theorem (Inclusion–Exclusion Principle):** Suppose  $A$  and  $B$  are finite sets. Then  $A \cup B$  and  $A \cap B$  are finite and

$$|A \cup B| = |A| + |B| - |A \cap B|$$

That is, we find the number of elements in A or B (or both) by first adding  $n(A)$  and  $n(B)$  (inclusion) and then subtracting  $n(A \cap B)$  (exclusion) since its elements were counted twice.

We can apply this result to obtain a similar formula for three sets:

**Corollary:**

If A,B,C are finite sets then

$$|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$$

**Example (1) :**

$$A = \{1,2,3\}$$

$$B = \{3,4\}$$

$$C = \{5,6\}$$

$$A \cup B \cup C = \{1,2,3,4,5,6\}$$

$$|A \cup B \cup C| = 6$$

$$|A| = 3, \quad |B| = 2, \quad |C| = 2$$

$$A \cap B = \{3\}, \quad |A \cap B| = 1$$

$$A \cap C = \{\}, \quad |A \cap C| = 0$$

$$B \cap C = \{\}, \quad |B \cap C| = 0$$

$$A \cap B \cap C = \{\}, \quad |A \cap B \cap C| = 0$$

$$|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$$

$$|A \cup B \cup C| = 3 + 2 + 2 - 1 - 0 - 0 + 0 = 6$$

**Example (2):**

Suppose a list A contains the 30 students in a mathematics class, and a list B contains the 35 students in an English class, and suppose there are 20 names on both lists. Find the number of students:

(a) only on list A

(b) only on list B

(c) on list  $A \cup B$

Solution:

(a) List A has 30 names and 20 are on list B; hence  $30 - 20 = 10$  names are only on list A.

(b) Similarly,  $35 - 20 = 15$  are only on list B.

(c) We seek  $n(A \cup B)$ . By inclusion-exclusion,

$$n(A \cup B) = n(A) + n(B) - n(A \cap B) = 30 + 35 - 20 = 45.$$

**Example (3):**

Suppose that 100 of 120 computer science students at a college take at least one of languages: French, German, and Russian and:

65 study French (F).

45 study German (G).

42 study Russian (R).


20 study French & German  $F \cap G$ .

25 study French & Russian  $F \cap R$ .

15 study German & Russian  $G \cap R$ .

Find the number of students who study:

- 1) All three languages ( $F \cap G \cap R$ )
- 2) The number of students in each of the eight regions of the Venn diagram



Solution:

$$\begin{aligned}
 |F \cup G \cup R| &= |F| + |G| + |R| - |F \cap G| - |F \cap R| - |G \cap R| + |F \cap G \cap R| \\
 100 &= 65 + 45 + 42 - 20 - 25 - 15 + |F \cap G \cap R| \\
 100 &= 92 + |F \cap G \cap R|
 \end{aligned}$$

$\therefore |F \cap G \cap R| = 8$  students study the 3 languages

$$20 - 8 = 12 \quad (F \cap G) - R$$

$$25 - 8 = 17 \quad (F \cap R) - G$$

$$15 - 8 = 7 \quad (G \cap R) - F$$

$$65 - 12 - 8 - 17 = 28 \quad \text{students study French only}$$

$$45 - 12 - 8 - 7 = 18 \quad \text{students study German only}$$

$$42 - 17 - 8 - 7 = 10 \quad \text{students study Russian only}$$

$$120 - 100 = 20 \quad \text{students do not study any language}$$