



جامعة الانبار

كلية العلوم

قسم علوم الحياة

Bacterial toxins

Assistant .prof .Dr.Muthanna Hamid Hassan

Lecture 1

Introduction of bacterial toxins

مقدمه عن السموم البكتيرية

Bacterial Toxins Lec:1

Introduction

A toxin is a poisonous substance produced within living cells or organisms. toxins can be small molecules, peptides, or proteins that are capable of causing disease on contact with or absorption by body tissues interacting with biological macromolecules such as enzymes or cellular receptors. Toxins were the first bacterial virulence factors to be identified and were also the first link between bacteria and cell biology.

Bacterial Toxigenesis

Toxigenesis, or the ability to produce toxins, is an underlying mechanism by which many bacterial pathogens produce disease. At a chemical level, there are two main types of bacterial toxins, **lipopolysaccharides**, which are associated with the cell wall of Gram-negative bacteria, and **proteins**, which are released from bacterial cells and may act at tissue sites removed from the site of bacterial growth. The **cell-associated** toxins are referred to as **endotoxins** and the **extracellular** diffusible toxins are referred to as **exotoxins**.

Endotoxins are cell-associated substances that are structural components of bacteria. Most endotoxins are located in the cell envelope. In the context of this article, endotoxin refers specifically to the lipopolysaccharide (LPS) or lipooligosaccharide (LOS) located in the outer membrane of Gram-negative bacteria. Although structural components of cells, soluble endotoxins may be released from growing bacteria or from cells that are lysed as a result of effective host defense

mechanisms or by the activities of certain antibiotics. Endotoxins generally act in the vicinity of bacterial growth or presence.

Exotoxins are usually secreted by bacteria and act at a site removed from bacterial growth. However, in some cases, exotoxins are only released by lysis of the bacterial cell. Exotoxins are usually proteins, minimally polypeptides, that act enzymatically or through direct action with host cells and stimulate a variety of host responses. Most exotoxins act at tissue sites remote from the original point of bacterial invasion or growth. However, some bacterial exotoxins act at the site of pathogen colonization and may play a role in invasion.

BACTERIAL PROTEIN TOXINS

Exotoxins are usually secreted by living bacteria during exponential growth. The production of the toxin is generally specific to a particular bacterial species that produces the disease associated with the toxin (e.g. only *Clostridium tetani* produces tetanus toxin; only *Corynebacterium diphtheriae* produces the diphtheria toxin). Usually, virulent strains of the bacterium produce the toxin while nonvirulent strains do not, and the toxin is the major determinant of virulence (e.g. tetanus and diphtheria). At one time, it was thought that exotoxin production was limited mainly to Gram-positive bacteria, but clearly both Gram-positive and Gram-negative bacteria produce soluble protein toxins.

- Bacterial protein toxins are the most powerful human poisons known and retain high activity at very high dilutions. The lethality of the most potent bacterial exotoxins is compared to the lethality of strychnine, snake venom, and endotoxin in Table 1 below.

TABLE 1. LETHALITY OF BACTERIAL PROTEIN TOXINS

Toxin	Toxic Dose (mg)	Host	Lethal toxicity compared with:		
			Strychnine	Endotoxin (LPS)	Snake Venom
Botulinum toxin	0.8×10^{-8}	Mouse	3×10^6	3×10^7	3×10^5
Tetanus toxin	4×10^{-8}	Mouse	1×10^6	1×10^7	1×10^5
Shiga toxin	2.3×10^{-6}	Rabbit	1×10^6	1×10^7	1×10^5
Diphtheria toxin	6×10^{-5}	Guinea pig	2×10^3	2×10^4	2×10^2

- Usually the site of damage caused by an exotoxin indicates the location for activity of that toxin. Terms such as **enterotoxin**, **neurotoxin**, **leukocidin** or **hemolysin** are descriptive terms that indicate the target site of some well-defined protein toxins.
- A few bacterial toxins that obviously bring about the death of an animal are known simply as **lethal toxins**, and even though the tissues affected and the target site or substrate may be known, the precise mechanism by which death occurs is not clear (e.g. anthrax LF).
- Some bacterial toxins are utilized as **invasins** because they act locally to promote bacterial invasion. Examples are extracellular enzymes that degrade tissue matrices or fibrin, allowing the bacteria to spread. This includes collagenase, hyaluronidase and streptokinase. Other toxins, also

considered invasins, degrade membrane components, such as phospholipases and lecithinases.

- The pore-forming toxins that insert a pore into eucaryotic membranes are considered as invasins, as well, but they will be reviewed here.

- Some protein toxins have very **specific cytotoxic activity** (i.e., they attack specific types of cells). For example, tetanus and botulinum toxins attack only neurons. But some toxins (as produced by staphylococci, streptococci, clostridia, etc.) have fairly **broad cytotoxic activity** and cause nonspecific death of various types of cells or damage to tissues, eventually resulting in necrosis. Toxins that are phospholipases act in this way. This is also true of pore-forming hemolysins and leukocidins.

- Bacterial protein toxins are strongly **antigenic**. *In vivo*, specific antibody neutralizes the toxicity of these bacterial exotoxins (**antitoxin**). However, *in vitro*, specific antitoxin may not fully inhibit their activity. This suggests that the antigenic determinant of the toxin may be distinct from the active portion of the protein molecule. The degree of neutralization of the active site may depend on the distance from the antigenic site on the molecule. However, since the toxin is fully neutralized *in vivo*, this suggests that other host factors must play a role in toxin neutralization in nature.

- Protein exotoxins are inherently **unstable**. In time they lose their toxic properties but retain their antigenic ones. This was first discovered by Ehrlich who coined the term "toxoid" for this product.

Toxoids are detoxified toxins which retain their antigenicity and their immunizing capacity. The formation of toxoids can be accelerated by treating toxins with a variety of reagents including formalin, iodine, pepsin, ascorbic acid, ketones, etc. The mixture is maintained at 37 degrees at pH range 6 to 9 for several weeks.

The resulting toxoids can be used for **artificial immunization** against diseases caused by pathogens where the primary determinant of bacterial virulence is toxin production. Toxoids are effective immunizing agents against diphtheria and tetanus that are part of the DPT (DTP) vaccine.

Nomenclature of toxins

1– Named for host cell attacked:

- **Neurotoxins** , **Enterotoxins**
- **Cytotoxins** (**Nephrotoxin** ,**Hepatotoxin** , **Cardiotoxin**)

2– Named for producer or disease: cholera, Shiga

3– Named for activity: lecithinase, adenylate cyclase

4-Letter designation: exotoxin A

Toxins with Enzymatic Activity

As proteins, many bacterial toxins **resemble enzymes** in a number of ways. Like enzymes, they are **denatured by heat**, acid and proteolytic enzymes, they **act catalytically**, and they exhibit **specificity of action**.

The **substrate** (in the host) may be a component of tissue cells, organs or body fluid.

A plus B Subunit Arrangement

Many protein toxins, notably those that act intracellularly (with regard to host cells), consist of two components:

- one component (**subunit A**) is responsible for the **enzymatic activity** of the toxin
- the other component (**subunit B**) is concerned with **binding** to a specific receptor on the host cell membrane and transferring the enzyme across the membrane.
- The enzymatic component is not active until it is released from the native (**A+B**) toxin.
- Isolated A subunits are enzymatically active but lack binding and cell entry capability.
- Isolated B subunits may bind to target cells (and even block the binding of the native toxin), but they are nontoxic.

There are a variety of ways that toxin subunits may be synthesized and arranged:

- **A + B** indicates that the toxin is synthesized and secreted as two separate protein subunits that interact at the target cell surface;
- **A-B** or **A-5B** indicates that the A and B subunits are synthesized separately, but associated by noncovalent bonds during secretion and binding to their target
- **5B** indicates that the binding domain of the protein is composed of 5 identical subunits.

- **A/B** denotes a toxin synthesized as a single polypeptide, divided into A and B domains that may be separated by proteolytic cleavage.

Attachments and mechanism of toxin entry

There are at least **two mechanisms of toxin entry into target cells**.



- **direct entry**, the B subunit of the native (A+B) toxin binds to a specific receptor on the target cell and induces the formation of a pore in the membrane through which the A subunit is transferred into the cell cytoplasm.

- an alternative mechanism, the native toxin binds to the target cell and the A+B structure is taken into the cell by the process of **receptor-mediated endocytosis (RME)**. The toxin is internalized in the cell in a membrane-enclosed vesicle called an **endosome**. H^+ ions enter the endosome lowering the internal pH which causes the A+B subunits to separate. The B subunit affects the release of the A subunit from the endosome so that it will reach its target in the cell cytoplasm. The B subunit remains in the endosome and is recycled to the cell surface.

In both cases above, a large protein molecule must insert into and cross a membrane lipid bilayer, either the cell membrane or the endosome membrane. This activity is reflected in the ability of most A+B or A/B toxins, or their B components, to insert into artificial lipid bilayers, creating ion permeable pathways.


If the B subunit contains a hydrophobic region (of amino acids) that insert into the membrane (as in the case of the diphtheria toxin), it may be referred to as the T (translocation) domain of the toxin.

Binding of A-B toxins




The B (binding) component of the exotoxin binds to a receptor on the surface of a susceptible host cell.

Entry of A Component of A-B Toxins by Direct Passage through the Host Cell's Membrane



Entry of A-B Toxins by Endocytosis



A few bacterial toxins (e.g. diphtheria) are known to utilize both direct entry and RME to enter into host cells, which is not surprising since both mechanisms are variations on a theme.

Bacterial toxins with similar enzymatic mechanisms may enter their target cells by different mechanisms. Thus, the diphtheria toxin and *Pseudomonas* exotoxin A, which have identical mechanisms of enzymatic activity, enter their host cells in slightly different ways.

The adenylate cyclase toxin of *Bordetella pertussis* (pertussis AC) and anthrax EF produced by *Bacillus anthracis*, act similarly to catalyze the production of cAMP from host cell intracellular ATP reserves. However, the anthrax toxin enters cells by receptor mediated endocytosis, whereas the pertussis adenylate cyclase traverses the cell membrane directly.

The **specific receptors** for the B subunit of toxins on target cells or tissues are usually sialogangliosides (glycoproteins) called **G-proteins** on the cell membrane. For example, the cholera toxin utilizes the ganglioside GM1, and tetanus toxin utilizes ganglioside GT1 and/or GD1b as receptors on host cells.

Reference

- *Microbial toxins : current research and future trends.*
Proft, Thomas. Norfolk: Caister Academic Press. 2009.
- Lectures by Prof. Dr. Mai Talib Falih - University of
Baghdad / college of Science .2011
- علم السموم البكتريا ، الاستاذ الدكتور رحاب العزاوي 2010
- السموم الميكروبية رشا حمد الميهي وجمهورية مصر العربية 2014