

University of Anbar

College of Science

Applied Mathematics

Numerical Analysis

Dr. Hamad Mohammed Salih

University of Anbar

College of Science

Applied Mathematics

Numerical Analysis

Dr. Hamad Mohammed Salih

## Numerical Analysis2

### **First Lecture**

For solving a linear system of n equations in n variable. Such a system has the form

$$\begin{aligned} E_1 : a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &= b_1 \\ E_2 : a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n &= b_2 \\ \cdot & \quad \cdot \quad \quad \quad \cdot \quad \quad \quad \cdot \\ \cdot & \quad \cdot \quad \quad \quad \cdot \quad \quad \quad \cdot \\ \cdot & \quad \cdot \quad \quad \quad \cdot \quad \quad \quad \cdot \\ E_n : a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nn}x_n &= b_n \end{aligned}$$

In this system we are given the constants  $a_{ij}$  for  $i, j = 1, 2, \dots, n$  and  $b_i = 1, 2, \dots, n$  and we need for solving system of this type. Direct methods find the solution in a finite number of steps or iterative methods start with an arbitrary first approximation to x and then improve this estimate in an infinite but convergent sequence of steps.

### **1-Direct methods:**

#### **1-1Gauss elimination method:**

Gauss elimination method is used to solve a system of linear equation by transforming it into an upper triangular system. (i.e. one in which all of the coefficients below the leading diagonal are zero) using elementary row operations. The solution of the upper triangular system. Then found using back substitution. We will describe the method in detail in the example down.

**Example 1:** The four equations

$$\begin{aligned} E_1 : x_1 + x_2 + 0 + 3x_4 &= 4 \\ E_2 : 2x_1 + x_2 - x_3 + x_4 &= 1 \\ E_3 : 3x_1 - x_2 - x_3 + 2x_4 &= -3 \\ E_4 : -x_1 + 2x_2 + 3x_3 - x_4 &= 4 \end{aligned}$$

Will be solved for  $x_1, x_2, x_3$  and  $x_4$ . We first use  $E_1$  to eliminate the unknown  $x_1$  from  $E_2, E_3$  and  $E_4$  by performing  $(E_2 - 2E_1) \rightarrow E_2, (E_3 - 3E_1) \rightarrow E_3$  and  $(E_4 + E_1) \rightarrow E_4$ . For example, in the second equation

$$(E_2 - 2E_1) \rightarrow E_2$$

Which simplifies to the result shown as  $E_2$  in

$$\begin{aligned} E_1 : x_1 + x_2 + 0 + 3x_n &= 4 \\ E_2 : -x_2 - x_3 - 5x_4 &= -7 \\ E_3 : -4x_2 - x_3 - 7x_4 &= -15 \\ E_4 : 3x_2 + 3x_3 + 2x_4 &= 8 \end{aligned}$$

For simplicity, the new equations are again labelled  $E_1, E_2, E_3$  and  $E_4$ . In the new system,

$E_2$  is used to eliminate the unknown  $x_2$  from  $E_3$  and  $E_4$  by performing

$(E_3 - 4E_2) \rightarrow E_3$  and  $(E_4 + 3E_2) \rightarrow E_4$  This results in

$$\begin{aligned} E_1 : x_1 + x_2 + 0 + 3x_n &= 4 \\ E_2 : -x_2 - x_3 - 5x_4 &= -7 \\ E_3 : 3x_3 + 13x_4 &= 13 \\ E_4 : -13x_4 &= -13 \end{aligned} \quad (*)$$

The system (\*) is now in triangular form and can be solved for the unknown by a backward – substitution process. Since  $E_4$  implies  $x_4 = 1$ , we can solve

$$E_3 \text{ for } x_3 \text{ to given } x_3 = \frac{1}{3}(13 - 13x_4) = \frac{1}{3}(13 - 13) = 0$$

Continuing,  $E_2$  given  $x_2 = -(-7 + 5x_4 + x_3) = -(-7 + 5 + 0) = 2$  and  $E_1$  given

$$x_1 = 4 - 3x_4 - x_2 = 4 - 3 - 2 = -1 .$$

The solution to system (\*), and consequently to system (1) is therefor

$$x_1 = -1, x_2 = 2, x_3 = 0 \text{ and } x_4 = 1$$

### **1-1-1 Gauss elimination method with partial pivoting.**

To reduce round-off error, it is often necessary to perform row interchanges even when the pivot element is not zero. If one of the pivot element is  $a_{ij} = 0$ . This row interchange has the form  $E_i \leftrightarrow E_k$ .

pivoting is performed by selecting an element  $a_{ij}$  with a larger magnitude as the pivot, and interchanging the  $k$ th and  $p$ th rows. This can be followed by the interchange of the  $k$ th and  $q$ th columns if necessary.

**Example2:**

Apply Gauss elimination to the system

$$E_1 : 0.003000x_1 + 59.14x_2 = 59.17$$

$$E_2 : 5.29x_1 - 6.130x_2 = 46.78$$

Using partial pivoting and four-digit arithmetic with rounding , and compare the results to the exact solution

$$x_1 = 10.00 \text{ and } x_2 = 1.$$

**Solution**

The partial –pivoting procedure first requires finding max

$$\{|a_{11}^{(1)}|, |a_{21}^{(1)}|\} = \max \{|0.00300|, |5.291|\} = \{|a_{21}^{(1)}|\}$$

This requires that the operation  $(E_2) \leftrightarrow (E_1)$  be performed to produce the equivalent system

$$E_1 : 5.29x_1 - 6.130x_2 = 46.78$$

$$E_2 : 0.003000x_1 + 59.14x_2 = 59.17$$

The multiplier for this system is

$$m_{21} = \frac{a_{21}^{(1)}}{a_{11}^{(1)}} = 0.0005670$$

And the operation  $(E_2 - m_{21}E_1) \rightarrow E_2$  reduces the system to

$$E_1 : 5.29x_1 - 6.130x_2 = 46.78$$

$$E_2 : \quad 0 \quad + 59.14x_2 = 59.17$$

## Homework

### Example 3

Use the G.E. method to solve the following linear system , if possible, and determined whether row interchanges are necessary.

(Exact solution

a)  $x_1 = 1.1875, x_2 = 1.8125$  and  $x_3 = 0.875$  with one row interchange required

b)  $x_1 = -1, x_2 = 0$  and  $x_3 = 1$  with no interchange required )

$$\begin{array}{ll} E_1 : x_1 - x_2 + 3x_3 = 2 & E_1 : 2x_1 - 1.5x_2 + 3x_3 = 1 \\ \text{a) } E_2 : 3x_1 - 3x_2 + x_3 = -1 & \text{b) } E_2 : -x_1 + 2x_3 = 3 \\ E_3 : x_1 + x_2 = 3 & E_3 : 4x_1 - 4.5x_2 + 5x_3 = 1 \end{array}$$

$$2x_1 - 1.5x_2 + 3x_3 = 1$$

$$-x_1 + 2x_3 = 3$$

$$4x_1 - 4.5x_2 + 5x_3 = 1$$

$$\begin{bmatrix} 2 & -1 & 3 \\ -1 & 0 & 2 \\ 4 & -4 & 5 \end{bmatrix} \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} \xrightarrow[R_1+2R_2]{2R_1-R_3} \begin{bmatrix} 2 & -1 & 3 \\ 0 & -2 & 7 \\ 0 & -6 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 7 \\ 1 \end{bmatrix} \xrightarrow{3R_2-R_3} \begin{bmatrix} 2 & -1 & 3 \\ 0 & -2 & 7 \\ 0 & 0 & 20 \end{bmatrix} \begin{bmatrix} 1 \\ 7 \\ 20 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -1 & 3 \\ A_1 & -2 & 7 \\ A_1 & A_3 & 20 \end{bmatrix} \begin{bmatrix} 1 \\ 7 \\ 20 \end{bmatrix} = \begin{bmatrix} 2 & -1 & 3 \\ 0 & A_1 & A_2 \\ 0 & 0 & A_3 \end{bmatrix} \begin{bmatrix} 1 \\ 7 \\ 20 \end{bmatrix}$$

$$A_1 = -2, A_2 = 7, A_3 = 20$$

### Example 4

Use the G.E. method with backward substitution and two-digit rounding arithmetic to solve the following linear system . Don't reader the equation

(Exact solution

a)  $x_1 = 1, x_2 = -1$  and  $x_3 = 3$

$$E_1 : 4x_1 - x_2 + 3x_3 = 8$$

a)  $E_2 : 2x_1 + 5x_2 + 2x_3 = 3$

$$E_3 : x_1 + 2x_2 + 4x_3 = 11$$

$$E_1 : 4x_1 + x_2 + 2x_3 = 9$$

b)  $E_2 : 2x_1 + 4x_2 - x_3 = -5$

$$E_3 : x_1 + x_2 - 3x_3 = -9$$

Second Lecture

### **2-Gauss Jordan method:**

The general idea of this method is to change the coefficients down the diagonal to zero (called upper-triangular form), as well as change the coefficients above the diagonal to zero (called Lower-triangular form), by the same steps used in the G.E. method. At last, we obtain the solution of system without using process backward substitution.

### **Example 5**

Use Gauss-Jordan method and two –digit chopping arithmetic to solve the linear system.

$$2x_1 + 3x_2 - x_3 = 5$$

$$4x_1 + 4x_2 - 3x_3 = 3$$

$$-2x_1 + 3x_2 - x_3 = 1$$

**Solution**



$$\begin{aligned}
 [A:B] &= \left[ \begin{array}{ccc|c} 2 & 3 & -1 & 5 \\ 4 & 4 & -3 & 3 \\ -2 & 3 & -1 & 1 \end{array} \right] \xrightarrow{-2R_1+R_2} \xrightarrow{R_1+R_3} \left[ \begin{array}{ccc|c} 2 & 3 & -1 & 5 \\ 0 & -2 & -1 & -7 \\ 0 & 6 & -2 & 6 \end{array} \right] \\
 &\xrightarrow{3r_2+r_3} \left[ \begin{array}{ccc|c} 4 & 0 & -5 & -11 \\ 0 & -2 & -1 & -7 \\ 0 & 0 & -5 & -15 \end{array} \right] \xrightarrow{-R_3+R_1} \xrightarrow{-5R_2+R_3} \left[ \begin{array}{ccc|c} 4 & 0 & 0 & 4 \\ 0 & 10 & 0 & 20 \\ 0 & 0 & -5 & -15 \end{array} \right]
 \end{aligned}$$

$$\Rightarrow -5x_3 = -15 \rightarrow x_3 = 3, x_2 = 2, x_1 = 1$$

### **3- Inverse Matrices method**

Certain  $n \times n$  matrices have the property that another  $n \times n$  matrix, which we will denote  $A^{-1}$

Exists with  $AA^{-1} = A^{-1}A = I$ . In this case  $A$  is said to be non-singular, or invertible, and the matrix  $A^{-1}$  is called the inverse to  $A$ . A matrix without an inverse is called singular or noninvertible.

#### **Example 6:**

Find the inverse of the following system

$$\begin{aligned}
 x_1 - x_2 + x_3 &= -4 \\
 5x_1 - 4x_2 + 3x_3 &= -12 \\
 2x_1 + x_2 + x_3 &= 11
 \end{aligned}$$

**Solution**

$$\begin{aligned}
& \begin{bmatrix} 1 & -1 & 1 & 1 & 0 & 0 \\ 5 & -4 & 3 & | & 0 & 1 & 0 \\ 2 & 1 & 1 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{-5R_1+R_2} \xrightarrow{-2R_1+R_3} \begin{bmatrix} 1 & -1 & 1 & 1 & 0 & 0 \\ 0 & 1 & -2 & | & -5 & 1 & 0 \\ 0 & 3 & -1 & -2 & 0 & 1 \end{bmatrix} \\
& \xrightarrow{-3R_2+R_3} \begin{bmatrix} 1 & -1 & 1 & 1 & 0 & 0 \\ 0 & 1 & -2 & | & -5 & 1 & 0 \\ 0 & 0 & 5 & 13 & -3 & 1 \end{bmatrix} \xrightarrow{-5R_1+R_3} \xrightarrow{2R_3+5R_2} \begin{bmatrix} -5 & 5 & 0 & 8 & -3 & 1 \\ 0 & 5 & 0 & | & 1 & -1 & 2 \\ 0 & 0 & 5 & 13 & -3 & 1 \end{bmatrix} \\
& \xrightarrow{-R_1+R_2} \begin{bmatrix} 5 & 0 & 0 & -7 & 2 & 1 \\ 0 & 5 & 0 & | & 1 & -1 & 2 \\ 0 & 0 & 5 & 13 & -3 & 1 \end{bmatrix} \xrightarrow{+\frac{1}{5}} \begin{bmatrix} 1 & 0 & 0 & -1.4 & 0.4 & 0.2 \\ 0 & 1 & 0 & | & 0.2 & -0.2 & 0.4 \\ 0 & 0 & 1 & 2.6 & 0.6 & 0.2 \end{bmatrix}
\end{aligned}$$

### Reference

- 1-Numerical Analysis. Richara L. Burden, and J. Douglas Faires .Ninth Edition.
- 2- Numerical Methods. J. Douglas Faires and Richara L. Burden. Fourth Edition.
- 3- Numerical mathematics and Computing. Ward Cheney and David Kincaid. Second Edition.